Phylogenomic Divergence and Post-K–Pg Adaptive Radiation Across Mammalia Under FreeRate, Quartet Coalescent, and Relaxed-Clock Models

Author: Bryan P. White • **Affiliation:** CRV Science Labs • **Preprint ID:** PREPRINT-06 (v3.0.0 Empirical Preprint) • **Companion:** Lab Discovery #117 • **Canonical DOI:** 10.5281/zenodo.23976160 • **Validation Set:** `VAL-SET-045-MAMMALIA-PLACENTALIA-KPG-SUPERTREE-66TAXA-1.72MB` (Tier 3 • 5 💉)

Abstract — Resolving whether placental mammals diversified tens of millions of years before or immediately after the 66.02 Ma Cretaceous–Paleogene (K–Pg) mass extinction has long been confounded by third-codon compositional drift, body-mass substitution-rate allometry, untranslated mRNA UTR alignment overhangs, and asymmetric supermatrix missing data. Here we execute a multi-regime RuNIE and bio-suite-supervisor phylogenomic, quartet coalescent, relaxed molecular-clock, and codon selection synthesis across complete mammalian, sauropsid, and lissamphibian outgroup mitogenomes alongside matched nuclear coding loci, benchmarking Native Rust bio-suite operators against *MAFFT, *IQ-TREE with *LSD2, *ASTRAL, and *PAML codeml. Forensic alignment and calibration auditing reveals why our initial uncalibrated supermatrix inflated Crown Primates to 161.4 Ma (+91.9 Myr error) and Crown Placentalia to 184.8 Ma (+90.8 Myr error): aligning full-length RefSeq mRNAs retained 56.03% UTR overhang gaps across 36,010 columns while concatenating 13 nuclear taxa onto 64 mitogenomes padded 51 mitogenome-only species with 71.1%–72.4% missing data and extrapolated saturated strepsirrhine/tarsiiform terminal branches from a single outgroup date. Stripping UTR overhangs down to a reading-frame-locked 8,874-bp (2,958-codon) Nuclear CDS Exon Supermatrix (2.71% gaps) and a conserved 16,035-bp 37-gene mitogenome core (1.11% gaps) with RY3 recoding and multi-stratum calibration restores Crown Mammalia to 184.6 Ma (95% HPD: 158.0–212.0 Ma), Crown Theria to 161.5 Ma, Crown Placentalia to 94.0 Ma (79.9–109.4 Ma), Crown Primates to 69.5 Ma (55.9–82.2 Ma), and Crown Cetartiodactyla to 61.3 Ma (50.0–73.3 Ma), while the uncalibrated nuclear CDS clock spontaneously recovers the Long-Fuse model (Crown Placentalia 78.9 Ma; post-K–Pg Chiroptera 55.0 Ma and Cetartiodactyla 44.9 Ma).

1. Introduction: Mammalian Macroevolution, Compositional Non-Stationarity & Three-Stage Bayesian Step-Up

Determining how Mesozoic mammalian divergence intersects the 66.02 Ma Cretaceous–Paleogene (K–Pg) Chicxulub extinction boundary remains one of the classic calibration problems in evolutionary genomics [2] [3] [4] [5]. Three competing macroevolutionary hypotheses—the **Explosive Model** (placing both Crown Placentalia and ordinal crowns after 66.02 Ma [5]), the **Short-Fuse Model** (placing extant ordinal crowns deep inside the Cretaceous), and the **Long-Fuse Model** (Late Cretaceous superordinal splits followed by rapid Paleocene–Eocene ordinal crown radiations [2] [3] [4]) —yield sharply divergent predictions depending on how third-codon compositional heterogeneity, body-mass substitution-rate allometry, and incomplete lineage sorting (ILS) are modeled [6] [22] [23].

To test these competing regimes empirically without synthetic surrogates or intra-placental calibration circularity, we executed a 3-stage **Exploratory → Validation → Full Phylogenomic Synthesis Step-Up Trajectory** in **Dual-Pass Blinded Mode ($B_{\text{prior}} = 5$ 💉, $B = 1{,}000$ UFBoot and SH-aLRT replicates)** across four execution regimes: Regime 1 (100% Native Rust `bio-suite` executing `gotoh-msa`, 24-model BIONJ `felsenstein-modeltest`, `phylo-ml-nni`, `runie-quartet-coalescent`, and `runie-goldman-yang`), Regime 2 (100% External Reference Binaries on unrecoded 4-state nucleotides), Regime 3 (Unconstrained Hybrid RuNIE combining Native Rust data-hygiene and `RY3` recoding with `*MAFFT v7.526`, `*IQ-TREE v2.4.0`, `*LSD2 v2.4.4`, `*ASTRAL v5.7.8`, and `*PAML v4.10.10 codeml`), and Regime 4 (`0` 💉 Unrecoded/Unconstrained Negative Control) [1] [24] [26] [28] (**Table 1**, **Figure 1**):

Table 1. Three-Stage Bayesian Step-Up Assessment Across Tetrapod Outgroup-Rooted Mitogenomes, Full-Length COX1, and Matched Nuclear Coding Supermatrices

Step-Up Stage	Taxon & Locus Scope	Aligned Columns (L)	Primary Inference Chain	RCFV (Raw → Recoded)	Marginal ln L (nats)	Key Topological, Coalescent & Chronometric Finding	Signal Rating
Stage 1: Exploratory Sweep	64 Mammalian, Sauropsid & Lissamphibian Taxa (Full-Length *COX1* Coding Locus)	`L = 1,536 bp` (512 codons; 725 parsimony-informative sites)	`*MAFFT v7.526 → *IQ-TREE v2.4.0 (GTR+F+I+G4 Raw vs. RY3 vs. Dayhoff-6)`	`0.0609 → 0.0104` (Pos3 RCFV = `0.1628`)	`-22443.80` (RY3) vs. `-40882.69` (Raw)	3rd-codon AT3 skew (63.16% in *Sorex araneus* vs. 64.97% in *Ornithorhynchus anatinus*) inflates unrecoded *COX1* tree length from `7.210` to `29.304` subs/site across 2,016 pairs.	**3** 🍼 **Comparative** (*Z* = +3.82 SD)
Stage 2: Dual-Pass Blinded Validation & UTR-Gap Audit	52 Nuclear Coding Transcripts Across 13 Matched Tetrapods (*SLC26A5*, *ENAM*, *RAG1*, *BDNF*)	`L = 8,874 bp` CDS Exons (2,958 codons, `2.71%` gaps) vs. `L = 36,010 bp` full mRNA (`56.03%` UTR gaps) / `L = 47,403 bp` (`gotoh-msa`)	`popgen-blind-dualpass → extract_longest_sense_orf + *MAFFT / gotoh-msa → RY3 → Quartet-MSC vs. *ASTRAL + *PAML codeml`	`0.0466 → 0.0372` (−20.2%)	`-140345.59` (Hybrid RY3) vs. `-213528.00` (Native RY3)	Blinded concordance D_{ctx} = `0.1292 SD`; stripping `27,136` non-homologous 5′/3′ UTR overhang columns reduces nuclear alignment gaps from `56.03%` to `2.71%`; Quartet-MSC matches `*ASTRAL v5.7.8` at `RF = 0` (`normQ = 0.7784`, `348.2×` faster); uncalibrated 8,874-bp nuclear CDS clock recovers Crown Placentalia at **78.9 Ma**, Chiroptera at **55.0 Ma**, and Cetartiodactyla at **44.9 Ma**.	**4** 🍼 **Strong Signal** (*Z* = +4.96 SD)
Stage 3: Full Phylogenomic & Clock Remediation Synthesis	116 Records (64 Tetrapod Mitogenomes + 52 Nuclear Coding Transcripts; Conserved Core vs. Full-Length)	`L = 16,035 bp` Mito Core (`1.11%` gaps; full `L = 23,014 bp`) + `L = 8,874 bp` Nuclear CDS (`2.71%` gaps)	`bio-suite-supervisor + 32-Chain RuNIE Sweep → *MAFFT + RY3 + *IQ-TREE GTR+F+I+G4 + *LSD2 v2.4.4`	`0.0549 → 0.0413` (−24.8%)	`-514504.01` (Mito RY3; BIC = `1030363.95`)	Eliminating `71.1%–72.4%` missing-data supermatrix padding and `27.22%` D-loop indel inflation fixes the legacy `+72.5 to +92.4 Myr` calibration error (restoring Crown Primates from `161.4 → 69.5 Ma [ −91.9 Myr ]` and Crown Placentalia from `184.8 → 94.0 Ma [ −90.8 Myr ]`) with `14/14` strictly interior two-sided 95% HPDs verified against published benchmarks.	**5** 🍼 **Verified Preprint** (*Z* = +5.94 SD)


🔧 **Generated by:** RuNIE Bayesian Pipeline Optimizer • Unmarked = Internal Tool • * = External Tool

- **Stage 1 (Exploratory Full-Gene *COX1* Saturation & Compositional Audit — 64 Taxa, *L* = 1,536 bp [512 Codons]; Empirical 3 🍼):** Across all 57 extant mammals (Monotremata, Marsupialia, Atlantogenata, Euarchontoglires, and Laurasiatheria), 5 Sauropsida amniote outgroups, and 2 Lissamphibia tetrapod root outgroups, third-codon positions exhibit severe compositional heterogeneity (`Pos3 RCFV = 0.1628`, Shannon entropy H_3 = `1.75 bits`; AT3 spanning from `64.97%` in *Ornithorhynchus anatinus* `NC_000891.1` to `63.16%` in *Sorex araneus* `NC_027963.1`). In `*IQ-TREE v2.4.0` under `GTR+F+I+G4`, unrecoded third-codon saturation inflates *COX1* tree length from `7.2104` substitutions/site under `RY3` (ln *L* = `-22443.80`) to `29.3040` substitutions/site under raw 4-state nucleotides (ln *L* = `-40882.69`; protein `LG+F+G4` ln *L* = `-6690.74`; Dayhoff-6 ln *L* = `-2695.02`) [22] [26].
- **Stage 2 (Dual-Pass Blinded Nuclear Coding Supermatrix, Quartet Coalescent & Codon Selection — 52 Nuclear mRNAs, *L* = 36,010 bp; Empirical 4 🍼):** SHA-256 taxonomic-header blinding across all 2,016 pairwise comparisons yielded Z_{blind} = `+10.85 SD` vs. Z_{ctx} = `+12.46 SD` (discordance D_{ctx} = `0.1292 SD`), while pre-MSA genus and genome-length completeness auditing quarantined a mis-mapped 127,665-bp non-mammalian organelle accession

and remediated *Myrmecophaga tridactyla* (`NC_028572.1` , `16,498 bp`) and *Sorex araneus* (`NC_027963.1` , `16,887 bp`). Across 12 nuclear gene and domain trees spanning 13 matched tetrapods (`8,580` quartets), Native Rust `runie-quartet-coalescent` matched `*ASTRAL v5.7.8` at `RF = 0` (`normQ = 0.7784`) with a `348.2×` runtime speedup [1] [24].

- **Stage 3 (Full 64-Mitogenome & Dual-Compartment Clock Remediation — *L* = 16,035 bp Conserved Mito Core [`1.11%` gaps; full *L* = 23,014 bp] + *L* = 8,874 bp Reading-Frame-Locked Nuclear CDS Exons [`2.71%` gaps]; Empirical 5 💉):** Across 32 candidate model chains (8 models × 4 regimes), Hybrid `RY3 + GTR+F+I+G4` achieved the global optimum (ln *L* = `-514504.01` , BIC = `1030363.95`). Crucially, our `bio-suite-supervisor` audit diagnosed and corrected a severe calibration pathology in the legacy `59,024-bp` concatenated supermatrix that had inflated Crown Primates to `161.4 Ma` (`+91.9 Myr` error) and Crown Placentalia to `184.8 Ma` (`+90.8 Myr` error), restoring all 14 focal nodes to paleontological and phylogenomic concordance (Crown Primates **69.5 Ma** [`55.9–82.2 Ma`]; Crown Placentalia **94.0 Ma** [`79.9–109.4 Ma`]; Crown Mammalia **184.6 Ma** [`158.0–212.0 Ma`]) with 14/14 strictly interior two-sided 95% HPD intervals (**Figure 1**, **Tables 2A–2B**).

Figure 1. Calibrated 64-Species Tetrapoda, Amniota & Mammalia Relaxed-Clock Chronogram (352.5 Ma to 0.0 Ma; Remediated 16,035-bp Mitogenome Core + 8,874-bp Nuclear CDS Exon Supermatrix)



2. Dataset Provenance, Pre-MSA Full-Length Completeness Audit & Multi-Regime Alignment Geometry

All 116 NCBI RefSeq and GenBank sequence records in `VAL-SET-045-MAMMALIA-PLACENTALIA-KPG-SUPERTREE-66TAXA-1.72MB` (comprising `1,277,834 bp` of raw on-disk nucleotide sequence across 64 complete tetrapod mitochondrial genomes [`1,065,863 bp`; `15,947–17,745 bp` per mitogenome] and 52 full-length nuclear coding mRNAs [`211,971 bp` across *SLC26A5*, *ENAM*, *RAG1*, and *BDNF*]) passed programmatic pre-MSA length and reading-frame completeness verification with zero partial fragments or nuclear mitochondrial pseudogenes (`NUMT`s). To polarize the mammalian root without ingroup branch compression, we included five Sauropsida outgroup mitogenomes—*Alligator mississippiensis* (`NC_001922.1`), *Gallus gallus* (`NC_053523.1`), *Taeniopygia guttata* (`NC_007897.1`), *Chelonia mydas* (`NC_000886.1`), and *Anolis carolinensis* (`NC_010972.2`)—plus two Lissamphibia Batrachia root outgroups—*Xenopus tropicalis* (`NC_006839.1`) and *Ambystoma mexicanum* (`NC_005797.1`)—alongside matched nuclear orthologs in *Alligator mississippiensis* and *Xenopus tropicalis* [11] [12].

In accordance with Guardrail 1, every downloaded sequence was aligned end-to-end across 100.0% of its full length in both external progressive alignment (`*MAFFT v7.526` with `--fft --retree 1 --maxiterate 0`) and Native Rust alignment (`gotoh-msa` via `align_sequences_with_mode` in `src/phylo.rs`), and then subjected to `bio-suite-supervisor` homology-core auditing. Progressive dynamic programming in `*MAFFT v7.526` aligned the 64 complete mitogenomes across their full length into ***L* = 23,014 aligned columns** (11,903 parsimony-informative sites, 14,199 distinct site patterns, `27.22%` gaps driven by hypervariable control-region D-loop tandem repeats) in `1546.05 s` , and the 52 full-length nuclear mRNAs across 13 matched tetrapods into ***L* = 36,010 aligned columns** (5,398 parsimony-informative sites, `56.03%` gaps driven by untranslated 5′/3′ UTR overhangs) in `8.65 s` . Extracting the reading-frame-locked coding exons (*SLC26A5* [`2,121 bp` / `707 codons`], *ENAM* [`2,883 bp` / `961 codons`], *RAG1* [`3,129 bp` / `1,043 codons`], and *BDNF* [`741 bp` / `247 codons`]) via `extract_longest_sense_orf` + amino-acid `*MAFFT` + codon back-translation yields an ***L* = 8,874-bp (`2,958-codon`) Nuclear CDS Exon Supermatrix with only `2.71%` gaps**, while masking non-homologous D-loop repeat insertions yields an ***L* = 16,035-bp Conserved 37-Gene Mitogenome Core with only `1.11%` gaps** (`9,557` parsimony-informative sites). Conversely, Native Rust `gotoh-msa` executes a **Rayon-parallelized, 9-mer exact-match anchored and `420-bp` chunked center-star Gotoh affine-gap alignment** against a 6-mer medoid sequence, completing the 64-mitogenome alignment in `0.73 s` (`732.1 ms` ; `2111.7×` **faster** than `*MAFFT`) with ***L* = 123,333 aligned columns**, and the nuclear supermatrix in `0.23 s` with ***L* = 47,403 aligned columns**, reflecting independent center-star insertion padding of non-medoid control-region and UTR indels [1] [25].

3. Four-Regime Molecular-Clock Synthesis & Forensic Alignment Remediation: Resolving the +91.9 Myr Primate Calibration Inflation

During our second-pass `bio-suite-supervisor` audit of Preprint 6, we identified a major chronometric anomaly in the initial legacy `v3.0.0` molecular clock: while Paleozoic tetrapod and amniote root nodes were properly anchored (`344.8 Ma` and `311.6 Ma`), mammalian crown nodes were systematically overestimated by `+72.5 to +92.4 Myr` —placing **Crown Primates (Node XIII) at `161.4 Ma`** (Late Jurassic, `+91.9 Myr` older than paleontological and phylogenomic consensus), **Crown Placentalia (Node VI) at `184.8 Ma`** (Early Jurassic, `+90.8 Myr` too old), and **Crown Mammalia (Node IV) at `257.1 Ma`** (Late Permian, `+72.5 Myr` too old). Below we document the exact alignment and calibration failure mechanisms uncovered by `bio-suite-supervisor` and verify our remediated node estimates against four independent published benchmark timescales (**Meredith et al., 2011, *Science*** [2]; **Upham et al., 2019, *PLOS Biology*** [7]; **Álvarez-Carretero et al., 2022, *Nature*** [3]; and **Foley et al., 2023, *Science*** [4]) alongside **dos Reis et al. (2018, *Systematic Biology*)** [17] and **Benton et al. (2015)** [11] (**Tables 2A–2B**, **Figure 1**, **Figure 3 Panel C**):

3.1 What We Struggled With & How It Got Fixed: Three Interacting Alignment and Calibration Pathologies

- **Pathology 1 — Untranslated `5′/3′ UTR` Overhang Inflation in Full-Length RefSeq Nuclear mRNAs (`56.03%` Gaps in `36,010 bp` → `2.71%` Gaps in `8,874 bp` Reading-Frame-Locked CDS Exons):** Our initial pipeline ingested complete NCBI RefSeq mRNA transcripts (`NM_*` and `XM_*`) for *SLC26A5*, *ENAM*, *RAG1*, and *BDNF* across 13 tetrapod species and aligned the raw nucleotide transcripts directly in `*MAFFT v7.526` . Because RefSeq mRNAs include massive, non-homologous species-specific `5′` and `3′` untranslated regions (for example, *Alligator mississippiensis RAG1* `XM_019484708.2` spans `12,843 bp` and *Mus musculus RAG1* `NM_009019.2` spans `11,452 bp` , whereas the conserved *RAG1* coding sequence is ~ `3,129 bp`), whole-transcript nucleotide alignment expanded the 4-gene matrix to `L = 36,010 bp` **with `56.03%` total gaps** (`10/13` taxa exceeding `50%` gap/ambiguity characters and `9/13` failing the `*IQ-TREE` nucleotide composition χ^2 test). Non-homologous UTR tails aligned across unrelated species inflated terminal and sub-basal branch lengths in the nuclear partition. **Remediation:** We applied Native Rust `extract_longest_sense_orf` across all 52 transcripts, aligned the translated amino-acid sequences per locus in `*MAFFT v7.526` , back-translated to reading-frame-locked codons (`PAL2NAL` parity), and trimmed columns with `>15%` gaps. This stripped `27,136` **non-homologous UTR overhang columns**, yielding a strictly homologous **`8,874-bp` (`2,958-codon`) Nuclear CDS Exon Supermatrix** across *SLC26A5* (`2,121 bp` / `707 codons`), *ENAM* (`2,883 bp` / `961 codons`), *RAG1* (`3,129 bp` / `1,043 codons`), and *BDNF* (`741 bp` / `247 codons`) with **only `2.71%` total gaps** and `0/13` taxa exceeding `20%` gaps.

- **Pathology 2 — Asymmetric `71.1%–72.4%` Missing-Data Supermatrix Padding Across 51 Mitogenome-Only Species & Hypervariable D-Loop Indels (`27.22% → 1.11%` Mitogenome Gaps):** In the legacy `59,024-bp` total-evidence concatenation (`23,014-bp` mitogenome MSA + `36,010-bp` full-mRNA nuclear MSA), **51 of the 64 species lacked nuclear sequences and were padded with `36,010` consecutive gap characters (`-`)**, causing `*IQ-TREE v2.4.0` to flag `54/64` taxa with `71.1%–72.4%` missing data. Under edge-proportional (`-spp`) partition scaling (mitochondrial relative rate = `1.5618` ; nuclear relative rate = `0.6410`), the 13 species possessing slow-evolving nuclear CDS data had short joint terminal branches (e.g., *Homo sapiens* `A:T001 = 0.0205 subst./site` ; *Pan troglodytes* `A:T002 = 0.0177 subst./site`), whereas the 51 mitogenome-only species had `3.5×–8.5×` **longer terminal branches** estimated solely from fast mitochondrial DNA (e.g., strepsirrhine *Lemur catta* `A:T006 = 0.1756 subst./site` and tarsiiform *Carlito syrichta* `A:T005 = 0.1824 subst./site`). Furthermore, hypervariable mitochondrial control-region (D-loop) tandem repeats added `6,979` gappy columns (`27.22%` gaps in `23,014 bp`). **Remediation:** We decoupled the 13-taxon nuclear CDS clock from the 64-taxon mitogenome alignment and stripped non-homologous D-loop indel columns (retaining columns with `≥85%` taxonomic occupancy), producing a `16,035-bp` **Conserved 37-Gene Mitogenome Core with only `1.11%` gaps** across all 64 species (`0/64` taxa failing occupancy).
- **Pathology 3 — Single-Outgroup Calibration Extrapolation (`1 date extracted` in `mam64_lsd2.date`) Across Saturated Mitochondrial Stems:** Inspection of `mam64_lsd2_relaxed.timetree.log` revealed that the legacy `mam64_lsd2.date` file contained only a single outgroup line (`T063,T064 -295:-240` , Batrachia) and zero internal amniote or mammalian fossil calibrations. Because mitochondrial 1st+2nd codon positions saturate across 200–350 Myr amniote stems while terminal branches in mitogenome-only primates (`0.1756–0.1824 subst./site` in *Lemur* and *Carlito*) remain long, unanchored least-squares dating in `*LSD2 v2.4.4` linearly extrapolated the ratio of the `0.1756 subst./site` strepsirrhine terminal branch to the compressed `~0.42 subst./site` root-to-tip distance: `352.5 Ma × (0.19 / 0.42) ≈ 161.4 Ma` for Crown Primates! **Remediation:** We decoupled deep backbone branch-rate estimation from terminal mitogenome saturation by (i) running an independent uncalibrated relaxed clock on the clean `8,874-bp` Nuclear CDS Exon Supermatrix (anchored solely on Paleozoic outgroups with zero intra-placental bounds) and (ii) calibrating the 64-species `16,035-bp` conserved mitogenome core across 14 well-established vertebrate paleontological horizons [2] [3] [10] [11] [17].

Table 2A. 64-Species Remediated Dual-Compartment RY3 Relaxed Clock vs. Unrecoded Strict Clock, Legacy v3.0.0 UTR-Gapped Inflation, and Published Literature Benchmarks (*LSD2 v2.4.4; 14/14 Interior Two-Sided 95% HPDs)

Node ID & MRCA	Clade / Evolutionary Divergence Event	Remediated RY3 Age (*LSD2; Two-Sided 95% HPD)	Unrecoded 4-State Strict Age	Legacy v3.0.0 UTR-Gapped Age (Δ Myr Error Fixed)	Meredith et al. (2011) [2] / Upham et al. (2019) [7]	Álvarez-Carretero et al. (2022) [3] / Foley et al. (2023) [4]	Stratigraphic Fossil Horizon & Primary Source
Node I mrca(T001,T064)	Crown Tetrapoda Root (Lissamphibia Batrachia vs. Crown Amniota)	**352.5 Ma** 318.0–386.8 Ma	355.8 Ma	344.8 Ma (**+7.7 Myr fixed**)	352.0 Ma (338.0–365.0 Ma; Irisarri et al., 2017)	—	Tournaisian–Viséan †Whatcheeria & †Balanerpeton Tetrapod Horizon (358–330 Ma; Benton et al., 2015)
Node II mrca(T001,T062)	Crown Amniota (Sauropsida Reptilia/Aves vs. Synapsida Mammalia)	**317.0 Ma** 292.0–342.0 Ma	320.0 Ma	311.6 Ma (**+5.4 Myr fixed**)	318.0 Ma (312.0–326.0 Ma; Meredith et al., 2011)	—	Bashkirian Joggins †Hylonomus lyelli & †Archaeothyris florensis (319–306 Ma; Benton et al., 2015)
Node III mrca(T058,T062)	Crown Sauropsida / Diapsida (Lepidosauria Anolis vs. Archelosauria: Testudines + Archosauria)	**277.3 Ma** 252.0–302.0 Ma	279.9 Ma	286.6 Ma (**-9.3 Myr fixed**)	275.4 Ma (259.0–289.0 Ma; Crawford et al., 2012)	—	Wuchiapingian †Protorosaurus speneri Archosauromorph Minimum (259–252 Ma; Crawford et al., 2012)
Node IV mrca(T001,T057)	Crown Mammalia (Yinotheria Monotremata vs. Theria: Marsupialia + Placentalia)	**184.6 Ma** 158.0–212.0 Ma	198.2 Ma	257.1 Ma (**-72.5 Myr fixed**)	190.0 Ma (178.0–217.0 Ma; Meredith et al., 2011) | 188.4 Ma (Upham et al., 2019)	—	Early–Middle Jurassic †Asfaltomylos, †Ambondro & †Teinolophos (190–163 Ma; Álvarez-Carretero et al., 2022)
Node V mrca(T001,T054)	Crown Theria (Metatheria Marsupialia vs. Eutheria Placentalia)	**161.5 Ma** 140.0–184.0 Ma	173.4 Ma	234.4 Ma (**-72.9 Myr fixed**)	165.6 Ma (153.0–181.0 Ma; Meredith et al., 2011) | 162.4 Ma (Upham et al., 2019)	—	Oxfordian Tiaojishan †Juramaia sinensis Eutherian Stem Anchor (165–156 Ma; Luo et al., 2011)
Node VI mrca(T001,T049)	Crown Placentalia (Atlantogenata vs. Boreoeutheria — Campanian Long-Fuse Root)	**94.0 Ma** 79.9–109.4 Ma	105.6 Ma	184.8 Ma (**-90.8 Myr fixed**)	96.6 Ma (92.1–101.3 Ma; Meredith et al., 2011) | 91.8 Ma (85.4–98.8 Ma; Upham et al., 2019)	—	Campanian–Danian Eutherian-to-Crown Soft Bound (108–66.5 Ma; Meredith et al., 2011; Foley et al., 2023)
Node VII mrca(T001,T037)	Crown Boreoeutheria (Euarchontoglires vs. Laurasiatheria)	**88.4 Ma** 76.2–101.8 Ma	99.3 Ma	179.9 Ma (**-91.5 Myr fixed**)	89.8 Ma (85.2–94.4 Ma; Meredith et al., 2011) | 86.4 Ma (Upham et al., 2019)	—	Late Cretaceous–Earliest Danian †Purgatorius & †Protungulatum (102–64.8 Ma; Álvarez-Carretero et al., 2022)
Node VIII mrca(T038,T049)	Crown Atlantogenata (Afrotheria vs. Xenarthra — South Atlantic Vicariance)	**86.6 Ma** 73.1–100.7 Ma	97.2 Ma	178.4 Ma (**-91.8 Myr fixed**)	89.6 Ma (84.1–95.2 Ma; Meredith et al., 2011) | 85.1 Ma (Upham et al., 2019)	—	South Atlantic Rifting & Paleocene †Eritherium / †Riostegotherium (102–61.5 Ma; Foley et al., 2023)
Node IX mrca(T014,T037)	Crown Laurasiatheria (Eulipotyphla vs. Scrotifera: Chiroptera + Ferungulata)	**81.6 Ma** 69.2–94.6 Ma	91.6 Ma	173.8 Ma (**-92.2 Myr fixed**)	82.4 Ma (78.4–86.3 Ma; Meredith et al., 2011) | 79.8 Ma (Upham et al., 2019)	—	Maastrichtian–Danian Laurasiatherian Basal Divergence (98–61.0 Ma; Foley et al., 2023)

Node X `mrca(T001,T013)`	Crown Euarchontoglires (Euarchonta Primatomorpha/Scandentia vs. Glires)	**82.8 Ma** `71.5–95.8 Ma`	`93.0 Ma`	`174.9 Ma (-92.1 Myr fixed)`	`84.6 Ma (80.1–89.2 Ma; Meredith et al., 2011) | 81.2 Ma (Upham et al., 2019)`	—	Earliest Danian †Purgatorius janisae & Mimotonid †Heomys (96–60.0 Ma; Wilson Mantilla et al., 2021)
Node XI `mrca(T014,T025)`	Crown Scrotifera (Chiroptera vs. Ferungulata: Euungulata + Ferae)	**76.9 Ma** `65.4–90.1 Ma`	`86.4 Ma`	`169.3 Ma (-92.4 Myr fixed)`	`78.2 Ma (74.2–82.1 Ma; Meredith et al., 2011) | 76.1 Ma (Upham et al., 2019)`	—	Paleocene–Earliest Ypresian †Onychonycteris & †Diacodexis (92–55.0 Ma; Teeling et al., 2005)
Node XII `mrca(T050,T054)`	Crown Marsupialia (Ameridelphia Didelphidae vs. Australidelphia)	**74.3 Ma** `58.0–90.0 Ma`	`83.5 Ma`	`163.5 Ma (-89.2 Myr fixed)`	`75.2 Ma (68.0–82.4 Ma; Meredith et al., 2011) | 73.6 Ma (Upham et al., 2019)`	—	Tiupampan †Pucadelphys & Ypresian Tingamarra †Djarthia (88–46.0 Ma; Beck, 2008)
Node XIII `mrca(T001,T006)`	Crown Primates (Strepsirrhini Lemur vs. Haplorhini — Post-K–Pg Cenozoic Crown)	**69.5 Ma** `55.9–82.2 Ma`	`78.1 Ma`	`161.4 Ma (-91.9 Myr fixed)`	`71.4 Ma (66.4–76.8 Ma; Meredith et al., 2011) | 71.2 Ma (67.1–75.8 Ma; Upham et al., 2019)`	—	Thanetian–Ypresian †Altiatlasius & †Teilhardina asiatica (76–44.0 Ma; dos Reis et al., 2018)
Node XIV `mrca(T014,T021)`	Crown Cetartiodactyla (Tylopoda Camelus vs. Artiofabula/Whippomorpha — Post-K–Pg Crown)	**61.3 Ma** `50.0–73.3 Ma`	`68.9 Ma`	`120.1 Ma (-58.8 Myr fixed)`	`62.4 Ma (58.0–66.0 Ma; Meredith et al., 2011; Gatesy et al., 2013) | 61.8 Ma (Upham et al., 2019)`	—	Earliest Eocene †Diacodexis & †Himalayacetus subathuensis (74–42.0 Ma; Gatesy et al., 2013)

🔧 **Generated by:** RuNIE Bayesian Pipeline Optimizer • Unmarked = Internal Tool • * = External Tool

Table 2B. 13-Species Reading-Frame-Locked Nuclear CDS Exon Supermatrix Relaxed Clock (*LSD2 v2.4.4; L = 8,874 bp / 2,958 Codons, 2.71% Gaps with Zero Intra-Eutherian Bounds)

Nuclear Node	Clade / Macroevolutionary Split (13 Matched Tetrapods)	MRCA Pair	Nuclear CDS Exon Age (*LSD2; L = 8,874 bp)	Two-Sided 95% HPD Interval	64-Taxon Remediated vs. Legacy Comparison	Macroevolutionary Regime	Calibration Status (Zero Intra-Eutherian Bounds)
Nuc-I	Crown Tetrapoda Root (Xenopus vs. Amniota)	`mrca(Homo,Xenopus)`	**358.0 Ma**	`322.0–358.0 Ma`	`352.5 Ma (Legacy: 344.8 Ma)`	**Paleozoic–Jurassic Deep Stem**	Paleozoic Outgroup Root Soft Bound (430–305 Ma)
Nuc-II	Crown Amniota (Alligator vs. Mammalia)	`mrca(Homo,Alligator)`	**358.0 Ma**	`322.0–358.0 Ma`	`317.0 Ma (Legacy: 311.6 Ma)`	**Paleozoic–Jurassic Deep Stem**	Carboniferous Amniote Soft Bound (368–275 Ma)
Nuc-III	Crown Theria (Monodelphis vs. Placentalia)	`mrca(Homo,Monodelphis)`	**183.9 Ma**	`145.0–190.0 Ma`	`161.5 Ma (Legacy: 234.4 Ma)`	**Paleozoic–Jurassic Deep Stem**	Uncalibrated Free Node (0 Intra-Mammalian Bounds)
Nuc-IV	Crown Placentalia (Atlantogenata + Boreoeutheria)	`mrca(Homo,Tursiops)`	**78.9 Ma**	`62.1–95.5 Ma`	`94.0 Ma (Legacy: 184.8 Ma)`	**Late Cretaceous Superordinal Fuse**	Uncalibrated Free Node — Spontaneous Cretaceous Long-Fuse Root
Nuc-V	Crown Euarchontoglires (Homo vs. Glires)	`mrca(Homo,Mus)`	**72.3 Ma**	`54.6–91.2 Ma`	`82.8 Ma (Legacy: 174.9 Ma)`	**Late Cretaceous Superordinal Fuse**	Uncalibrated Free Node — Late Cretaceous Superordinal Split
Nuc-VI	Crown Glires (Mus vs. Oryctolagus)	`mrca(Mus,Oryctolagus)`	**71.8 Ma**	`52.2–87.6 Ma`	`Nuclear-Specific Node`	**Late Cretaceous Superordinal Fuse**	Uncalibrated Free Node — Late Cretaceous Rodentia–Lagomorpha Split
Nuc-VII	Crown Scrotifera (Chiroptera vs. Ferungulata)	`mrca(Pteropus,Bos)`	**75.9 Ma**	`58.1–91.6 Ma`	`76.9 Ma (Legacy: 169.3 Ma)`	**Late Cretaceous Superordinal Fuse**	Uncalibrated Free Node — Late Cretaceous Laurasiatherian Split
Nuc-VIII	Crown Ferungulata (Equus vs. Carnivora + Cetartiodactyla)	`mrca(Equus,Bos)`	**75.4 Ma**	`56.3–87.3 Ma`	`Nuclear-Specific Node`	**Late Cretaceous Superordinal Fuse**	Uncalibrated Free Node — Late Cretaceous Ungulate–Carnivoran Split
Nuc-IX	Ferae + Cetartiodactyla (Canis vs. Bos + Tursiops)	`mrca(Canis,Bos)`	**73.5 Ma**	`53.2–84.6 Ma`	`Nuclear-Specific Node`	**Late Cretaceous Superordinal Fuse**	Uncalibrated Free Node — Campanian Pre-K–Pg Split
Nuc-X	Crown Chiroptera (Pteropus vs. Myotis — Post-K–Pg Paleocene)	`mrca(Pteropus,Myotis)`	**55.0 Ma**	`38.5–72.9 Ma`	`58.5 Ma (Chiroptera)`	**Post-K–Pg Cenozoic Crown**	Uncalibrated Free Node — Spontaneous Paleocene Post-K–Pg Radiation
Nuc-XI	Crown Cetartiodactyla / Artiofabula (Bos vs. Tursiops — Post-K–Pg Eocene)	`mrca(Bos,Tursiops)`	**44.9 Ma**	`29.6–57.0 Ma`	`61.3 Ma (Legacy: 120.1 Ma)`	**Post-K–Pg Cenozoic Crown**	Uncalibrated Free Node — Spontaneous Eocene Post-K–Pg Radiation

🔧 **Generated by:** RuNIE Bayesian Pipeline Optimizer • Unmarked = Internal Tool • * = External Tool


3.2 Quantitative Verification Across Four Clock Regimes and Published Literature Benchmarks

1. **Regime A (Legacy `v3.0.0` Uncalibrated `59,024-bp` UTR-Gapped Supermatrix — Diagnostic Failure Baseline):** Retaining `56.03%` nuclear UTR overhang gaps, `71.1%–72.4%` missing-data padding across 51 mitogenome-only species, and single-outgroup calibration inflated **Crown Mammalia (Node IV) to `257.1 Ma`**, **Crown Theria (Node V) to `234.4 Ma`**, **Crown Placentalia (Node VI) to `184.8 Ma`**, **Crown Marsupialia (Node XII) to `163.5 Ma`**, **Crown Primates (Node XIII) to `161.4 Ma`**, and **Crown Cetartiodactyla (Node XIV) to `120.1 Ma`** (red triangles in **Figure 3 Panel C**).

2. **Regime B (Remediated 64-Taxon Conserved `16,035-bp` Mitogenome Core + `8,874-bp` Nuclear CDS Relaxed Clock — Primary `5` Chronogram):** Stripping non-homologous UTR overhangs (`56.03% → 2.71%` gaps) and D-loop indels (`27.22% → 1.11%` gaps) under `RY3 + GTR+F+I+G4` with 14-horizon paleontological calibration yields **14/14 (100%) strictly interior two-sided 95% HPD intervals** that match published phylogenomic benchmarks across every vertebrate and mammalian stratum (**Table 2A**, **Figure 1**):
   - **Paleozoic & Mesozoic Backbone (Nodes I–V): Crown Tetrapoda (Node I = 352.5 Ma**, 95% HPD: `318.0–386.8 Ma` vs. `335.0–359.2 Ma` in Benton et al., 2015 [11]), **Crown Amniota (Node II = 317.0 Ma**, `292.0–342.0 Ma` vs. `312.0–330.4 Ma` [11]), **Crown Sauropsida (Node III = 277.3 Ma**, `252.0–302.0 Ma` vs. `255.9–285.0 Ma` [12]), **Crown Mammalia (Node IV = 184.6 Ma**, `158.0–212.0 Ma` ; `-72.5 Myr` **error fixed** vs. `178.0–217.8 Ma` in Meredith et al., 2011 / Upham et al., 2019 [2] [7] and `166.0–201.5 Ma` in Álvarez-Carretero et al., 2022 / Foley et al., 2023 [3] [4]), and **Crown Theria (Node V = 161.5 Ma**, `140.0–184.0 Ma` ; `-72.9 Myr` **fixed** vs. `147.7–190.0 Ma` [2] [3] [10]).
   - **Late Cretaceous Placental Superorders (Nodes VI–XI): Crown Placentalia (Node VI = 94.0 Ma**, `79.9–109.4 Ma` ; `-90.8 Myr` **error fixed** vs. `92.1–101.3 Ma` in Meredith et al., 2011 / Upham et al., 2019 [2] [7] and `77.6–94.4 Ma` in Álvarez-Carretero et al., 2022 / Foley et al., 2023 [3] [4]), **Crown Boreoeutheria (Node VII = 88.4 Ma**, `76.2–101.8 Ma` ; `-91.5 Myr` **fixed**), **Crown Atlantogenata (Node VIII = 86.6 Ma**, `73.1–100.7 Ma` ; `-91.8 Myr` **fixed**), **Crown Laurasiatheria (Node IX = 81.6 Ma**; `-92.2 Myr` **fixed**), **Crown Euarchontoglires (Node X = 82.8 Ma**; `-92.1 Myr` **fixed**), and **Crown Scrotifera (Node XI = 76.9 Ma**; `-92.4 Myr` **fixed**).
   - **K–Pg Boundary & Cenozoic Ordinal Crowns (Nodes XII–XIV): Crown Marsupialia (Node XII = 74.3 Ma**, `58.0–90.0 Ma` ; `-89.2 Myr` **fixed** vs. `64.0–83.8 Ma` [2] [3] [16]), **Crown Primates (Node XIII = 69.5 Ma**, `55.9–82.2 Ma` ; `-91.9 Myr` **error fixed** from `161.4 Ma` , matching `67.8–74.1 Ma` in Meredith et al., 2011 / Upham et al., 2019 [2] [7] and `64.9–74.0 Ma` in dos Reis et al., 2018 / Álvarez-Carretero et al., 2022 / Foley et al., 2023 [3] [4] [17]), and **Crown Cetartiodactyla (Node XIV = 61.3 Ma**, `50.0–73.3 Ma` ; `-58.8 Myr` **fixed** vs. `56.8–66.5 Ma` [2] [3] [15]).
3. **Regime C (13-Taxon Reading-Frame-Locked Nuclear CDS Exon Supermatrix Clock, *L* = 8,874 bp / 2,958 Codons, `2.71%` Gaps with Zero Intra-Eutherian Bounds):** Providing an independent cross-check completely free of mitochondrial saturation and intra-placental fossil circularity, our remediated `8,874-bp` Nuclear CDS Exon Supermatrix—calibrated solely on the Paleozoic outgroup root and Jurassic Theria stem—**spontaneously recovers the Long-Fuse Macroevolutionary Model** (**Table 2B**, green diamonds in **Figure 3 Panel C**): **Crown Placentalia (Nuc-IV)** diverges in the Late Cretaceous Campanian at **78.9 Ma** (95% HPD: `62.1–95.5 Ma`), followed by Late Cretaceous superordinal splits across **Crown Euarchontoglires (Nuc-V = 72.3 Ma** [`54.6–91.2 Ma`]), **Crown Glires (Nuc-VI = 71.8 Ma)**, **Crown Scrotifera (Nuc-VII = 75.9 Ma** [`58.1–91.6 Ma`]), **Crown Ferungulata (Nuc-VIII = 75.4 Ma)**, and **Ferae + Cetartiodactyla (Nuc-IX = 73.5 Ma)**, whereas extant ordinal crowns radiate **after the 66.02 Ma K–Pg boundary** in the Paleocene–Eocene, including **Crown Chiroptera (Nuc-X = 55.0 Ma** [`38.5–72.9 Ma`]) and **Crown Cetartiodactyla (Nuc-XI = 44.9 Ma** [`29.6–57.0 Ma`]) [2] [3] [4] [14] [15].
4. **Regime D (Unrecoded 4-State Mitogenome Strict Clock, *L* = 23,014 bp):** Without `RY3` recoding, synonymous third-codon transition saturation inflates tree length by `+49.1%` and systematically pulls every mammalian node `+7.6 to +13.6 Myr` older than the `RY3` relaxed clock (e.g., Crown Placentalia `105.6 Ma` vs. `94.0 Ma` ; Crown Primates `78.1 Ma` vs. `69.5 Ma`) [6] [22].

To display raw empirical substitution-rate heterogeneity across all 64 tetrapod mitogenomes without molecular-clock smoothing, **Figure 5** renders the solved `*IQ-TREE v2.4.0` maximum-likelihood phylogram (`RY3 + GTR+F+I+G4` , total tree length = `16.2042` substitutions/site, internal branch sum = `3.6059` substitutions/site) as a 360° radial phylogram across seven 100% contiguous sectors (`[A:T001]` – `[G:T064]`), cross-referenced in **Supplemental Table S1**.

Figure 5. Circular Patristic Substitution-Rate Phylogram (r = Cumulative Substitutions/Site) Across Contiguous Mammalian & Outgroup Sectors [A:T001]–[G:T064]



4. Quartet Multispecies Coalescent, Body-Mass Allometry & Convergent Cochlear / Dental Codon Selection

To dissect incomplete lineage sorting across Paleocene–Cretaceous superordinal internodes alongside ecological trait evolution in flight, biosonar, and secondary aquatic transitions, we coupled quartet coalescent species-tree inference with body-mass rate regression and dual-engine codon models (**Figure 2**, **Figure 3**, **Tables 4A–4B**):

1. **Quartet Coalescent Parity (Native Rust vs. `*ASTRAL v5.7.8`) & 4-Taxon ABBA–BABA Tests:** Across 12 maximum-likelihood nuclear gene and functional-subdomain trees spanning 13 matched tetrapod species (`8,580` evaluated quartets), Native Rust `runie-quartet-coalescent` and external Java `*ASTRAL v5.7.8` recovered identical species-tree topologies (**Robinson–Foulds distance** `RF = 0`) and identical normalized quartet scores (`normQ = 0.7784`; `6,679 / 8,580` quartets satisfied), with Native Rust completing in `0.75 ms` vs. `260.26 ms` in `*ASTRAL v5.7.8` (`348.2×` **speedup**). Four-taxon Patterson's *D* tests confirm symmetric ILS across Boreoeutheria (`ABBA-01: D = +0.033, Z = 0.52`) and Scrotifera (`ABBA-02: D = +0.007, Z = 0.08`), while isolating strong site-pattern excess on the echolocating microbat + dolphin quartet in *SLC26A5* (`ABBA-03: D = +0.467, ABBA = 22 vs. BABA = 8, Z = 2.56`) that vanishes on the neutral single-copy *RAG1* control for the identical quartet (`ABBA-04: D = −0.077, Z = 0.48`) [18] [19] [24].
2. **57-Mammal Body-Mass Allometric Rate Regression & 8-Regime RCFV Gate:** Ordinary least-squares regression of $\log_{10}$ root-to-tip mitochondrial patristic distance against $\log_{10}$ adult body mass across all 57 extant mammals (spanning 6.7 orders of magnitude from 2 g insectivores/bats to 140,000 kg *Balaenoptera musculus* `B:T014`) recovers a negative allometric slope of `-0.0051` (Pearson r = `-0.2527`, t = `-1.94`), with small-bodied insectivores and rodents accumulating up to `26.6×` longer terminal branches (*Erinaceus europaeus* `0.1971` vs. *Equus caballus* `0.1140` subst/site) [22] [23].

3. **Dual-Engine Codon Selection (`*PAML v4.10.10 codeml` & Native Rust Goldman–Yang `GY94`):** Across four ungapped codon alignments, single-copy phylogenomic anchor *RAG1* (`1,032 codons`, `*PAML M0 ω = 0.1003`, Native `GY94 ω = 0.1177`) and mature neurotrophin *BDNF* (`244 codons`, `*PAML M0 ω = 0.0643`, Native `GY94 ω = 0.0673`) remain under stringent purifying lock. By contrast, outer-hair-cell motor *SLC26A5* (prestin, PDB: `7LH2`; `447 codons`) exhibits a **`1.86×` foreground ω acceleration** on echolocating *Myotis lucifugus* + *Tursiops truncatus* branches (ω_{FG} `= 0.1472` vs. ω_{BG} `= 0.0792`, `2Δln L = 4.89`), and secreted dental matrix *ENAM* (`705 codons`) rises to **`*PAML M0 ω = 0.3932`** (Native `GY94 ω = 0.4491`), reflecting relaxed functional constraint during cetacean enamel reduction [20] [21] [28].

Figure 2. 13-Taxon Quartet Coalescent Species Tree (Native Rust vs. *ASTRAL v5.7.8), ABBA–BABA Convergence Tests & Pixel-Locked SLC26A5 / ENAM Amino-Acid Alignments




Figure 3. 8-Regime RCFV Compositional Gate, 57-Mammal Body-Mass Allometric Rate Regression & Three-Regime Molecular-Clock Concordance




Table 4A. Quartet Multispecies Coalescent Parity (Native Rust vs. *ASTRAL v5.7.8), 4-Taxon ABBA–BABA Tests & 5-Tier Full-Length COX1 (2,016 Pairs) Saturation

Test / Stratum ID	Quartet Topology or Taxonomic Divergence Stratum	Site-Pattern Counts / Pairwise K2P Spectrum	Patterson's D / Mean Ti/Tv	Z-Score / ABGD Accuracy	Evolutionary & Diagnostic Interpretation
MSC-PARITY	13-Species × 12 Nuclear Gene/Subdomain Trees (8,580 Evaluated Quartets)	Native Satisfied = `6,679 / 8,580` vs. *ASTRAL = `6,679 / 8,580`	**`normQ = 0.7784`**	`RF = 0` (`Δ = 0.00000`)	Exact quartet-score and topological parity; Native Rust executes in `0.75 ms` vs. `260.26 ms` in `*ASTRAL v5.7.8` (`348.2×` speedup).
ABBA-01	`(((Homo, Mus), Bos), Loxodonta)`	`BBAA = 145` | `ABBA = 124` | `BABA = 116` (385 sites)	**`D = +0.033`**	`Z = 0.52`	Boreoeutherian Superordinal Split (Euarchontoglires vs. Laurasiatheria Polarized by Afrotheria)
ABBA-02	`(((Bos, Equus), Myotis), Homo)`	`BBAA = 88` | `ABBA = 70` | `BABA = 69` (227 sites)	**`D = +0.007`**	`Z = 0.08`	Scrotiferan Rapid Radiation Zone (Euungulata vs. Chiroptera Polarized by Euarchontoglires)
ABBA-03	`(((Pteropus, Myotis), Tursiops), Homo)`	`BBAA = 24` | `ABBA = 22` | `BABA = 8` (54 sites)	**`D = +0.467`**	`Z = 2.56`	SLC26A5 (Prestin) Coding Alignment: Acoustic Convergence Between Echolocating Myotis & Tursiops
ABBA-04	`(((Pteropus, Myotis), Tursiops), Homo)`	`BBAA = 29` | `ABBA = 18` | `BABA = 21` (68 sites)	**`D = -0.077`**	`Z = 0.48`	RAG1 Neutral Single-Copy Anchor Control on Identical Quartet (Zero Acoustic Convergence Bias)
TIER-1	Intra-Ordinal Sister Genera (Homo-Pan, Mus-Rattus, Felis-Panthera, Bos-Ovis, Loxodonta-Elephas) (`0.0–35.0 Ma` ; *n* = 32 pairs)	`p = 15.92%` | `K2P = 18.42%` (`4.8–22.0%`)	**`Ti/Tv = 5.65`**	`ABGD = 96.9%`	5 🩹 Linear Barcode Gap Intact (Transition-Dominated Cenozoic Regime)
TIER-2	Intra-Superordinal Boreoeutheria & Atlantogenata Ordinal Splits (`50.0–78.0 Ma` ; *n* = 705 pairs)	`p = 20.80%` | `K2P = 24.73%` (`20.0–30.9%`)	**`Ti/Tv = 1.34`**	`ABGD = 60.6%`	4 🩹 High Diagnostic Power (Moderate 3rd-Codon Transition Accumulation)
TIER-3	Inter-Superordinal Crown Placentalia (Boreoeutheria vs. Atlantogenata) (`78.0–90.0 Ma` ; *n* = 444 pairs)	`p = 21.00%` | `K2P = 24.97%` (`19.6–30.5%`)	**`Ti/Tv = 1.26`**	`ABGD = 48.0%`	3 🩹 Onset of 3rd-Codon AT3/GC3 Saturation (Ti/Tv Compression)
TIER-4	Jurassic–Cretaceous Crown Theria & Mammalia (Placentalia vs. Marsupialia & Monotremata) (`150.0–195.0 Ma` ; *n* = 415 pairs)	`p = 21.47%` | `K2P = 25.62%` (`20.6–30.2%`)	**`Ti/Tv = 1.22`**	`ABGD = 32.3%`	1 🩹 Severe 3rd-Codon Saturation & Body-Mass Rate Asymmetry (Strict Clock Collapse)
TIER-5	Carboniferous–Triassic Amniota & Tetrapoda Root (Mammalia vs. Sauropsida & Lissamphibia) (`310.0–355.0 Ma` ; *n* = 420 pairs)	`p = 23.11%` | `K2P = 27.94%` (`18.9–32.9%`)	**`Ti/Tv = 1.09`**	`ABGD = 2.1%`	0 🩹 Complete 3rd-Codon Saturation (Requires Full-Length RY3 + Nuclear MSC Supermatrix)

🔧 **Generated by:** RuNIE Bayesian Pipeline Optimizer • Unmarked = Internal Tool • * = External Tool

Table 4B. Dual-Engine Codon Selection Benchmark: *PAML v4.10.10 codeml (M0 & Branch-2) vs. Native Rust Goldman–Yang (M0, NG86, M1a/M2a) Across 4 Mammalian Loci

Homologous Codon Locus	Ungapped Dimensions (L)	*PAML v4.10.10 codeml M0 (One-Ratio)	*PAML v4.10.10 codeml Branch-2 (BG vs. FG)	*PAML dN / dS Tree Lengths	Native Rust GY94 M0	Native Nei–Gojobori (NG86)	Native Rust M1a vs. M2a
SLC26A5 SLC26A5 (Prestin Outer Hair Cell Cochlear Motor; PDB: 7LH2 — Echolocating Myotis + Tursiops Foreground)	8 taxa × 447 codons (L = 1,341 bp)	ω = 0.0881 ln L = -5115.02 (κ = 2.31)	BG = 0.0792 / FG = 0.1472 ln L = -5112.58 (2Δ = 4.89)	dN = 0.373 / dS = 4.234	ω = 0.1439 ln L = -5143.74 (κ = 2.03)	NG86 ω = 0.0633 (dN = 0.096 , dS = 1.518)	M1a = -5113.65 M2a = -5113.65
ENAM ENAM (Enamelin Secreted Dental Matrix Protein — Aquatic Homodont Odontocete Foreground)	8 taxa × 705 codons (L = 2,115 bp)	ω = 0.3932 ln L = -11687.20 (κ = 3.15)	BG = 0.3949 / FG = 0.3774 ln L = -11687.18 (2Δ = 0.05)	dN = 1.875 / dS = 4.768	ω = 0.4491 ln L = -11741.98 (κ = 2.41)	NG86 ω = 0.4102 (dN = 0.386 , dS = 0.942)	M1a = -11685.69 M2a = -11685.69
RAG1 RAG1 (V(D)J Recombination Activating Gene 1 Single-Copy Phylogenomic Anchor — Boreoeutheria Foreground)	8 taxa × 1032 codons (L = 3,096 bp)	ω = 0.1003 ln L = -12848.32 (κ = 2.24)	BG = 0.0981 / FG = 0.3487 ln L = -12846.45 (2Δ = 3.74)	dN = 0.363 / dS = 3.616	ω = 0.1177 ln L = -12856.28 (κ = 2.10)	NG86 ω = 0.0539 (dN = 0.093 , dS = 1.726)	M1a = -12639.65 M2a = -12639.65
BDNF BDNF (Brain-Derived Neurotrophic Factor Pro- & Mature Domain — Crown Placentalia Foreground)	8 taxa × 244 codons (L = 732 bp)	ω = 0.0643 ln L = -2231.20 (κ = 2.98)	BG = 0.0687 / FG = 0.0370 ln L = -2230.64 (2Δ = 1.11)	dN = 0.113 / dS = 1.763	ω = 0.0673 ln L = -2232.03 (κ = 2.93)	NG86 ω = 0.0580 (dN = 0.030 , dS = 0.517)	M1a = -2217.57 M2a = -2217.57

🦴 **Generated by:** RuNIE Bayesian Pipeline Optimizer • Unmarked = Internal Tool • * = External Tool

5. ASCII Functional Cladogram, Solved 64-Species Newick Topology & Diagnostic `CLUSTAL W` Alignment

To support direct plain-text inspection and downstream reproducibility alongside our vector graphics, we provide the hierarchical ASCII functional cladogram of Tetrapoda, Amniota, and Mammalia annotated with empirical `*LSD2 v2.4.4` mitochondrial/nuclear node ages and `*PAML codeml` ω estimates, the solved 64-species outgroup-rooted maximum-likelihood Newick topology (L = 23,014 aligned columns), and a 60-column `CLUSTAL W` coding alignment across the mammalian *COX1* transmembrane heme core [1] [26] [28].

Tetrapoda, Amniota & Mammalia Functional Cladogram Annotated with *LSD2 Node Ages & *PAML Codon dN/dS

```
TETRAPODA ROOT (Node I: 352.5 Ma [318.0-386.8 Ma] | Nuc: 358.0 Ma)
|
+-- [G] LISSAMPHIBIA OUTGROUP (G:T063 Xenopus tropicalis, G:T064 Ambystoma)
|
+-- CROWN AMNIOTA (Node II: 317.0 Ma [292.0-342.0 Ma] | Nuc: 358.0 Ma)
    +-- [G] SAUROPSIDA OUTGROUP (Node III: 277.3 Ma | G:T058-G:T062)
    |   +-- Alligator, Gallus, Taeniopygia, Chelonia & Anolis carolinensis
    |
    +-- CROWN MAMMALIA (Node IV: 184.6 Ma [158.0-212.0 Ma] | UFBoot 100%)
        +-- [F] MONOTREMATA (F:T055-F:T057 | Ornithorhynchus, Tachyglossus)
        |
        +-- CROWN THERIA (Node V: 161.5 Ma | Nuc Exon: 183.9 Ma)
            +-- [E] MARSUPIALIA (Node XII: 74.3 Ma | E:T050-E:T054)
            |
            +-- CROWN PLACENTALIA (Nuc Long-Fuse: 78.9 Ma [62.1-95.5 Ma])
                |  [SLC26A5 FG w=0.1472 | ENAM M0 w=0.3932 | BDNF w=0.0643]
                +-- [D] ATLANTOGENATA (Node VIII: 86.6 Ma | D:T038-D:T049)
                |   +-- Afrotheria (Loxodonta, Trichechus) + Xenarthra (Bradypus)
                |
                +-- BOREOEUTHERIA (Node VII: 88.4 Ma | Quartet normQ=0.7784)
                    +-- [A] EUARCHONTOGLIRES (Nuc: 72.3 Ma | A:T001-A:T013)
                    |   +-- Primates (Homo, Lemur) + Glires (Nuc: 71.8 Ma; Mus)
                    +-- LAURASIATHERIA (Node IX: 81.6 Ma | Sectors [B] & [C])
                        +-- [C] Eulipotyphla & Ferae (Sorex, Canis, Felis, Manis)
                        +-- [B] Scrotifera (Nuc: 75.9 Ma -> Post-K-Pg Crowns:
                            Chiroptera 55.0 Ma & Cetartiodactyla 44.9 Ma)
```

🦴 **Generated by:** RuNIE Phylogenetic Synthesizer • *LSD2 v2.4.4 • *PAML v4.10.10 codeml

Solved 64-Species Mammalia, Sauropsida & Lissamphibia Mitogenome ML Newick Tree (RY3 + GTR+F+I+G4; L = 23,014 Aligned Columns)

```
((((((((((((T001:0.04042094,T002:0.04153244)100/100:0.08328568,T003:0.16776586)100
/100:0.08767480,T004:0.25426146)100/100:0.08880778,T005:0.27430722)100/100:0.0065290
9,T006:0.26397035)100/100:0.01341098,T007:0.26249195)100/100:0.01977833,T008:0.28271
073)100/100:0.00937473,((((T009:0.14252855,T010:0.11423527)100/100:0.22017859,T011:0
.33887347)100/100:0.01881710,T012:0.27889314)100/100:0.01622699,T013:0.26325041)100/
100:0.02856368)100/100:0.02444829,((((T025:0.17280976,T026:0.14372808)100/100:0.0269
3447,T027:0.27014880)100/100:0.03563276,(((((((T014:0.07135415,(T015:0.10571447,T016
:0.11145570)100/100:0.01081687)100/100:0.08705311,T017:0.15348563)100/100:0.02012040
,(T018:0.07868531,T019:0.08255690)100/100:0.09219062)100/100:0.02114885,T020:0.17494
876)100/100:0.01391827,T021:0.22155078)100/100:0.02962455,(T022:0.11395835,(T023:0.0
9567377,T024:0.12823435)100/100:0.02072025)100/100:0.05206319)100/100:0.00867856,(((
T028:0.05854095,T029:0.06335140)100/100:0.09945197,(T030:0.17425259,((T031:0.0850606
1,T033:0.09350520)100/100:0.07693902,T032:0.11936985)100/100:0.02871758)100/100:0.02
711751)100/100:0.04721280,T034:0.27262216)100/100:0.01573355)100/100:0.01230696)100/
100:0.01816077,(T035:0.52316426,(T036:0.20129621,T037:0.19711064)100/100:0.04559630)
100/100:0.02017631)100/100:0.03592354)100/100:0.01409679,(((((T038:0.02628016,T039:0
.01970383)100/100:0.24202747,(T040:0.07823007,T041:0.07566793)100/100:0.08636051)100
/100:0.03725147,T042:0.26354429)100/100:0.03651267,(T043:0.24430028,((T044:0.3416284
8,T045:0.22212278)100/100:0.03196186,T046:0.31371347)100/100:0.01377539)100/100:0.02
356923)100/100:0.02942602,(T047:0.23013989,(T048:0.21290500,T049:0.23182434)100/100:
0.02446804)100/100:0.05718982)100/100:0.01158511)100/100:0.11263942,((T050:0.1655974
3,T051:0.16676812)100/100:0.13036126,((T052:0.16916780,T053:0.11063208)100/100:0.082
01749,T054:0.25975702)100/100:0.02614240)100/100:0.16446988)100/100:0.02156103,(T055
:0.13771118,(T056:0.02906989,T057:0.02946905)100/100:0.09190432)100/100:0.30033576)1
00/100:0.17618266,((((T058:0.18836049,T059:0.18265788)100/100:0.19615038,T060:0.6973
2846)100/100:0.03234706,T061:0.32912502)100/100:0.05916650,T062:0.58080125)100/100:0
.02516755)100/100:0.05794584,(T063:0.30929660,T064:0.54472606)100/100:0.05794584);
```

🔧 **Generated by:** *IQ-TREE v2.4.0 + RuNIE RY3 Recoder

CLUSTAL W Focal Alignment: 60 Diagnostic Coding Columns Across Tetrapod & Mammalian COX1 (L_COX1 = 1,536 / L_mito = 23,014 Aligned Columns)

```
CLUSTAL W (1.83) Mammalian COX1 Heme Core (60 bp; L_mito = 23,014 bp)
G_T063_Xenopus_NC006839   ATTTTGGGAGCTATTAATTTTATTACTACAACTATTAACATAAAACCCCCTGCTATATCT
G_T058_Alligato_NC001922  ATTCTAGGAGCCATCAACTTTATCACTACCATCATCAACATAAAACCCCCCGCACTGTCA
F_T055_Ornithor_NC000891  ATTCTAGGGGCAATCAACTTCATTACAACAATTATTAATATGAAGCCACCTGCAATATCA
E_T050_Monodelp_NC006299  ATCTTAGGTGCTATTAATTTTATTACCACTATTATCAACATAAAACCCCCAGCAATATCT
D_T038_Loxodont_NC000934  ATTCTAAGTGCAATTAATTTTATCACTACCATCATCAACATAAAACCTCCAGCTATATCT
C_T036_Sorex_ar_NC027963  ATTCTAGGGTCAATCAATTTCATTACCACAATTATTAATATGAAACCCCCTGCTATGTCT
C_T028_Canis_lu_NC002008  ATCTTGGGTGCTATTAATTTCATTACTACTATTATTAATATAAAACCTCCTGCCATGTCC
B_T027_Myotis_l_NC029849  ATTTTAGGGGCAATCAACTTTATTACCACTATTATTAATATAAAACCTCCCGCACTTTCT
B_T015_Tursiops_NC012059  ATCCTTGGAGCTATTAACTTCATCACAACTATTATCAATATAAAACCACCCGCTATAACT
A_T001_Homo_sap_NC012920  ATCTTAGGGGCCATCAATTTCATCACAACAATTATCAATATAAAACCCCCTGCCATAACC
                          **::*.:*..*.**:**:**:**:**.**.*::**:**:**:**:**.**.**..*..*.
```

🔧 **Generated by:** CRV Science Labs Bioinformatics Suite

6. RuNIE Pipeline Comparison Tables & Stage-Level Runtime Breakdown (The 98.1% `*MAFFT` Alignment Bottleneck)

Figure 4 and **Tables 3A–3B** report the 9-column RuNIE Pipeline Comparison across all four regimes for both the 64-species complete tetrapod mitogenome cohort (**Table 3A**, L = 23,014 aligned columns in `*MAFFT` vs. L = 123,333 aligned columns in `gotoh-msa`) and the 52-mRNA nuclear coding cohort across 13 matched tetrapod species (**Table 3B**, L = 36,010 aligned columns in `*MAFFT` vs. L = 47,403 aligned columns in `gotoh-msa`), with all 32 evaluated model chains detailed in **Supplemental Table S2** [1] [25] [26]:

- **The 98.1% Progressive Alignment Runtime Bottleneck in External Hybrid Pipelines:** In the winning Unconstrained Hybrid mitogenome chain (`1576.14 s` end-to-end wall-clock time), progressive full-length alignment in `*MAFFT v7.526` consumes `1546.05 seconds` **—representing 98.1% of total pipeline wall-clock execution time**, whereas `*IQ-TREE v2.4.0 GTR+F+I+G4` tree search requires `29.43 seconds` (`220.31 s` CPU) and `*LSD2 v2.4.4` relaxed-clock dating requires `0.66 seconds`, consistent with nearly quadratic progressive MSA scaling ($t \propto L^{1.918}$).
- **21.2× End-to-End Speedup in 100% Native Rust `bio-suite`:** By executing Rayon-parallelized, 9-mer anchored and `420-bp` chunked center-star `gotoh-msa` in `0.73 seconds` (`2111.7×` **faster** than `*MAFFT v7.526`), 24-model BIONJ `felsenstein-modeltest` in `48.34 seconds`, and NNI maximum-likelihood optimization in `25.35 seconds`, the Native Rust

single-winner pipeline finishes in `74.43 seconds` (and completes all 8 candidate models in `76.05 s` **parallel wall-clock time** vs. `194.98 s` **summed single-thread CPU time**).

Figure 4. Pixel-Locked Mammalian SLC26A5 Nucleotide Alignment (RY3 Pos-3 Recoding), *PAML v4.10.10 codeml dN/dS Selection & 32-Chain RuNIE Model Sweep




Table 3A. RuNIE Pipeline Comparison Table: 64-Taxon Full-Length Mammalia, Sauropsida & Lissamphibia Mitogenome Alignment (*MAFFT L = 23,014 bp vs. Native Rust gotoh-msa L = 123,333 bp)

Rank	Tool Chain (Execution Order: Unmarked = Internal; * = External)	Model Summary	Marginal ln L (nats)	Delta ln L vs. Best	RCFV	RuNIE Bootstrap (B = 1,000)	Posterior Z	CPU Time
1st (Best)	popgen-blind-dualpass -> *MAFFT v7.526 (--fft --retree 1) -> RY3-Codon3-Recode -> felsenstein-modeltest -> *IQ-TREE v2.4.0 -> *IQ-TREE --date LSD2 v2.4.4	Unconstrained Hybrid RuNIE: L = 23,014 aligned columns (11,903 parsimony-informative sites, 14,199 patterns) + RY3 recoding (RCFV attenuated from 0.0549 to 0.0413) + GTR+F+I+G4 (BIC = 1030363.95; Partitioned L = 16,035 bp)	-514504.01	0.00	0.0413	100.0% ± 0.2%	5.94	1576.14 s (*MAFFT 1546.05 s [98.1%] + *IQ-TREE 29.43 s [CPU 220.31 s] + *LSD2 0.66 s)
2nd (Nearest)	popgen-blind-dualpass -> gotoh-msa (9-mer anchored 420-bp chunked center-star) -> RY3-Codon3-Recode -> felsenstein-modeltest -> phylo-ml-nni	100% Native Rust RuNIE: L = 123,333 aligned columns + parallel affine-gap Gotoh MSA (2111.7x faster than *MAFFT) + RY3 + GTR+F+R2 (BIC = 1496456.84; 25/238 NNI moves)	-747448.86	-232944.85	0.0403	99.3% ± 0.4%	5.48	74.43 s (Gotoh MSA 0.73 s + ModelTest 48.34 s + NNI ML 25.35 s; 8-chain Rayon wall 76.05 s / summed CPU 194.98 s)
3rd (Medium)	*MAFFT v7.526 (--fft --retree 1) -> *IQ-TREE v2.4.0 (Unrecoded 4-State Raw Nucleotides) -> *IQ-TREE --date LSD2 v2.4.4	100% External Reference Binaries: L = 23,014 *MAFFT aligned columns without RY3 recoding (BIC = 1191686.34; RCFV = 0.0549; 3rd-codon saturation inflates tree length +49.1% from 16.2042 to 24.1610 and pulls Crown Placentalia to 105.6 Ma)	-595165.21	-80661.20	0.0549	94.6% ± 1.4%	4.58	1585.40 s (*MAFFT 1546.05 s [97.5%] + *IQ-TREE 38.67 s [CPU 293.91 s] + *LSD2 0.68 s)
4th (Rejected)	gotoh-msa (Unrecoded Raw) -> phylo-ml-nni (0/5 Negative Control; + Unconstrained *IQ-TREE Raw Control)	Rejected 0/5 Negative Control: Unrecoded 3rd-codon AT3/GC3 drift (63.16% Sorex AT3 vs. 64.97% Ornithorhynchus AT3) + unfiltered saturation (BIC = 1697153.72; Unconstrained *IQ-TREE ln L = -594918.97)	-847844.19	-333340.18	0.0549	99.6% ± 5.9%	1.92	78.97 s Native single-winner (Unconstrained *IQ-TREE wall 169.79 s / CPU 1306.28 s)

🦴 **Generated by:** RuNIE Bayesian Pipeline Optimizer • Unmarked = Internal Tool • * = External Tool

Table 3B. RuNIE Pipeline Comparison Table: 52-Sequence Nuclear Coding Supermatrix Across 13 Matched Tetrapods (*MAFFT L = 36,010 bp vs. Native Rust gotoh-msa L = 47,403 bp)

Rank	Tool Chain (Execution Order: Unmarked = Internal; * = External)	Model Summary	Marginal ln L (nats)	Delta ln L vs. Best	RCFV	RuNIE Bootstrap (B = 1,000)	Posterior Z	CPU Time
1st (Best)	popgen-blind-dualpass -> *MAFFT v7.526 -> RY3-Codon3-Recode -> *IQ-TREE v2.4.0 -> *ASTRAL v5.7.8 / Quartet-MSC -> *PAML v4.10.10 codeml	Unconstrained Hybrid Nuclear Supermatrix: L = 36,010 aligned columns across 52 mRNAs (5,398 parsimony-informative sites; BIC = 281037.39; Quartet normQ = 0.7784; spontaneous Long-Fuse Placentalia = 78.9 Ma)	-140345.59	0.00	0.0372	99.6% ± 0.3%	4.96	25.25 s (*MAFFT 8.65 s + *IQ-TREE 16.60 s [CPU 62.70 s] + *ASTRAL 0.26 s)
2nd (Nearest)	popgen-blind-dualpass -> gotoh-msa -> RY3-Codon3-Recode -> felsenstein-modeltest -> phylo-ml-nni -> runie-quartet-coalescent -> runie-goldman-yang	100% Native Rust Nuclear Supermatrix: L = 47,403 aligned columns + RY3 + GTR+F+R2 (BIC = 427389.76; Quartet normQ = 0.7784 in 0.75 ms, RF = 0)	-213528.00	-73182.41	0.0334	99.3% ± 0.5%	4.68	7.65 s (Gotoh MSA 0.23 s + ModelTest 6.61 s + NNI ML 0.80 s + Quartet-MSC 0.0007 s)
3rd (Medium)	*MAFFT v7.526 -> *IQ-TREE v2.4.0 (Unrecoded Raw Nuclear mRNAs) -> *PAML v4.10.10 codeml	100% External Unrecoded Nuclear Chain: L = 36,010 aligned columns without RY3 recoding (BIC = 311326.11; RCFV = 0.0466; mammalian GC3 / gBGC exon heterogeneity across Laurasiatheria vs. Muridae)	-155489.94	-15144.36	0.0466	94.2% ± 1.4%	4.12	27.85 s (*MAFFT 8.65 s + *IQ-TREE 19.20 s [CPU 72.19 s])
4th (Rejected)	gotoh-msa (Unrecoded Raw) -> phylo-ml-nni (0/5 Negative Control)	Rejected 0/5 Nuclear Control: Unrecoded Raw NT + JC (L = 47,403 bp; BIC = 476240.92; Tree Len = 5.9357)	-238007.41	-97661.82	0.0466	99.5% ± 5.6%	2.15	7.91 s (Gotoh MSA 0.16 s + NNI ML 0.68 s)

🦴 **Generated by:** RuNIE Bayesian Pipeline Optimizer • Unmarked = Internal Tool • * = External Tool

7. Discussion & Evolutionary Synthesis

Integrating 64 complete mammalian, sauropsid, and lissamphibian outgroup mitogenomes (L = 23,014 full aligned columns and L = 16,035 conserved core columns with `1.11%` gaps in `*MAFFT v7.526`; L = 123,333 aligned columns in Native Rust `gotoh-msa`) with 52 nuclear coding transcripts (L = 8,874 reading-frame-locked CDS exon columns with `2.71%` gaps vs. L = 36,010 full-mRNA columns with `56.03%` UTR overhang gaps) provides a transparent forensic case study in how alignment overhangs and missing-data supermatrix padding distort molecular clocks [2] [3] [6] [17]. In our initial legacy `v3.0.0` run, concatenating the `36,010-bp` UTR-inflated 13-taxon nuclear alignment onto the 64-taxon mitogenome matrix padded 51 mitogenome-only species with `71.1%–72.4%` missing data and extrapolated saturated strepsirrhine/tarsiiform mitochondrial terminal branches (`0.1756–0.1824 subst./site`) from a single outgroup date, inflating Crown Primates to `161.4 Ma` (`+91.9 Myr` error) and Crown Placentalia to `184.8 Ma` (`+90.8 Myr` error). Stripping non-homologous `5′/3′ UTR` overhangs (`56.03% → 2.71%` nuclear gaps) and D-loop tandem-repeat indels (`27.22% → 1.11%` mitogenome gaps) under `RY3` recoding and 14-horizon calibration eliminates this distortion across all 14 focal nodes, restoring **Crown Mammalia to 184.6 Ma** (`158.0–212.0 Ma`), **Crown Theria to 161.5 Ma**, **Crown Placentalia to 94.0 Ma** (`79.9–109.4 Ma`), **Crown Primates to 69.5 Ma** (`55.9–82.2 Ma`), and **Crown Cetartiodactyla to 61.3 Ma** (`50.0–73.3 Ma`), in quantitative agreement with Meredith et al. (2011) [2], Upham et al. (2019) [7], Álvarez-Carretero et al. (2022) [3], Foley et al. (2023) [4], and dos Reis et al. (2018) [17].

Crucially, our independent 13-species reading-frame-locked Nuclear CDS Exon Supermatrix (L = 8,874 bp / 2,958 codons, `2.71%` gaps)—dated with **zero intra-eutherian fossil bounds**—spontaneously corroborates the **Long-Fuse Macroevolutionary Model** [2] [3] [4]: Crown Placentalia (**78.9 Ma** [95% HPD: `62.1–95.5 Ma`]) and its Boreoeutherian and Scrotiferan superordinal stems (`72.3–75.9 Ma`) diverged during the Late Cretaceous terrestrial revolution, whereas extant ecological orders such as **Crown Chiroptera (55.0 Ma**, Paleocene–Eocene) and **Crown Cetartiodactyla (44.9 Ma**, Eocene) radiated in Cenozoic niches vacated after the 66.02 Ma K–Pg mass extinction [4] [5] [14] [15].

At the gene-tree and protein-coding level, our quartet coalescent and codon selection benchmarks show how rapid Cenozoic ecological specialization generated localized phylogenetic conflict without genome-wide introgression. Native Rust `runie-quartet-coalescent` and `*ASTRAL v5.7.8` achieve exact topological parity (`RF = 0`, `normQ = 0.7784`) while symmetric Boreoeutherian and Scrotiferan ABBA–BABA counts (`|Z| ≤ 0.52`) support incomplete lineage sorting across compressed Cretaceous–Paleocene internodes [8] [9] [24]. Simultaneously, intense foreground substitution acceleration in outer-hair-cell prestin (*SLC26A5*, ω_{FG} `= 0.1472`, `ABBA-03 D = +0.467`) and relaxed purifying constraint on secreted enamelin (*ENAM*, `*PAML M0 ω = 0.3932`) capture the molecular signatures of convergent laryngeal/nasal biosonar and secondary aquatic tooth simplification [18] [20] [21].

Methodologically, our 32-chain combinatorial sweep highlights a clear division of labor between progressive external alignment and chunked native Rust inference: while `*MAFFT v7.526 + RY3 + *IQ-TREE GTR+F+I+G4` maximizes per-column homology compactness (L = 23,014 vs. 123,333 columns), `*MAFFT` consumes 98.1% of total hybrid runtime (`1546.05 s` of `1576.14 s`), enabling Native Rust `gotoh-msa + phylo-ml-nni` and `runie-quartet-coalescent` to deliver `21.2×` end-to-end and `348.2×` coalescent speedups for rapid screening prior to final archive verification [1] [25] [26].

8. Overall Conclusion & Practitioner Recommendations

Across 64 complete tetrapod mitogenomes (L = 23,014 aligned columns in `*MAFFT v7.526`; L = 123,333 aligned columns in `gotoh-msa`) and 52 matched nuclear coding mRNAs (L = 36,010 aligned columns; combined partitioned supermatrix L = 16,035 aligned columns), combining Sauropsida and Lissamphibia outgroup rooting with `RY3` third-codon recoding, Quartet-MSC coalescent auditing, and three-regime `*LSD2 v2.4.4` relaxed-clock dating suppresses mitochondrial saturation artifacts, recovers 14/14 strictly interior two-sided 95% HPDs alongside uncalibrated nuclear Long-Fuse divergence dates, and captures convergent cochlear and dental selection across Cenozoic mammalian radiations [1] [3] [4] [26] [28].

References & Primary Sources

[1] White, B. P. (2026). "RuNIE (Rust Native Inference Engine): An Open-Source Architecture for Autonomous Bioinformatics Pipeline Synthesis, Blinded Validation, and Bayesian Scope-Depth Auditing." CRV Science Labs Preprints (PREPRINT-01). https://doi.org/10.5281/zenodo.23969275

[2] Meredith, R. W., Janečka, J. E., Gatesy, J., Ryder, O. A., Fisher, C. A., Teeling, E. C., ... & Murphy, W. J. (2011). "Impacts of the Cretaceous Terrestrial Revolution and KPg extinction on mammal diversification." Science, 334(6055), 521-524. https://doi.org/10.1126/science.1211028

[3] Álvarez-Carretero, S., Tamuri, A. U., Battini, M., Nascimento, F. F., Carlisle, E., Asher, R. J., ... & dos Reis, M. (2022). "A species-level timeline of mammal evolution integrating phylogenomic data." Nature, 602(7896), 263-267. https://doi.org/10.1038/s41586-021-04341-1

[4] Foley, N. M., Mason, V. C., Harris, A. J., Bredemeyer, K. R., Damas, J., Lewin, H. A., ... & Murphy, W. J. (2023). "A genomic timescale for placental mammal evolution." Science, 380(6643), eabl8189. https://doi.org/10.1126/science.abl8189

[5] O'Leary, M. A., Bloch, J. I., Flynn, J. J., Gaudin, T. J., Giallombardo, A., Giannini, N. P., ... & Cirranello, A. L. (2013). "The placental mammal ancestor and the post-K-Pg radiation of placentals." Science, 339(6120), 662-667. https://doi.org/10.1126/science.1229237

[6] dos Reis, M., Inoue, J., Hasegawa, M., Asher, R. J., Donoghue, P. C. J., & Yang, Z. (2012). "Phylogenomic datasets provide both precision and accuracy in estimating the timescale of placental mammal phylogeny." Proceedings of the Royal Society B, 279(1742), 3491-3500. https://doi.org/10.1098/rspb.2012.0683

[7] Upham, N. S., Esselstyn, J. A., & Jetz, W. (2019). "Inferring the mammal tree: species-level sets of phylogenies for questions in ecology, evolution, and conservation." PLOS Biology, 17(12), e3000494. https://doi.org/10.1371/journal.pbio.3000494

[8] Liu, L., Zhang, J., Rheindt, F. E., Lei, F., Qu, Y., Wang, Y., ... & Edwards, S. V. (2017). "Genomic evidence reveals a radiation of placental mammals uninterrupted by the KPg boundary." Proceedings of the National Academy of Sciences, 114(35), E7282-E7290. https://doi.org/10.1073/pnas.1616744114

[9] Springer, M. S., Emerling, C. A., Meredith, R. W., Janečka, J. E., Eizirik, E., & Murphy, W. J. (2017). "Waking the undead: implications of a soft explosive model for placental mammal origins." Journal of Heredity, 108(1), 62-77. https://doi.org/10.1093/jhered/esw052

[10] Luo, Z. X., Yuan, C. X., Meng, Q. J., & Ji, Q. (2011). "A Jurassic eutherian mammal and divergence of marsupials and placentals." Nature, 476(7361), 442-445. https://doi.org/10.1038/nature10291

[11] Benton, M. J., Donoghue, P. C. J., Asher, R. J., Friedman, M., Near, T. J., & Vinther, J. (2015). "Constraints on the timescale of animal evolutionary history." Palaeontologia Electronica, 18.1.1FC, 1-106. https://doi.org/10.26879/424

[12] Crawford, N. G., Faircloth, B. C., McCormack, J. E., Brumfield, R. T., Winker, K., & Glenn, T. C. (2012). "More than 1000 ultraconserved elements provide evidence that turtles are sister to archosaurs." Biology Letters, 8(5), 783-786. https://doi.org/10.1098/rsbl.2012.0331

[13] Wilson Mantilla, G. P., Chester, S. G. B., Clemens, W. A., Moore, J. R., Sprain, C. J., Hovatter, B. T., ... & Renne, P. R. (2021). "Earliest Palaeocene purgatoriids and the initial radiation of stem primates." Royal Society Open Science, 8(2), 210050. https://doi.org/10.1098/rsos.210050

[14] Teeling, E. C., Springer, M. S., Madsen, O., Bates, P., O'Brien, S. J., & Murphy, W. J. (2005). "A molecular phylogeny for bats illuminates biogeography and the fossil record." Science, 307(5709), 580-584. https://doi.org/10.1126/science.1105113

[15] Gatesy, J., Geisler, J. H., Chang, J., Buell, C., Berta, A., Meredith, R. W., ... & McGowen, M. R. (2013). "A phylogenetic blueprint for a modern whale." Molecular Phylogenetics and Evolution, 66(2), 479-506. https://doi.org/10.1016/j.ympev.2012.10.022

[16] Beck, R. M. D. (2008). "A dated phylogeny of marsupials using a molecular supermatrix and multiple fossil constraints." Journal of Mammalogy, 89(1), 175-189. https://doi.org/10.1644/06-MAMM-A-437.1

[17] dos Reis, M., Gunnell, G. F., Barba-Montoya, J., Wilkins, A., Yang, Z., & Yoder, A. D. (2018). "Using phylogenomic data to explore the effects of relaxed clocks and calibration strategies on divergence time estimation: primates as a test case." Systematic Biology, 67(4), 594-615. https://doi.org/10.1093/sysbio/syy001

[18] Li, Y., Liu, Z., Shi, P., & Zhang, J. (2010). "The hearing gene Prestin unites echolocating bats and whales." Current Biology, 20(2), R55-R56. https://doi.org/10.1016/j.cub.2009.11.042

[19] Liu, Y., Cotton, J. A., Shen, B., Han, X., Rossiter, S. J., & Zhang, S. (2010). "Convergent sequence evolution between echolocating bats and dolphins." Current Biology, 20(2), R53-R54. https://doi.org/10.1016/j.cub.2009.11.058

[20] Bavi, N., Clark, M. D., Contreras, G. F., Shen, R., Reddy, B. G., Milewski, W., & Perozo, E. (2021). "The conformational cycle of prestin underlies outer-hair-cell electromotility (PDB: 7LH2)." Nature, 600(7889), 553-558. https://doi.org/10.1038/s41586-021-04152-4

[21] Meredith, R. W., Gatesy, J., Murphy, W. J., Ryder, O. A., & Springer, M. S. (2009). "Molecular decay of the tooth gene Enamelin (ENAM) mirrors the loss of enamel in the fossil record of placental mammals." PLOS Genetics, 5(9), e1000634. https://doi.org/10.1371/journal.pgen.1000634

[22] Nabholz, B., Glémin, S., & Galtier, N. (2008). "Strong variations of mitochondrial mutation rate across mammals—the longevity hypothesis." Molecular Biology and Evolution, 25(1), 120-130. https://doi.org/10.1093/molbev/msm248

[23] Romiguier, J., Ranwez, V., Delsuc, F., Galtier, N., & Douzery, E. J. P. (2013). "Less is more in mammalian phylogenomics: AT-rich genes minimize tree conflicts and unravel the root of placental mammals." Molecular Biology and Evolution, 30(9), 2134-2144. https://doi.org/10.1093/molbev/mst116

[24] Zhang, C., Rabiee, M., Sayyari, E., & Mirarab, S. (2018). "ASTRAL-III: polynomial time species tree reconstruction from partially resolved gene trees." BMC Bioinformatics, 19(6), 153. https://doi.org/10.1186/s12859-018-2129-y

[25] Katoh, K., & Standley, D. M. (2013). "MAFFT multiple sequence alignment software version 7: improvements in performance and usability." Molecular Biology and Evolution, 30(4), 772-780. https://doi.org/10.1093/molbev/mst010

[26] Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von Haeseler, A., & Lanfear, R. (2020). "IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era." Molecular Biology and Evolution, 37(5), 1530-1534. https://doi.org/10.1093/molbev/msaa015

[27] To, T. H., Jung, M., Lycett, S., & Gascuel, O. (2016). "Fast dating using least-squares criteria and algorithms." Systematic Biology, 65(1), 82-97. https://doi.org/10.1093/sysbio/syv068

[28] Yang, Z. (2007). "PAML 4: phylogenetic analysis by maximum likelihood." Molecular Biology and Evolution, 24(8), 1586-1591. https://doi.org/10.1093/molbev/msm088

Supplemental Table S1. Complete 64-Species Mammalia, Sauropsida & Lissamphibia Mitogenome Roster (Sectors A:T001–G:T064), Body Mass, Compositional Heterogeneity & Solved Branch Lengths

Supplemental Table S1. Complete 64-Species Tetrapod Mitogenome Inventory (A:T001–G:T064), NCBI RefSeq Accessions, Body Mass, GC% / AT3%, Taxon RCFV & Solved Patristic Branch Lengths

Sector:Tip ID	Species	Order & Family	NCBI RefSeq Accession	Adult Body Mass	Mitogenome Length (bp)	GC% / AT3%	Taxon RCFV	Terminal (Cum.) Branch Len	Parent Node (UFBoot %)
A:T001	*Homo sapiens*	Primates: Hominidae	NC_012920.1	62.0 kg	16,569	44.36% / 56.32%	0.0975	0.04042 (0.7562)	N14 (100/100)
A:T002	*Pan troglodytes*	Primates: Hominidae	NC_001643.1	45.0 kg	16,554	43.69% / 56.54%	0.0881	0.04153 (0.7573)	N14 (100/100)
A:T003	*Macaca mulatta*	Primates: Cercopithecidae	NC_005943.1	7.7 kg	16,564	43.23% / 56.58%	0.0802	0.16777 (0.8002)	N13 (100/100)
A:T004	*Callithrix pygmaea*	Primates: Callitrichidae	NC_021942.1	0.120 kg	16,520	39.68% / 61.55%	0.0093	0.25426 (0.7990)	N12 (100/100)
A:T005	*Carlito syrichta*	Primates: Tarsiidae	NC_012774.1	0.120 kg	17,004	39.14% / 58.82%	0.0227	0.27431 (0.7303)	N11 (100/100)
A:T006	*Lemur catta*	Primates: Lemuridae	NC_004025.1	2.2 kg	17,036	38.09% / 64.05%	0.0354	0.26397 (0.7134)	N10 (100/100)
A:T007	*Galeopterus variegatus*	Dermoptera: Cynocephalidae	NC_004031.1	1.3 kg	16,748	43.21% / 56.09%	0.0670	0.26249 (0.6985)	N09 (100/100)
A:T008	*Tupaia belangeri*	Scandentia: Tupaiidae	NC_002521.1	0.150 kg	16,754	40.75% / 59.42%	0.0181	0.28271 (0.6990)	N08 (100/100)
A:T009	*Mus musculus*	Rodentia: Muridae	NC_005089.1	0.020 kg	16,299	36.74% / 60.08%	0.0623	0.14253 (0.8332)	N18 (100/100)
A:T010	*Rattus norvegicus*	Rodentia: Muridae	NC_001665.2	0.300 kg	16,313	38.57% / 62.88%	0.0258	0.11424 (0.8049)	N18 (100/100)
A:T011	*Cavia porcellus*	Rodentia: Caviidae	NC_000884.1	0.750 kg	16,801	39.31% / 61.57%	0.0456	0.33887 (0.8094)	N17 (100/100)
A:T012	*Sciurus vulgaris*	Rodentia: Sciuridae	NC_002369.1	0.330 kg	16,507	37.03% / 61.54%	0.0684	0.27889 (0.7306)	N16 (100/100)
A:T013	*Oryctolagus cuniculus*	Lagomorpha: Leporidae	NC_001913.1	1.8 kg	17,245	40.21% / 59.85%	0.0244	0.26325 (0.6987)	N15 (100/100)
B:T014	*Balaenoptera musculus*	Cetartiodactyla: Balaenopteridae (Mysticeti)	NC_001601.1	136000.0 kg	16,402	40.65% / 59.74%	0.0254	0.07135 (0.7007)	N29 (100/100)
B:T015	*Tursiops truncatus*	Cetartiodactyla: Delphinidae (Odontoceti)	NC_012059.1	260.0 kg	16,388	38.92% / 59.96%	0.0188	0.10571 (0.7459)	N30 (100/100)
B:T016	*Physeter catodon*	Cetartiodactyla: Physeteridae (Odontoceti)	NC_002503.2	35000.0 kg	16,428	43.06% / 56.12%	0.0649	0.11146 (0.7516)	N30 (100/100)
B:T017	*Hippopotamus amphibius*	Cetartiodactyla: Hippopotamidae	NC_000889.1	1500.0 kg	16,407	42.56% / 56.39%	0.0549	0.15349 (0.6958)	N28 (100/100)
B:T018	*Bos taurus*	Cetartiodactyla: Bovidae (Ruminantia)	NC_006853.1	650.0 kg	16,338	39.42% / 59.92%	0.0143	0.07869 (0.6931)	N31 (100/100)
B:T019	*Ovis aries*	Cetartiodactyla: Bovidae (Ruminantia)	NC_001941.1	70.0 kg	16,616	38.94% / 62.21%	0.0193	0.08256 (0.6969)	N31 (100/100)
B:T020	*Sus scrofa*	Cetartiodactyla: Suidae (Suina)	NC_000845.1	130.0 kg	16,613	39.46% / 61.82%	0.0402	0.17495 (0.6760)	N26 (100/100)
B:T021	*Camelus bactrianus*	Cetartiodactyla: Camelidae (Tylopoda)	NC_009628.2	550.0 kg	16,659	41.92% / 58.49%	0.0413	0.22155 (0.7087)	N25 (100/100)
B:T022	*Equus caballus*	Perissodactyla: Equidae	NC_001640.1	450.0 kg	16,660	41.96% / 59.19%	0.0428	0.11396 (0.6235)	N32 (100/100)
B:T023	*Ceratotherium simum*	Perissodactyla: Rhinocerotidae	NC_001808.1	2100.0 kg	16,832	40.85% / 61.44%	0.0455	0.09567 (0.6260)	N33 (100/100)

B:T024	*Tapirus terrestris*	Perissodactyla: Tapiridae	NC_053962.1	`225.0 kg`	16,772	37.77% / 60.21%	0.0418	0.12823 (0.6585)	N33 (100/100)
B:T025	*Pteropus vampyrus*	Chiroptera: Pteropodidae (Megabat)	NC_026542.1	`1.0 kg`	16,554	41.52% / 57.87%	0.0405	0.17281 (0.6719)	N22 (100/100)
B:T026	*Rhinolophus ferrumequinum*	Chiroptera: Rhinolophidae (CF Sonar)	NC_016191.1	`0.023 kg`	16,839	42.77% / 56.64%	0.0582	0.14373 (0.6428)	N22 (100/100)
B:T027	*Myotis lucifugus*	Chiroptera: Vespertilionidae (FM Sonar)	NC_029849.1	`0.008 kg`	17,038	36.86% / 63.55%	0.0608	0.27015 (0.7423)	N21 (100/100)
C:T028	*Felis catus*	Carnivora: Felidae (Feliformia)	NC_001700.1	`4.2 kg`	17,009	40.33% / 57.84%	0.0137	0.05854 (0.6698)	N36 (100/100)
C:T029	*Panthera tigris*	Carnivora: Felidae (Feliformia)	NC_010642.1	`190.0 kg`	16,990	41.15% / 59.44%	0.0257	0.06335 (0.6746)	N36 (100/100)
C:T030	*Canis lupus*	Carnivora: Canidae (Caniformia)	NC_002008.4	`35.0 kg`	16,727	39.65% / 60.29%	0.0392	0.17425 (0.7131)	N37 (100/100)
C:T031	*Ursus maritimus*	Carnivora: Ursidae (Caniformia)	NC_003428.1	`380.0 kg`	17,017	41.36% / 59.01%	0.0538	0.08506 (0.7296)	N39 (100/100)
C:T032	*Phoca vitulina*	Carnivora: Phocidae (Pinnipedia)	NC_001325.1	`95.0 kg`	16,826	41.72% / 59.06%	0.0428	0.11937 (0.6870)	N38 (100/100)
C:T033	*Ailuropoda melanoleuca*	Carnivora: Ursidae (Giant Panda)	NC_009492.1	`105.0 kg`	16,805	38.79% / 61.69%	0.0706	0.09351 (0.7380)	N39 (100/100)
C:T034	*Manis pentadactyla*	Pholidota: Manidae (Edentulous Pangolin)	NC_016008.1	`4.5 kg`	16,578	43.81% / 55.63%	0.0872	0.27262 (0.7372)	N34 (100/100)
C:T035	*Erinaceus europaeus*	Eulipotyphla: Erinaceidae	NC_002080.2	`0.850 kg`	17,447	32.60% / 65.25%	0.1452	0.52316 (0.9617)	N40 (100/100)
C:T036	*Sorex araneus*	Eulipotyphla: Soricidae	NC_027963.1	`0.009 kg`	16,887	38.28% / 63.16%	0.0315	0.20130 (0.6854)	N41 (100/100)
C:T037	*Talpa europaea*	Eulipotyphla: Talpidae	NC_002391.1	`0.090 kg`	16,884	38.92% / 62.92%	0.0448	0.19711 (0.6812)	N41 (100/100)
D:T038	*Loxodonta africana*	Proboscidea: Elephantidae	NC_000934.1	`4800.0 kg`	16,866	38.75% / 62.38%	0.0235	0.02628 (0.7514)	N46 (100/100)
D:T039	*Elephas maximus*	Proboscidea: Elephantidae	NC_005129.2	`3800.0 kg`	16,902	38.72% / 62.99%	0.0229	0.01970 (0.7448)	N46 (100/100)
D:T040	*Trichechus manatus*	Sirenia: Trichechidae	NC_010302.1	`480.0 kg`	16,882	42.85% / 57.44%	0.0598	0.07823 (0.6477)	N47 (100/100)
D:T041	*Dugong dugon*	Sirenia: Dugongidae	NC_003314.1	`300.0 kg`	16,850	43.35% / 57.96%	0.0699	0.07567 (0.6451)	N47 (100/100)
D:T042	*Procavia capensis*	Hyracoidea: Procaviidae	NC_004919.1	`3.6 kg`	16,743	41.05% / 58.57%	0.0630	0.26354 (0.7094)	N44 (100/100)
D:T043	*Orycteropus afer*	Tubulidentata: Orycteropodidae	NC_002078.1	`55.0 kg`	16,816	38.05% / 63.28%	0.0363	0.24430 (0.6772)	N48 (100/100)
D:T044	*Echinops telfairi*	Afrosoricida: Tenrecidae	NC_002631.2	`0.140 kg`	16,549	37.00% / 64.09%	0.0810	0.34163 (0.8203)	N50 (100/100)
D:T045	*Chrysochloris asiatica*	Afrosoricida: Chrysochloridae	NC_004920.1	`0.045 kg`	16,652	36.56% / 62.95%	0.0660	0.22212 (0.7008)	N50 (100/100)
D:T046	*Macroscelides proboscideus*	Macroscelidea: Macroscelididae	NC_004026.1	`0.042 kg`	16,641	38.77% / 61.91%	0.0218	0.31371 (0.7604)	N49 (100/100)
D:T047	*Dasypus novemcinctus*	Cingulata: Dasypodidae	NC_001821.1	`4.2 kg`	17,056	38.86% / 60.79%	0.0348	0.23014 (0.6672)	N51 (100/100)
D:T048	*Choloepus didactylus*	Pilosa: Choloepodidae (Folivora)	NC_006924.1	`6.5 kg`	16,543	43.26% / 58.41%	0.0711	0.21291 (0.6745)	N52 (100/100)
D:T049	*Myrmecophaga tridactyla*	Pilosa: Myrmecophagidae (Vermilingua)	NC_028572.1	`33.0 kg`	16,498	39.10% / 62.05%	0.0283	0.23182 (0.6934)	N52 (100/100)
E:T050	*Monodelphis domestica*	Marsupialia: Didelphidae	NC_006299.1	`0.100 kg`	17,079	35.00% / 66.87%	0.0973	0.16560 (0.7161)	N54 (100/100)
E:T051	*Didelphis virginiana*	Marsupialia: Didelphidae	NC_001610.1	`2.8 kg`	17,084	33.18% / 64.47%	0.1335	0.16677 (0.7173)	N54 (100/100)

E:T052	*Phascolarctos cinereus*	Marsupialia: Phascolarctidae	NC_008133.1	`8.5 kg`	16,357	36.69% / 61.34%	0.0633	0.16917 (0.6975)	N56 (100/100)
E:T053	*Vombatus ursinus*	Marsupialia: Vombatidae	NC_003322.1	`26.0 kg`	16,996	39.61% / 61.11%	0.0275	0.11063 (0.6390)	N56 (100/100)
E:T054	*Sarcophilus harrisii*	Marsupialia: Dasyuridae	NC_018788.1	`8.0 kg`	17,117	37.37% / 62.19%	0.0574	0.25976 (0.7061)	N55 (100/100)
F:T055	*Ornithorhynchus anatinus*	Monotremata: Ornithorhynchidae	NC_000891.1	`1.5 kg`	17,019	37.13% / 64.97%	0.0833	0.13771 (0.6722)	N57 (100/100)
F:T056	*Tachyglossus aculeatus*	Monotremata: Tachyglossidae	NC_003321.1	`4.5 kg`	16,360	40.06% / 59.42%	0.0438	0.02907 (0.6554)	N58 (100/100)
F:T057	*Zaglossus bruijni*	Monotremata: Tachyglossidae	NC_006364.1	`9.0 kg`	16,367	39.92% / 57.95%	0.0471	0.02947 (0.6558)	N58 (100/100)
G:T058	*Gallus gallus*	Sauropsida: Aves (Phasianidae)	NC_001323.1	`1.8 kg`	16,775	45.96% / 52.69%	0.1220	0.18836 (0.5591)	N62 (100/100)
G:T059	*Taeniopygia guttata*	Sauropsida: Aves (Estrildidae)	NC_007897.1	`0.015 kg`	16,853	45.94% / 53.85%	0.1216	0.18266 (0.5534)	N62 (100/100)
G:T060	*Alligator mississippiensis*	Sauropsida: Archosauria (Alligatoridae)	NC_001922.1	`180.0 kg`	16,646	43.03% / 57.35%	0.0634	0.69733 (0.8720)	N61 (100/100)
G:T061	*Chelonia mydas*	Sauropsida: Testudines (Cheloniidae)	NC_000886.1	`140.0 kg`	16,497	39.45% / 60.03%	0.0762	0.32913 (0.4714)	N60 (100/100)
G:T062	*Anolis carolinensis*	Sauropsida: Lepidosauria (Dactyloidae)	NC_010972.2	`0.005 kg`	17,223	38.40% / 61.52%	0.0449	0.58080 (0.6639)	N59 (100/100)
G:T063	*Xenopus tropicalis*	Lissamphibia: Pipidae (Tetrapod Outgroup)	NC_006839.1	`0.025 kg`	17,610	42.50% / 58.19%	0.0528	0.30930 (0.3672)	N63 (100/100)
G:T064	*Ambystoma mexicanum*	Lissamphibia: Ambystomatidae (Tetrapod Outgroup)	NC_005797.1	`0.110 kg`	16,369	33.26% / 67.23%	0.1321	0.54473 (0.6027)	N63 (100/100)

🦴 **Generated by:** RuNIE Bayesian Pipeline Optimizer • Unmarked = Internal Tool • * = External Tool

Supplemental Table S2. Complete 32-Chain Multi-Model Combinatorial Optimization Sweep Across 8 Candidate Substitution Models × 4 RuNIE Regimes

Supplemental Table S2. Complete 32-Chain Multi-Model Combinatorial Sweep Across 8 Candidate Substitution Models × 4 RuNIE Regimes (L = 23,014 / 123,333 Aligned Columns)

Candidate Substitution Model (*IQ-TREE / Native Rust)	Regime 3 Hybrid RY3 ln L	Regime 3 Hybrid BIC	Hybrid CPU	Regime 1 Native RY3 ln L	Regime 1 Native BIC	Native CPU	Regime 2 External Raw ln L	Regime 2 External BIC	External CPU	Regime 4 0/5 Control ln L	Regime 4 0/5 Control BIC	0/5 CPU
Chain C1: JC / GTR+F+R2	-626162.08	1253579.65	0.48 s	-747448.86	1496456.84	25.35 s	-715013.49	1431282.46	0.38 s	-847844.19	1697153.72	28.22 s
Chain C2: K2P+G4 / K80+G4	-548372.62	1098020.82	5.38 s	-761275.92	1524017.18	26.02 s	-621278.64	1243832.86	10.23 s	-860291.54	1722142.19	26.04 s
Chain C3: HKY+F+I+G4 / HKY+F+I+G4	-516522.18	1034360.10	135.75 s	-775262.75	1552037.72	25.84 s	-597535.99	1196387.72	170.02 s	-880040.37	1761604.68	27.92 s
Chain C4: TN+F+I+G4 / TN+F+I+G4	-516461.17	1034248.13	149.58 s	-776293.58	1554111.11	26.19 s	-596573.00	1194471.79	187.42 s	-880266.02	1762044.27	28.19 s
Chain C5: TIM2+F+I+G4 / TIM2+F+I+G4	-515317.31	1031970.45	160.11 s	-781802.63	1565140.92	26.02 s	-595218.13	1191772.09	256.83 s	-884122.44	1769780.55	28.15 s
Chain C6: GTR+F+G4 / GTR+F+I+G4	-516449.67	1034245.22	25.20 s	-782617.46	1566794.04	26.21 s	-596737.11	1194820.10	39.53 s	-884761.00	1771081.11	28.15 s
Chain C7: GTR+F+I+G4 / GTR+F+I+G4 (Fast)	-514504.01	1030363.95	220.31 s	-782929.76	1567418.62	16.10 s	-595165.21	1191686.34	293.91 s	-884942.35	1771443.80	17.14 s
Chain C8: GTR+F+R4 / JC	-514594.23	1030584.56	63.55 s	-815920.97	1633283.83	23.25 s	-595619.52	1192635.13	76.02 s	-912500.33	1826442.54	24.55 s

🔧 **Generated by:** RuNIE Bayesian Pipeline Optimizer • Unmarked = Internal Tool • * = External Tool

Supplemental Table S3. Complete 52-Record Matched Nuclear Coding mRNA Inventory (N001–N052 Across SLC26A5, ENAM, RAG1 & BDNF)

Supplemental Table S3. Complete 52-Sequence Nuclear Coding mRNA Inventory Across 13 Matched Tetrapods (L = 36,010 Aligned Columns in *MAFFT / 47,403 in gotoh-msa)

Nuclear ID (Mito Tip)	Species	Orthologous Nuclear Locus & Functional Role	NCBI Accession	mRNA Length (bp)	GC% / AT3%	Transcript RCFV	Species Term. (Cum.) Branch	Parent Node (UFBoot %)
N001 (`A:T001`)	*Homo sapiens*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	NM_001167962.2	2,623	46.32% / 58.35%	0.0168	0.06927 (0.3389)	N05 (94/94)
N002 (`A:T009`)	*Mus musculus*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	NM_001289787.1	4,430	44.97% / 52.51%	0.0303	0.17360 (0.4569)	N06 (93/93)
N003 (`A:T013`)	*Oryctolagus cuniculus*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	NM_001142261.1	2,235	52.84% / 33.15%	0.0484	0.09785 (0.3811)	N06 (93/93)
N004 (`B:T015`)	*Tursiops truncatus*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	XM_004320267.4	7,278	40.74% / 58.90%	0.0726	0.03506 (0.3552)	N10 (100/100)
N005 (`B:T018`)	*Bos taurus*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	NM_001192878.2	2,226	45.64% / 53.23%	0.0236	0.05043 (0.3706)	N10 (100/100)
N006 (`B:T022`)	*Equus caballus*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	XM_070265357.1	2,462	51.18% / 56.83%	0.0318	0.17332 (0.4523)	N08 (100/100)
N007 (`B:T025`)	*Pteropus vampyrus*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	XM_011365196.2	2,841	43.33% / 52.48%	0.0467	0.04801 (0.3319)	N11 (100/100)
N008 (`B:T027`)	*Myotis lucifugus*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	XM_023748393.1	2,483	46.48% / 61.79%	0.0152	0.05040 (0.3343)	N11 (100/100)
N009 (`C:T030`)	*Canis lupus familiaris*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	XM_038423862.1	4,606	42.57% / 62.35%	0.0543	0.09176 (0.3769)	N09 (91/91)
N010 (`D:T038`)	*Loxodonta africana*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	XM_064289890.1	1,716	44.81% / 54.55%	0.0319	0.09356 (0.3537)	N04 (55/55)
N011 (`E:T050`)	*Monodelphis domestica*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	XM_007504069.4	7,037	39.90% / 61.54%	0.0810	0.26654 (0.4049)	N02 (100/100)
N012 (`G:T060`)	*Alligator mississippiensis*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	XM_019487482.2	5,158	44.86% / 56.02%	0.0314	0.39571 (0.3957)	N01 (100/100)
N013 (`G:T063`)	*Xenopus tropicalis*	***SLC26A5*** — Prestin Outer Hair Cell Cochlear Motor (PDB: 7LH2)	XM_031899020.1	5,872	38.73% / 63.41%	0.0927	0.49596 (0.4960)	N01 (100/100)
N014 (`A:T001`)	*Homo sapiens*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	NM_001368133.1	4,943	41.43% / 57.92%	0.0657	0.06927 (0.3389)	N05 (94/94)
N015 (`A:T009`)	*Mus musculus*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	NM_017468.3	5,490	42.95% / 51.37%	0.0505	0.17360 (0.4569)	N06 (93/93)
N016 (`A:T013`)	*Oryctolagus cuniculus*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	XM_002717035.4	4,320	43.26% / 65.00%	0.0474	0.09785 (0.3811)	N06 (93/93)
N017 (`B:T015`)	*Tursiops truncatus*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	XM_033857118.1	2,912	44.09% / 53.09%	0.0391	0.03506 (0.3552)	N10 (100/100)
N018 (`B:T018`)	*Bos taurus*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	XM_002688339.6	5,512	40.75% / 56.51%	0.0725	0.05043 (0.3706)	N10 (100/100)
N019 (`B:T022`)	*Equus caballus*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	XM_001487894.3	3,763	45.95% / 62.84%	0.0205	0.17332 (0.4523)	N08 (100/100)

N020 (`B:T025`)	*Pteropus vampyrus*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	XM_011360502.1	3,459	46.08% / 63.31%	0.0192	0.04801 (0.3319)	N11 (100/100)
N021 (`B:T027`)	*Myotis lucifugus*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	XM_006094414.1	3,450	46.70% / 62.52%	0.0130	0.05040 (0.3343)	N11 (100/100)
N022 (`C:T030`)	*Canis lupus familiaris*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	XM_038684621.1	3,447	46.07% / 63.97%	0.0193	0.09176 (0.3769)	N09 (91/91)
N023 (`D:T038`)	*Loxodonta africana*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	XM_003414167.3	6,723	42.51% / 62.07%	0.0549	0.09356 (0.3537)	N04 (55/55)
N024 (`E:T050`)	*Monodelphis domestica*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	XM_056803382.2	3,906	44.96% / 61.06%	0.0304	0.26654 (0.4049)	N02 (100/100)
N025 (`G:T060`)	*Alligator mississippiensis*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	XM_019490832.2	6,169	46.56% / 54.18%	0.0144	0.39571 (0.3957)	N01 (100/100)
N026 (`G:T063`)	*Xenopus tropicalis*	***ENAM*** — Enamelin Secreted Dental Matrix Protein	NM_001145743.1	3,440	38.81% / 52.62%	0.0919	0.49596 (0.4960)	N01 (100/100)
N027 (`A:T001`)	*Homo sapiens*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	XM_054369590.1	7,759	41.82% / 56.65%	0.0618	0.06927 (0.3389)	N05 (94/94)
N028 (`A:T009`)	*Mus musculus*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	NM_009019.2	6,669	44.02% / 54.84%	0.0398	0.17360 (0.4569)	N06 (93/93)
N029 (`A:T013`)	*Oryctolagus cuniculus*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	XM_008268573.5	6,668	42.32% / 56.71%	0.0568	0.09785 (0.3811)	N06 (93/93)
N030 (`B:T015`)	*Tursiops truncatus*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	XM_004329819.4	3,146	52.07% / 37.12%	0.0407	0.03506 (0.3552)	N10 (100/100)
N031 (`B:T018`)	*Bos taurus*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	XM_024975902.2	7,349	43.34% / 56.31%	0.0466	0.05043 (0.3706)	N10 (100/100)
N032 (`B:T022`)	*Equus caballus*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	NM_001256901.1	3,129	51.36% / 38.35%	0.0336	0.17332 (0.4523)	N08 (100/100)
N033 (`B:T025`)	*Pteropus vampyrus*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	XM_011356837.1	4,413	46.02% / 61.79%	0.0198	0.04801 (0.3319)	N11 (100/100)
N034 (`B:T027`)	*Myotis lucifugus*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	XM_023763518.1	3,132	52.91% / 33.91%	0.0491	0.05040 (0.3343)	N11 (100/100)
N035 (`C:T030`)	*Canis lupus familiaris*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	XM_038423502.1	3,132	61.33% / 18.58%	0.1333	0.09176 (0.3769)	N09 (91/91)
N036 (`D:T038`)	*Loxodonta africana*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	XM_003412274.3	3,268	48.38% / 45.55%	0.0038	0.09356 (0.3537)	N04 (55/55)
N037 (`E:T050`)	*Monodelphis domestica*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	XM_056803782.2	3,714	47.47% / 46.45%	0.0053	0.26654 (0.4049)	N02 (100/100)
N038 (`G:T060`)	*Alligator mississippiensis*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	XM_006275935.3	7,051	37.95% / 63.06%	0.1005	0.39571 (0.3957)	N01 (100/100)
N039 (`G:T063`)	*Xenopus tropicalis*	***RAG1*** — V(D)J Recombination Activating Gene 1 Anchor	XM_002937292.4	4,840	45.41% / 50.22%	0.0259	0.49596 (0.4960)	N01 (100/100)
N040 (`A:T001`)	*Homo sapiens*	***BDNF*** — Brain-Derived Neurotrophic Factor	NM_170734.4	3,979	41.74% / 60.56%	0.0626	0.06927 (0.3389)	N05 (94/94)
N041 (`A:T009`)	*Mus musculus*	***BDNF*** — Brain-Derived Neurotrophic Factor	NM_001316310.1	3,950	40.20% / 57.52%	0.0780	0.17360 (0.4569)	N06 (93/93)
N042 (`A:T013`)	*Oryctolagus cuniculus*	***BDNF*** — Brain-Derived Neurotrophic Factor	XM_089310241.1	1,562	44.24% / 60.58%	0.0376	0.09785 (0.3811)	N06 (93/93)
N043 (`B:T015`)	*Tursiops truncatus*	***BDNF*** — Brain-Derived Neurotrophic Factor	XM_019947884.3	1,580	50.51% / 41.63%	0.0251	0.03506 (0.3552)	N10 (100/100)

N044 (`B:T018`)	*Bos taurus*	***BDNF*** — Brain-Derived Neurotrophic Factor	NM_001046607.2	1,206	51.33% / 38.06%	0.0333	0.05043 (0.3706)	N10 (100/100)
N045 (`B:T022`)	*Equus caballus*	***BDNF*** — Brain-Derived Neurotrophic Factor	NM_001163828.2	3,187	56.92% / 56.69%	0.0892	0.17332 (0.4523)	N08 (100/100)
N046 (`B:T025`)	*Pteropus vampyrus*	***BDNF*** — Brain-Derived Neurotrophic Factor	XM_011362569.2	4,028	41.73% / 59.24%	0.0627	0.04801 (0.3319)	N11 (100/100)
N047 (`B:T027`)	*Myotis lucifugus*	***BDNF*** — Brain-Derived Neurotrophic Factor	XM_023756376.1	1,605	46.98% / 60.00%	0.0102	0.05040 (0.3343)	N11 (100/100)
N048 (`C:T030`)	*Canis lupus familiaris*	***BDNF*** — Brain-Derived Neurotrophic Factor	NM_001002975.1	1,146	46.07% / 61.78%	0.0193	0.09176 (0.3769)	N09 (91/91)
N049 (`D:T038`)	*Loxodonta africana*	***BDNF*** — Brain-Derived Neurotrophic Factor	XM_023550772.2	4,009	40.00% / 60.22%	0.0800	0.09356 (0.3537)	N04 (55/55)
N050 (`E:T050`)	*Monodelphis domestica*	***BDNF*** — Brain-Derived Neurotrophic Factor	XM_085329484.1	1,767	43.97% / 54.84%	0.0403	0.26654 (0.4049)	N02 (100/100)
N051 (`G:T060`)	*Alligator mississippiensis*	***BDNF*** — Brain-Derived Neurotrophic Factor	XM_019476631.2	4,199	37.56% / 64.26%	0.1044	0.39571 (0.3957)	N01 (100/100)
N052 (`G:T063`)	*Xenopus tropicalis*	***BDNF*** — Brain-Derived Neurotrophic Factor	NM_001097298.1	865	43.93% / 54.17%	0.0407	0.49596 (0.4960)	N01 (100/100)

🔧 **Generated by:** RuNIE Bayesian Pipeline Optimizer • Unmarked = Internal Tool • * = External Tool