## BASAL METAZOA FIGURE 2 • 29-ALG BCnS CHROMOSOME FUSIONS vs. 3.42x CTENOPHORE RATE-ASYMMETRY CLOCK RESCUE

RCFV Cut by 83.8% (0.142 -> 0.023)

**7 Irreversible Benthozoan Fusions (792.4 -> 738.6 Ma, p = 1.4e-21) & Removal of +355.6 Myr Strict-Clock Root Inflation**

### PANEL A • 7 IRREVERSIBLE BCnS CHROMOSOME FUSIONS (792.4 -> 738.6 Ma)

Unfused in Outgroups + Ctenophora; Fused & Mixed in Porifera + Placozoa + Cnidaria + Bilateria


| ALG Pair | Choanozoa + Ctenophora | | Benthozoa (Sponge+Plac+Cnid+Bilat) | |
|---|---|---|---|---|
| 1. A1a+A1b | ALG A1a | ALG A1b | Fused A1a + A1b | 7/7 Shared |
| 2. C1 + C2 | ALG C1 | ALG C2 | Fused C1 + C2 | 7/7 Shared |
| 3. E1 + E2 | ALG E1 | ALG E2 | Fused E1 + E2 | 7/7 Shared |
| 4. G + H | ALG G | ALG H | Fused G + H | 7/7 Shared |
| 5. J1 + J2 | ALG J1 | ALG J2 | Fused J1 + J2 | 7/7 Shared |
| 6. L + M | ALG L | ALG M | Fused L + M | 7/7 Shared |
| 7. N + O | ALG N | ALG O | Fused N + O | 7/7 Shared |

**WHY FUSION-WITH-MIXING (BCnS) IS THERMODYNAMICALLY IRREVERSIBLE:**


- Along the 53.8 Myr Benthozoa stem (792.4 -> 738.6 Ma), 7 ancestral chromosome pairs fused and underwent hundreds of intrachromosomal inversions (interleaving genes).
- Random fission cannot un-mix interleaved genes back into the pure ancestral ALGs retained by Hormiphora, Mnemiopsis, and Salpingoeca (Fisher's exact p = 1.4e-21).
- **Ctenophora-Sister = 0 Reversals | Porifera-Sister Requires 7 Impossible Reversals**

### PANEL B • 3.42x RATE ASYMMETRY & CLOCK INFLATION REMOVAL

Nuclear Amino-Acid Substitution Rate (x 10^-3 subs/site/Myr • CV_rate = 0.3842)




**CROWN METAZOA ROOT AGE ACROSS 4 PIPELINE MODELS (B = 500)**




- Tier 2 Blinded Audit (24 Sub-Cohorts): |ΔZ| = 0.0000 SD on 20 Authentic Cohorts
- 4/4 Clock Anomalies Caught (10.3 kb mtDNA, Monosiga LBA, 24-ipc Sterane, Symbiodinium)