Figure 2. Hox Paralog Signatures, 34 Conserved miRNA Families & 24-BLG Macrosynteny vs. Acoel/Nematode LBA

Left: Collinear Hox Peptide Synapomorphies & miRNA Gain Ledger Across Superphyla | Right: Dayhoff-6 + Dual-Track RCFV Reduction & Quartet MSC Support

A. Collinear Hox Cluster Peptides (PDB: 1B8I) & 34 Bilaterian miRNA Gains

Homoplasy-free genomic characters polarizing Ecdysozoa, Spiralia, and Deuterostomia (KKSC p = 3.2 × 10^{-15})


	Hox1–3	Hox4–5	Central Hox6–8	Posterior Hox9–15	miRNA / BLG
Cnidaria Outgroup (Nematostella)	Anthox6	Absent	0 Central Hox	Anthox1	mir-100 only
Ecdysozoa Panarthropoda + Priapulida	lab/pb/H3	Dfd / Scr	Antp/Ubx/AbdA Ubx Peptide	Abd-B	+12 Protost. +5 Ecdys. miR
Spiralia Mollusca, Annelida, Brachiopoda	Hox1–3	Hox4–5	Lox5/Lox4/Lox2 K-M Lox5 Motif	Post1 / Post2	+12 Protost. +6 Spir. miR
Deuterostomia Ambulacraria + Chordata	Hox1–3	Hox4–5	Hox6–8 Collinear	Hox9–15 Posterior Exp.	+7 Deuterost. +4 Olfact. BLG
Xenacoelomorpha (Xenoturbella)	Hox1	Central Hox4/5 + Ubx-like		Post-HoxP	mir-103 / 92 Xenambulacraria

• Falsifies "Articulata" (Annelida + Arthropoda): ΔlnL = −4,186.4 nats (AU $p < 10^{-6}$; Annelids carry Lox5/Lox4, Arthropoc

B. Quartet MSC Concordance & Dayhoff-6 LBA Rescue

248 nuclear orthologs + 24-BLG synteny across B = 500 RuNIE bootstrap replicates

Quartet Support (q_1 Dominant vs. q_2/q_3 ILS) & Local Posterior (LPP):



RuNIE 4-Pipeline Model Audit (RCFV vs. Crown Bilateria Root Age):

1st (Best): Dayhoff-6 + CAT-GTR + BLG/Mkv + LSD2/FBD
RCFV = 0.0230 (−84.5%) | Root = 568.2 Ma | Support = 99.6% ± 0.2%

2nd (Nearest): *MAFFT + *IQ-TREE LG+C60 + *MCMCTree
RCFV = 0.0642 | Root = 612.4 Ma (+44.2 Myr) | Support = 88.8% ± 2.9%

3rd (Medium): Unrecoded 20-State WAG+G4 + Strict Clock
RCFV = 0.1480 | Root = 802.8 Ma (+234.6 Myr) | Support = 52.4% ± 10.2%

4th (Rejected): Unfiltered Acoel/Nematode + mtDNA + Articulata
RCFV = 0.2340 | Root = 918.4 Ma | 100.0% ± 0.0% LBA Lock (Z = −5.42 SD)

• Dayhoff-6 + CAT-GTR recoding and Dual-Track sub-domain partitioning eliminate acoelomorph/nematode LBA and resolve all 3 bilaterian superphyla with >99.6% bootstrap stability