
Figure 3. The Cambrian Stage 2–3 Ecological Arms Race: Biomineralization Co-Option & Visual Opsin Surge

Top: Coupled Phenotypic (3.86×) and Molecular Clock (2.44×) Rate Surge Across 539–508 Ma | Bottom: α-Carbonic Anhydrase (PDB: 1CA2) & Opsin (PDB: 1F88) Selection

A. Coupled Phenotypic Character Rate (σ^2_morph, Mkv) & Molecular Clock Rate Surge Across the Ediacaran–Cambrian Transition



B. Parallel Biomineralization Toolkit Co-Option (PDB: 1CA2)

Conserved Track A α-Carbonic Anhydrase Core (TM = 0.908) + Lineage-Specific Acidic IDR Track B

- Mollusca (Nacrein): CA Zn^{2+} Core + (Gly-X-Asn)n Acidic Repeat → Aragonite/Calcite Shell
- Brachiopoda (Lingula / †Heliomedusa): CA Core + Asp-Rich IDR → Ca-Phosphate / Calcite
- Euarthropoda (Trilobita / Crustacea): Chitin Synthase CS1 + Calcified Cuticle Matrix
- Echinodermata & Vertebrata: SM50 Spicule Matrix / BMP2/4 + Enamel/Dentin (ω_2 = 3.62)

C. Visual Opsin Divergence (PDB: 1F88 / 2Z73) & Predatory Arms Race

Retinal Schiff-Base Lys296 Pocket Split: Rhabdomeric r-Opsin vs. Ciliary c-Opsin

Protostome Rhabdomeric r-Opsin (Gq-PLC-TRP Cascade, PDB: 2Z73)

- Counterion Tyr111/Glu181 • High-acuity compound eyes in †Anomalocaris (>16,000 lenses)
- Coupled with raptorial frontal appendages & Distal-less (Dll) limb patterning (ω_2 = 3.4

Deuterostome Ciliary c-Opsin (Gt-PDE Cascade, PDB: 1F88)

- Counterion Glu113 stabilizes protonated Lys296 Schiff base ($\Delta\lambda$_max spectral tuning)
- Paired camera eyes + Pax6/Rx neural crest in †Haikouichthys (518 Ma) & †Metaspriggina

• Ediacaran genetic toolkit assembly (568–540 Ma) preceded the synchronous Early Cambrian (539–514 Ma) biomineralization and visual predation burst