
HUMAN OUT-OF-AFRICA FIGURE 2 • HAPLOTYPE MINIMUM SPANNING NETWORK (MSN) & k-CORE HUB TOPOLOGY

Reticulation Cut by 87.4% (Z = +5.28 SD)

East African L3* k = 5 Connector Hub (C_B = 0.518) Bridging Sub-Saharan L0-L2 to Eurasian M/N/R vs. D-Loop Homoplasy


PANEL A • PARTITIONED 13-GENE OXPHOS + rRNA/tRNA MINIMUM SPANNING NETWORK (R_retic = 0.042)

All non-African mitogenomes radiate strictly through the East African L3* founder hub via M (10400T) and N (10873T)



PANEL B • UNPARTITIONED D-LOOP HOMOPLASY

Recurrent HVS-I 16189C / 16519C / 16223T Loops



QUARANTINED TIER-2 BLINDED ANOMALIES (4/4)

1. **BLIND-OOAMT-05 (NT_006576.16 Chr5 NUMT)**
   dN/dS = 0.884, 2 stops • D_ctx = 5.52 SD (Quarantined)
2. **BLIND-OOAMT-11 (HVS-I 16189C Chimera)**
   Poly-C slippage loop • D_ctx = 5.18 SD (Down-weighted)
3. **BLIND-OOAMT-17 (5' C->T aDNA Damage)**
   False 10400T/10873T call • D_ctx = 4.92 SD (Trimmed)
4. **BLIND-OOAMT-22 (Recurrent 9-bp Deletion)**
   L2 vs. B4indel homoplasy • D_ctx = 4.74 SD (Phased)

East African L3* Founder Hub (k = 5, C_B = 0.518) | M (10400T) & N (10873T) Primary Out-of-Africa Bridges | Quarantined D-Loop / NUMT Homoplasy