
Figure 2: Frequentist Supermatrix Variance Collapse vs. RuNIE 3-Tier Scope x 0-5 Beaker Bayesian Calibration

A. Supermatrix Sampling-Variance Collapse (sigma^2 ~ 1/L)

Unpartitioned Concatenation (RCFV = 0.118, epsilon_sys > 0)



B. Orthogonal Scope Tier x Bayesian Posterior Beaker Matrix

Mb (L)

Tier = Epistemic Scope | Beakers = Posterior Signal Strength Z_post

```
0 Beakers (Reject): Z < 2.50 SD or Fails K_min (54 Logged Controls)
1 Beaker  (Note):   Single-locus baseline / MIBiG Jaccard >= 0.80
2 Beakers (Valid.): >= 5 phased loci / r >= 0.9906 vs Reference
3 Beakers (Synth.): >= 25 taxa / >= 250 kb comparative synthesis
4 Beakers (Strong): >= 250 kb, >= 3 chained tools, Z >= 3.50 SD
5 Beakers (Signif): >= 1.0 Mb, >= 50 taxa, Z >= 4.50 SD + H4 Test
```

Bayesian Prior -> Posterior Adjustment Rule:

B_post = f(ln BF_10, Z_post, D_ctx, H4_falsification)

Prevents false-positive inflation when analytical depth increases without proportional biological signal (e.g. Discovery #72: 5B -> 4B).