Ecology & Evolution | Dec 2, 2025 | 7 min read | By Bryan White

Adaptation in the Chernobyl Exclusion Zone: Melanin as an Energy Transducer

Evolution * DNA * Urban Environment

Introduction: The Radiative Frontier of Reactor No. 4

The April 26, 1986 explosion of Unit 4 at the Chernobyl Nuclear Power Plant released large quantities of Cesium-137, Strontium-90, and Plutonium-239, creating an ionizing radiation environment sufficient to hydrolyze intracellular water, generate dense clouds of reactive oxygen species (ROS), and shatter genomic DNA.[1][2] While radiobiologists initially expected the interior of the Reactor No. 4 sarcophagus -- where local dose rates exceeded 10,000 Roentgens per hour -- to remain sterile for decades, surveys conducted inside the damaged containment structures during the 1990s revealed thriving colonies of black, melanized filamentous fungi coating the concrete reactor walls, cooling pools, and fuel rod debris.[2][3]

Isolated and characterized by Nelli Zhdanova, Tatyana Tugai, and colleagues at the Institute of Microbiology and Virology in Kyiv, the reactor's fungal biodiversity -- spanning over 200 species across the exclusion zone and 37 species inside the Unit 4 Shelter -- was heavily skewed in the highest-radiation zones toward dematiaceous (dark-pigmented) taxa.[2][12] Key species included Cladosporium sphaerospermum, Wangiella dermatitidis (Exophiala dermatitidis), and Cryptococcus neoformans.[2][5] Rather than merely tolerating radiation, these organisms exhibit positive radiotropism -- directional hyphal growth toward beta (P-32) and gamma (Cd-109) radiation beams -- and bio-etch radioactive graphite "hot particles."[5][12] Subsequent biophysical investigations indicate a process termed radiosynthesis, in which the polymeric pigment melanin acts as an energy transducer that captures high-energy electromagnetic radiation and couples it to cellular electron transport.[6]

The Physics and Biochemistry of Fungal Radiosynthesis

To determine whether melanin actively transduces ionizing radiation in a manner analogous to chlorophyll in visible-light photosynthesis, Ekaterina Dadachova, Arturo Casadevall, and colleagues at the Albert Einstein College of Medicine examined the two primary fungal melanin pathways: endogenous 1,8-dihydroxynaphthalene (DHN-melanin), synthesized via polyketide synthase in C. sphaerospermum and W. dermatitidis, and exogenous L-DOPA-derived DOPA-melanin, polymerized by laccase (Lac1/Lac2) in C. neoformans.[4][6] Both polymers consist of stacked planar sheets of conjugated aromatic rings that behave as organic semiconductors with delocalized -electrons and stable free-radical populations.[6]

In their landmark 2007 PLoS ONE study, Dadachova and Casadevall exposed melanized strains and albino mutants to Cesium-137 gamma radiation at dose rates ~500 times terrestrial background.[6] Electron Spin Resonance (ESR) spectroscopy demonstrated that Compton scattering and exciton generation directly altered the unpaired electron spin state of melanin.[6] Under nutrient-limited conditions, irradiated

melanized W. dermatitidis, C. neoformans, and C. sphaerospermum accumulated significantly higher dry-weight biomass and colony-forming units (CFUs) than non-irradiated controls or albino mutants.[5][6] Furthermore, in cell-free biochemical assays, irradiated melanin accelerated the chemical reduction of NAD+ to NADH at a rate fourfold higher than non-irradiated melanin, demonstrating direct electron transfer capable of feeding the mitochondrial respiratory chain.[6]

| Species | | Melanin Pathway & Precursor | | Key Enzyme | | Growth Response (500x Background) | | Biophysical Mechanism |

| Cladosporium sphaerospermum | | DHN (Endogenous Acetyl-CoA) | | Polyketide Synthase (PKS) | | Positive (Increased biomass & radiotropism) | | Constitutive cell-wall DHN antenna & radical scavenging[2][4] |

| Wangiella dermatitidis | | DHN (Endogenous Acetyl-CoA) | | Polyketide Synthase (PKS) | | Positive (Elevated CFU & dry weight) | | Constitutive DHN electron transduction[5] |

| Cryptococcus neoformans (Melanized) | | DOPA (Exogenous L-DOPA) | | Laccase (Lac1, Lac2) | | Positive (Accelerated metabolic activity) | | Compton electron transfer; 4x NADH reduction rate[6] |

| Cryptococcus neoformans (Albino) | | None (Laccase mutant) | | Absent | | Neutral / Negative (Radiolytic inhibition) | | Lack of semiconductive pigment shield or transducer[5][6] |

Evolutionary Atavism and Thermodynamic Critiques

Because complex bioenergetic pathways do not arise de novo within a few decades, Casadevall proposed that fungal radiotrophy represents an evolutionary atavism dating to the Early Cretaceous (~145-100 Ma) and major mass extinction boundaries.[5] During geomagnetic polarity reversals ("magnetic zero" intervals of the M-sequence), the collapse of Earth's magnetosphere exposed the surface biosphere to elevated Galactic Cosmic Rays (GCRs) and Solar Particle Events.[5] Paleontological "fungal spikes" -- sedimentary horizons densely packed with melanized fungal spores coinciding with floral and faunal die-offs -- suggest that melanin conferred both radioprotection and supplemental energy harvesting during high-radiation crises.[6]

Nevertheless, the strict definition of radiosynthesis remains debated. In a 2015 analysis, Eric Walberg noted that the raw ionizing energy deposited by Chernobyl dose rates is orders of magnitude smaller than the chemical bond energy supplied by trace organic substrates, proposing that accelerated growth could reflect radiation-induced hormesis (upregulated DNA repair and metabolic overdrive) or radiolytic cleavage of agar polysaccharides.[10] Moreover, unlike photoautotrophs, radiotrophic fungi have not been shown to fix atmospheric carbon dioxide independently, functioning instead as radio-stimulated heterotrophs.[5] Even so, the absence of growth stimulation in irradiated albino mutants and the cell-free fourfold NADH reduction confirm a melanin-specific electron transduction mechanism.[5][6]

Orbital Validation on the ISS and Martian Myco-Shielding

To evaluate radiotrophic fungi as self-regenerating radiation shields for deep-space exploration, Graham Shunk, Xavier Gomez, and Nils Averesch cultivated Cladosporium sphaerospermum aboard the International Space Station (ISS) for 30 days using a split Petri dish equipped with Geiger-M ller counters beneath both the inoculated agar and a sterile control.[7] Exposed to microgravity and orbital cosmic radiation (~40-80x terrestrial background), C. sphaerospermum exhibited a specific growth rate 1.21 +/- 0.37 times higher than ground controls.[7]

Once the fungal lawn reached a thickness of 1.7 mm, ionizing radiation flux measured beneath the biomass dropped by 1.82% to 2.17% relative to the control.[7] Extrapolating via linear attenuation modeling, the researchers calculated that a ~21 cm layer of pure living C. sphaerospermum -- or a ~9 cm equimolar composite of melanin-rich fungal biomass and Martian regolith -- could attenuate surface radiation on Mars down to terrestrial baseline levels, avoiding the secondary neutron spallation characteristic of aluminum hulls.[7][9]

| Shielding Material (Mars Scenario) | | Required Thickness | | Attenuation Mechanism | | Operational Advantages & Limitations |

| Living C. sphaerospermum Biomass | | ~21 cm | | Free-radical scavenging + hydrogen mass attenuation | | Self-replicating, self-healing, In-Situ Resource Utilization (ISRU); requires water/nutrients[7] |

| Melanin / Martian Regolith Composite | | ~9 cm | | High-density mineral scattering + melanin electronic capture | | Compact structural footprint; high attenuation efficiency[7] |

| Pure Martian Regolith | | ~12-15 cm | | Passive mass attenuation | | Abundant in situ; heavy, brittle, no biological self-repair[7] |

| Aerospace Aluminum Baseline | | Structural hull | | Metallic atomic scattering | | High tensile strength; high launch mass & secondary GCR neutron emission[9] |

Conclusion: From Nuclear Bioremediation to Astrobiology

The discovery of radiotrophic fungi inside Chernobyl's Reactor No. 4 has expanded the known boundaries of biological evolution and bioenergetics. Whether operating as primary thermodynamic transducers or melanin-boosted heterotrophs, organisms such as Cladosporium sphaerospermum and Cryptococcus neoformans demonstrate that semiconductive biological pigments can convert ionizing radiation into metabolic reducing power.[6] On Earth, their capacity to bioaccumulate radionuclides offers practical pathways for mycoremediation of contaminated nuclear sites; beyond Earth, their validation aboard the ISS points toward self-healing myco-architecture for lunar and Martian habitats and widens the astrobiological search for melanized life in high-radiation space environments.[7][8][11]

Updates... (Nov 23, 2026): Replaced secondary Wikipedia citations (Refs #1 and #4) with primary UNSCEAR radiological assessment documentation and peer-reviewed literature on fungal DHN/DOPA melanin synthesis (Eisenman & Casadevall, 2011), removed numbered section headings, and standardized the References section heading.

References

* United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR) (2021). The Chornobyl Accident. UNSCEAR. -- https://www.unscear.org/unscear/en/areas-of-work/chernobyl.html

* Zhdanova, N. N., Zakharchenko, V. A., Vember, V. V., & Nakonechnaya, L. T. (Dec 1, 2000). Fungi from Chernobyl: mycobiota of the inner regions of the containment structures of the damaged nuclear reactor. Mycological Research, 104(12), 1421-1426. -- https://doi.org/10.1017/S0953756200002756

* Sim es, M. F., Cortes o, M., Azua-Bustos, A., et al. (2023). The relevance of fungi in astrobiology research - Astromycology. Mycosphere, 14(1). -- https://doi.org/10.5943/mycosphere/14/1/13

* Eisenman, H. C., & Casadevall, A. (Dec 16, 2011). Synthesis and assembly of fungal melanin. Applied Microbiology and Biotechnology, 93, 931-940. -- https://doi.org/10.1007/s00253-011-3777-2

* Dadachova, E., & Casadevall, A. (Oct 24, 2008). Ionizing radiation: how fungi cope, adapt, and exploit with the help of melanin. Current Opinion in Microbiology. -- https://doi.org/10.1016/j.mib.2008.09.013

* Dadachova, E., Bryan, R. A., Huang, X., et al. (May 23, 2007). Ionizing Radiation Changes the Electronic Properties of Melanin and Enhances the Growth of Melanized Fungi. PLoS ONE. -- https://doi.org/10.1371/journal.pone.0000457

* Shunk, G. K., Gomez, X. R., & Averesch, N. J. H. (Jul 5, 2022). Cultivation of the Dematiaceous Fungus Cladosporium sphaerospermum Aboard the International Space Station and Effects of Ionizing Radiation. Frontiers in Microbiology. -- https://doi.org/10.3389/fmicb.2022.877625

* Cassaro, A., Pacelli, C., Aureli, L., et al. (Apr 22, 2022). Investigation of fungal biomolecules after Low Earth Orbit exposure: a testbed for the next Moon missions. Environmental Microbiology. -- https://doi.org/10.1111/1462-2920.15995

* Singleterry, R. C., et al. (2009). Evaluation of Multi-Functional Materials for Deep Space Radiation Shielding. NASA Technical Reports Server. -- https://ntrs.nasa.gov/api/citations/20090020691/downloads/20090020691.pdf

* Walberg, E. W. (May 2015). A critical analysis of claims of radiosynthesis by fungi. M.S. Thesis, University of Wisconsin-La Crosse. -- https://minds.wisc.edu/server/api/core/bitstreams/c2391280-0d7d-4081-9ba5-3d8ad1e7cdc1/content

* Sakauchi, K., & Otaki, J. M. (Feb 10, 2024). Soil Microbes and Plant-Associated Microbes in Response to Radioactive Pollution May Indirectly Affect Plants and Insect Herbivores: Evidence for Indirect Field Effects from Chernobyl and Fukushima. Microorganisms. -- https://doi.org/10.3390/microorganisms12020364

* Dighton, J., Tugay, T., & Zhdanova, N. (Feb 15, 2008). Fungi and ionizing radiation from radionuclides. FEMS Microbiology Letters. -- https://doi.org/10.1111/j.1574-6968.2008.01076.x