Ecology & Evolution | Jan 11, 2026 | 9 min read | By Bryan White

An Integrative Perspective on Bat Evolution: From Eocene Origins to Genomic Frontiers

Biology * Evolution * DNA

The Chiropteran Enigma and Laurasiatherian Origins

With over 1,460 recognized species constituting roughly 20% of extant mammalian diversity, the order Chiroptera represents the only mammalian lineage to achieve true powered flight.[1][3] While early anatomical classifications grouped bats within the "Archonta" alongside primates and colugos, molecular phylogenetics firmly placed Chiroptera within the superorder Laurasiatheria alongside carnivores, pangolins, and ungulates.[1][4] Synthesizing paleontological, biomechanical, and comparative genomic advances reviewed through 2025,[4] modern research demonstrates that bat evolution is an integrated physiological cascade: the metabolic demands of powered flight reshaped cranial morphology, ultrasonic sensory systems, intestinal sugar transport, DNA repair, and innate viral tolerance -- placing bats at the center of global epidemiological and evolutionary genetics research.[2][5][6]

Eocene Fossil Chronicles and Developmental Origins of Flight

Bats appear across multiple continents during the Early Eocene greenhouse (~52-56 million years ago), with exceptional skeletal preservation in Wyoming's Green River Formation Lagerst tte resolving the longstanding "flight-first" versus "echolocation-first" debate:[3][9][11]

* Onychonycteris finneyi (~52.5 Ma): Retained claws on all five digits of its handwing and intermediate forelimb-to-hindlimb proportions suited for agile arboreal climbing.[9][10][11] Aerodynamic reconstructions indicate broad, short wings operating as a flapping-gliding hybrid, while its small cochlea -- comparable to non-echolocating pteropodids -- confirms that powered flight evolved prior to laryngeal echolocation.[9][12][13]

* Icaronycteris gunnelli & I. index (~52 Ma): Described in 2023 as the oldest known bat skeletons (~25 g body mass), I. gunnelli and its sister taxon I. index possess enlarged cochleae and reduced outer-digit claws indicative of active laryngeal sonar and rapid Early Eocene diversification across multiple families.[3][7][8]

Embryonically, the transition from terrestrial forelimb to wing (patagium) is governed by shifts in Bmp (Bone Morphogenetic Protein) and Fgf (Fibroblast Growth Factor) signaling that promote longitudinal cartilage proliferation in digits II-V while inhibiting interdigital apoptosis (programmed cell death) to preserve the aerodynamic wing membrane.[4][14]

| Eocene Taxon | | Age (Ma) | | Locomotor Mode | | Laryngeal Sonar | | Diagnostic Skeletal & Ecological Traits |

| Onychonycteris finneyi | | ~52.5 | | Flapping-Gliding Hybrid | | Absent | | Claws on all 5 digits; small cochlea; climbing limb proportions |

| Icaronycteris gunnelli | | ~52.0 | | Powered Flight | | Present | | ~25 g body mass; claws on digits I-II only; expanded cochlea |

| Icaronycteris index | | ~52.0 | | Powered Flight | | Present | | Expanded cochlea; preserved stomach fish scales indicating gleaning |

| Palaeochiropteryx | | ~48.0 | | Maneuverable Flight | | Present (High Duty) | | Low-aspect wing geometry; fossilized lepidopteran/insect stomach contents |

The Sensory Revolution: Echolocation, Prestin Convergence, and Cranial Trade-offs

While Old World fruit bats (Pteropodidae) rely primarily on vision and olfaction (with Rousettus utilizing lingual clicks), most Yangochiroptera and Yinochiroptera employ high-frequency laryngeal echolocation.[1] Comparative genomics reveals remarkable convergent evolution in the cochlear motor protein gene Prestin: echolocating bats and odontocete whales share identical amino acid substitutions that tune outer hair cell electromotility to ultrasonic frequencies.[15] Simultaneously, the vocalization transcription factor FOXP2 exhibits accelerated mutation rates in laryngeal echolocators, supporting sensorimotor calls emitted up to 200 times per second during terminal prey buzzes.[16]

Recent 2024-2025 acoustic modeling and onboard microphone telemetry in dense swarms of Brazilian free-tailed bats (Tadarida brasiliensis) have overturned the classic "spectral jamming avoidance response" hypothesis, demonstrating that emerging swarms navigate via collective amplitude gradients (acoustic shadows) and rapid spatial dispersion rather than frequency shifting.[17][18] Furthermore, sound emission pathways dictate cranial architecture: oral emitters (e.g., Vespertilionidae) evolve dorsally upturned rostra to align open-mouth calls with flight, whereas nasal emitters (e.g., Rhinolophidae, Phyllostomidae) feature downturned rostra and acoustic nose-leaves -- decoupling the jaws from sonar emission and unlocking extensive masticatory diversification.[19][20]

Cranial Biomechanics and the Genomic Architecture of Diet

Freed from oral sonar constraints, Neotropical leaf-nosed bats (Phyllostomidae) radiated across insectivorous, frugivorous, nectarivorous, carnivorous, and sanguivorous niches.[21] Hard-fruit specialists (Artibeus) evolved short rostra and broad zygomatic arches that maximize masseter lever mechanical advantage,[20][22] whereas nectarivores (Anoura, Glossophaga) evolved elongated, delicate rostra housing brush-tipped extensible tongues.[23]

At the level of the genome and intestinal biochemistry, dietary specialization is driven by transport upregulation and adaptive gene loss:

* Hyperglycemic Sugar Assimilation: Nectar bats (Anoura geoffroyi) survive postprandial blood glucose exceeding 700 mg/dL via high constitutive apical expression of intestinal Slc2a2 (GLUT2), whereas frugivores (Artibeus jamaicensis) upregulate Slc5a1 (SGLT1) and Slc2a5 (GLUT5) for fructose/glucose co-absorption.[23]

* Sanguivorous Gene Pseudogenization: Common vampire bats (Desmodus rotundus) have lost at least 13 functional genes, including insulin-secretion regulators (adapting to carbohydrate-poor blood meals) and the iron-recycling gene REP15 -- allowing

enterocytes to slough and excrete toxic iron loads 800-fold higher than human diets -- alongside widespread pseudogenization of sweet (TAS1R2) and umami (TAS1R1) taste receptors.[24][25][26]

Flight Metabolism, Viral Tolerance, and Ecosystem Services

Powered flight elevates metabolic rates 3- to 5-fold above terrestrial locomotion, generating high fluxes of reactive oxygen species (ROS) and cytosolic oxygen radicals.[2] To prevent oxidative genomic damage, bats evolved positive selection in DNA double-strand break repair pathways, upregulated autophagy, and replication-independent telomere maintenance -- allowing 7-gram Myotis brandtii bats to live over 40 years.[6][27]

Crucially, because flight-induced mitochondrial DNA leakage into the cytoplasm would otherwise trigger chronic sterile inflammation, bats evolved a dampened NLRP3 inflammasome and a mutated S358 residue in the STING DNA-sensing pathway, paired with constitutive baseline expression of antiviral Interferon-alpha (IFN-).[5][28][29][30] This immunotolerance prevents lethal cytokine storms when hosting filoviruses or coronaviruses.[2][6] Ecologically, bats provide an estimated USD 3.7-53 billion annually in U.S. agricultural pest suppression and pollinate over 500 plant species, though North American hibernating populations face high mortality from White-Nose Syndrome (Pseudogymnoascus destructans) and climate change.[31][32]

| Trait / Pathway | | Insectivorous Laryngeal Bats | | Pteropodidae (Fruit Bats) | | Sanguivorous Bats (Desmodus) |

| Primary Orientation | | High-duty laryngeal echolocation (Prestin / FOXP2) | | Large-eye vision & olfaction (lingual clicks in Rousettus) | | Nasal-emission echolocation, olfaction & infrared pit sensors |

| Taste Receptors | | Functional Umami (TAS1R1); frequent Sweet loss | | Conserved Sweet (TAS1R2); frequent Umami loss | | Pseudogenized Sweet (TAS1R2) & Umami (TAS1R1) |

| Metabolic / Gut Genetics | | Standard mammalian proteolytic profile | | Apical GLUT2 (nectar) & SGLT1/GLUT5 (fruit) | | Loss of insulin regulators & REP15 iron-recycling gene |

| Innate Immunity | | Dampened NLRP3 & STING S358 mutation | | Constitutive IFN- & dampened NLRP3 inflammasome | | Dampened inflammasome & antiviral tolerance |

Conclusion: Evolutionary Integration Across Scales

From the five-clawed wings of Eocene Onychonycteris to the apical GLUT2 transporters of Neotropical nectar bats and the S358 STING mutation enabling viral tolerance, Chiroptera exemplifies how a single biomechanical innovation -- powered flight -- can reorganize an entire mammalian genome.[4][9][23] Protecting bat diversity against fungal pathogens and habitat fragmentation is essential both for global agricultural stability and for translating chiropteran longevity and immunotolerance into human medicine.[2][31][32]

Updates... (Nov 22, 2026): Added verified publication and last-updated dates to National Park Service Ref [11] (Sep 26, 2025) and USGS Ref [32] (Sep 26, 2019);

earlier (Nov 19, 2026) upgraded citations to verified primary journal DOIs and institutional releases.

Works Cited

* [1] Teeling, E. C., Dool, S., & Springer, M. S. (2012). Phylogenies, fossils and functional genes: the evolution of echolocation in bats. In Evolutionary History of Bats, Cambridge University Press. https://doi.org/10.1017/cbo9781139045599.002

* [2] Ng, W. L., & Wang, L.-F. (2025). Translational lessons from the balanced immune system in bats. Disease Models & Mechanisms. https://doi.org/10.1242/dmm.050763

* [3] Rietbergen, T. B., van den Hoek Ostende, L. W., Aase, A., et al. (2023). The oldest known bat skeletons and their implications for Eocene chiropteran diversification. PLOS ONE. https://doi.org/10.1371/journal.pone.0283505

* [4] Santana, S. E., Bernal-Rivera, A., Chhaya, V., et al. (2025). An Integrative Perspective on Bat Evolution. Annual Review of Ecology, Evolution, and Systematics. https://doi.org/10.1146/annurev-ecolsys-102723-045407

* [5] Ahn, M., Anderson, D. E., Zhang, Q., et al. (2019). Dampened NLRP3-mediated inflammation in bats and implications for a special viral reservoir host. Nature Microbiology. https://doi.org/10.1038/s41564-019-0371-3

* [6] Pei, G., Balkema-Buschmann, A., & Dorhoi, A. (2024). Disease tolerance as immune defense strategy in bats: One size fits all? PLOS Pathogens. https://doi.org/10.1371/journal.ppat.1012471

* [7] Natural History Museum (London) (2023). Oldest bat skeletons ever found described as new species. https://www.nhm.ac.uk/discover/news/2023/april/oldest-bat-skeletons-ever-found-described-new-species.html

* [8] National Geographic (2023). 52-million-year-old bat skeletons are the oldest ever -- and tell a 'really weird' tale. https://www.nationalgeographic.com/science/article/52-million-year-old-bat-skeletons-are-the-oldest-ever

* [9] Simmons, N. B., Seymour, K. L., Habersetzer, J., & Gunnell, G. F. (2008). Primitive Early Eocene bat from Wyoming and the evolution of flight and echolocation. Nature. https://doi.org/10.1038/nature06549

* [10] Royal Ontario Museum (2008). Researchers Discover 'Missing Link' in Bat Evolution (Onychonycteris finneyi). https://www.rom.on.ca/news-releases/researchers-discover-missing-link-bat-evolution

* [11] U.S. National Park Service (Sep 26, 2025). Fossil Mammals -- Fossil Butte National Monument. https://www.nps.gov/fobu/learn/nature/fossil-mammals.htm

* [12] Amador, L. I., Simmons, N. B., & Giannini, N. P. (2019). Aerodynamic reconstruction of the primitive fossil bat Onychonycteris finneyi (Mammalia: Chiroptera). Biology Letters. https://doi.org/10.1098/rsbl.2018.0857

* [13] Burtner, A. E., Grossnickle, D. M., & Santana, S. E. (2024). Gliding toward an understanding of the origin of flight in bats. PeerJ. https://doi.org/10.7717/peerj.17824

* [14] Sears, K. E., Behringer, R. R., Rasweiler, J. J., & Niswander, L. A. (2006). Development of bat flight: Morphologic and molecular evolution of bat wing digits. Proceedings of the National Academy of Sciences. https://doi.org/10.1073/pnas.0509716103

* [15] Li, Y., Liu, Z., Shi, P., & Zhang, J. (2010). The hearing gene Prestin unites echolocating bats and whales. Current Biology. https://doi.org/10.1016/j.cub.2009.11.042

* [16] Li, G., Wang, J., Rossiter, S. J., Jones, G., & Zhang, S. (2007). Accelerated FoxP2 evolution in echolocating bats. PLOS ONE. https://doi.org/10.1371/journal.pone.0000900

* [17] Vanderelst, D., & Peremans, H. (2025). How swarming bats can use the collective soundscape for obstacle avoidance. PLOS Computational Biology. https://doi.org/10.1371/journal.pcbi.1013013

* [18] Goldshtein, A., Mazar, O., Harten, L., et al. (2025). Onboard recordings reveal how bats maneuver under severe acoustic interference. Proceedings of the National Academy of Sciences. https://doi.org/10.1073/pnas.2407810122

* [19] Gilley, D. T., Santana, S. E., & Arbour, J. H. (2024). Macroevolutionary and biomechanical implications of rostral flexion in bat skulls: a major early driver of cranial evolution in bats. Biological Journal of the Linnean Society. https://doi.org/10.1093/biolinnean/blae123

* [20] Arbour, J. H., Curtis, A. A., & Santana, S. E. (2019). Signatures of echolocation and dietary ecology in the adaptive evolution of skull shape in bats. Nature Communications. https://doi.org/10.1038/s41467-019-09951-y

* [21] Santana, S. E., Sadier, A., & Mello, M. A. R. (2024). The ecomorphological radiation of phyllostomid bats. Evolutionary Journal of the Linnean Society. https://doi.org/10.1093/evolinnean/kzae032

* [22] Santana, S. E., Grosse, I. R., & Dumont, E. R. (2012). Dietary hardness, loading behavior, and the evolution of skull form in bats. Evolution. https://doi.org/10.1111/j.1558-5646.2012.01615.x

* [23] Camacho, J., Bernal-Rivera, A., Pe a, V., et al. (2024). Sugar assimilation underlying dietary evolution of Neotropical bats. Nature Ecology & Evolution. https://doi.org/10.1038/s41559-024-02485-7

* [24] Blumer, M., Brown, T., Freitas, M. B., et al. (2022). Gene losses in the common vampire bat illuminate molecular adaptations to blood feeding. Science Advances. https://doi.org/10.1126/sciadv.abm6494

* [25] Max Planck Institute of Molecular Cell Biology and Genetics (MPI-CBG) (2022). Vampires with genetic defects. https://www.mpi-cbg.de/news-outreach/news-media/article/vampires-with-genetic-defects

* [26] Zhao, H., Zhou, Y., Pinto, C. M., et al. (2010). Evolution of the Sweet Taste Receptor Gene Tas1r2 in Bats. Molecular Biology and Evolution. https://doi.org/10.1093/molbev/msq152

* [27] Wilkinson, G. S., & Adams, D. M. (2019). Recurrent evolution of extreme

longevity in bats. Biology Letters. https://doi.org/10.1098/rsbl.2018.0860

* [28] Das, S., Jain, A., Chaudhary, S., et al. (2025). Bat adaptations in inflammation and cell death regulation contribute to viral tolerance. mBio. https://doi.org/10.1128/mbio.03204-23

* [29] Fu, Z., Shao, S., Zhang, J., et al. (2023). Bat STING drives IFN-beta production in anti-RNA virus innate immune response. Frontiers in Microbiology. https://doi.org/10.3389/fmicb.2023.1232314

* [30] Bondet, V., Le Baut, M., Le Poder, S., et al. (2021). Constitutive IFN Protein Production in Bats. Frontiers in Immunology. https://doi.org/10.3389/fimmu.2021.735866

* [31] Boyles, J. G., Cryan, P. M., McCracken, G. F., & Kunz, T. H. (2011). Economic Importance of Bats in Agriculture. Science. https://doi.org/10.1126/science.1201366

* [32] U.S. Geological Survey (USGS) (Sep 26, 2019). Why are bats important? https://www.usgs.gov/faqs/why-are-bats-important