Ecology & Evolution | Dec 3, 2025 | 7 min read | By Bryan White

Apex Predators of the Aptian: How Cardabiodontid Sharks Challenged Marine Reptiles

Biology * Evolution * Paleontology

Introduction: The Fifteen-Million-Year Shift in Shark Gigantism

In traditional reconstructions of Mesozoic vertebrate evolution, Early Cretaceous seas are depicted as the undisputed domain of giant marine reptiles -- pliosaurs, plesiosaurs, and ichthyosaurs -- with chondrichthyan sharks relegated to secondary, mid-trophic tiers until the Late Cretaceous radiations of the Cenomanian and Turonian stages (~100-90 Ma).[1] However, a landmark paleontological discovery from the Darwin Formation in Australia's Northern Territory has fundamentally overturned this gradualist timeline.[2] Published in Communications Biology by Mohamad Bazzi (Uppsala University and Stanford University), Mikael Siversson (Western Australian Museum), and an international team of paleontologists and ichthyologists, the study describes five associated vertebral centra (specimen NTM P22-33) belonging to a colossal cardabiodontid mackerel shark (Order Lamniformes).[2][3]

Dating to the late Aptian stage (~115 Ma), these fossilized vertebrae demonstrate that macrophagous lamniforms achieved total body lengths of 6 to 8 meters and body masses exceeding 3 metric tons at least 10 to 15 million years earlier than previously documented.[1][3] Rather than waiting for the Cenomanian extinction of ichthyosaurs or Late Cretaceous thermal maxima to attain mega-predator dimensions (defined as total length >6 meters), lamniform sharks established the genetic, anatomical, and physiological pathways for gigantism shortly after their Jurassic divergence, competing directly alongside marine reptiles at the apex of the Early Cretaceous food web.[1]

Geologic Setting: The Darwin Formation and the Eromanga Sea

To evaluate the ecological context of the Darwin cardabiodontid, researchers reconstructed the paleogeography of the Australian Cretaceous. During the Aptian (125-113 Ma), Australia formed the high-latitude eastern sector of the Gondwana supercontinent, where a major marine transgression flooded the continental interior to create the vast epicontinental Eromanga Sea.[5] Located near modern-day Darwin in the Northern Territory, the Darwin Formation represents the northern gateway of this inland waterway where it opened into the ancient Tethys Ocean, functioning as a vital biogeographic corridor that connected tropical Tethyan waters with the cooler, high-latitude basins of the Australian interior.[1]

Because chondrichthyan skeletons consist of prismatic cartilage that rapidly disaggregates after death, preserving three-dimensional shark vertebrae requires exceptional taphonomic conditions. The Darwin Formation comprises radiolarian claystones and glauconitic siltstones deposited across a low-energy, shallow marine shelf with a muddy substrate, characterized by slow sedimentation and dysaerobic (low-oxygen) conditions at the sediment-water interface.[7][8] This benthic geochemistry inhibited scavenging and bacterial decomposition long enough for

partial early diagenetic mineralization to preserve the three-dimensional structure of the five associated centra across 115 million years.[3]

Furthermore, despite the global greenhouse climate of the Cretaceous, Australian epeiric seas were cool. Whereas southern stratigraphic units such as the contemporaneous Bulldog Shale preserve glendonite pseudomorphs (derived from ikaite crystals forming in near-freezing water) and ice-rafted dropstones, the northern Darwin shelf supported a cool-temperate coastal ecosystem.[5] The survival and dominance of gigantic sharks across these high-latitude Gondwanan shelves suggest that early cardabiodontids had already evolved regional endothermy (mesothermy), enabling them to maintain elevated cruising metabolic rates in cool waters.[1]

Systematic Paleontology and Allometric Size Reconstruction

While fossil shark taxonomy is most commonly anchored by isolated teeth, vertebral centrum architecture is diagnostic at the family level within the Lamniformes. Specimen NTM P22-33 consists of five associated asterospondylous vertebral centra displaying the tightly spaced, radiating calcified lamellae characteristic of large mackerel sharks.[9] Based on the robust, heavy-set centrum proportions and internal lamellar geometry, Siversson and colleagues assigned the specimen to the extinct family Cardabiodontidae -- a clade originally erected by Siverson (Siversson) in 1999 from the Cenomanian (~95 Ma) type species Cardabiodon ricki of the Giralia Range in Western Australia.[2][4][10]

To rigorously reconstruct the body dimensions of the Darwin cardabiodontid and account for non-linear ontogenetic scaling, Bazzi et al. compiled morphometric measurements from nearly 2,000 extant shark specimens across multiple lamniform families and applied log-transformed bivariate allometric regression relating maximum vertebral centrum diameter to total body length (TL) and body mass.[1][3] Whereas a 6-meter adult modern Great White Shark (Carcharodon carcharias) possesses vertebral centra approximately 80 mm (8 cm) in diameter, the recovered Darwin centra exceed 120 mm (12.6 cm) across.[3] The regression models yield an estimated total length of 6 to 8 meters (20-26 feet) and a body mass exceeding 3,000 kilograms (3.3 tons), placing the Aptian predator in the same length class as the largest extant Great Whites and the Late Cretaceous Cretoxyrhina mantelli, but with a substantially heavier, stockier build.[3][10]

| Anatomical & Ecological Metric | | Darwin Cardabiodontid (NTM P22-33) | | Cardabiodon ricki (Type Species) | | Great White Shark (Carcharodon carcharias) |

| Geological Age | | ~115 Ma (Late Aptian, Early Cretaceous) | | ~95 Ma (Cenomanian, Late Cretaceous) | | ~16 Ma (Miocene) - Present |

| Max Vertebral Centrum Width | | >120 mm (12.6 cm) | | ~90-100 mm | | ~80 mm (at 6 m TL) |

| Estimated Total Length | | 6 - 8 meters (20 - 26 ft) | | 5 - 5.5 meters | | Up to ~6 meters |

| Estimated Body Mass | | >3,000 kg (>3.3 metric tons) | | ~1,500 - 2,000 kg | | ~1,900 - 2,200 kg (at 6 m TL) |

| Body Plan & Ecology | | Stocky, heavy-bodied coastal/shelf apex predator | |

Robust antitropical macrophagous predator | | Fusiform, regionally endothermic apex predator |

Paleoecology: Challenging the Marine Reptile Empire

The presence of a 3-ton lamniform predator in the late Aptian compels a major revision of Early Cretaceous marine food webs. Sustaining a macrophagous predator -- one capable of dismembering large vertebrate prey -- requires a calorie-dense biomass base. The northern gateway of the Eromanga Sea supported abundant marine reptiles, including the 7-meter ophthalmosaurid ichthyosaur Platypterygius australis and the 2.5-meter leptocleidid plesiosaur Umoonasaurus demoscyllus, alongside large Tethyan teleost fishes.[5][7][11]

Although giant thalassophonean pliosaurs such as Kronosaurus queenslandicus (10-11 meters) occupied the absolute size ceiling of the Australian inland sea, the Darwin cardabiodontid co-ruled the apex guild via ecological niche partitioning.[6] Its heavy-set body plan suited high-torque ambush predation and scavenging of large marine reptile carcasses, while regional endothermy allowed cardabiodontids to forage in deeper, colder water columns that were metabolically expensive for air-breathing reptiles.[1] Moreover, the rise of giant Aptian sharks coincided with the protracted decline of ichthyosaur diversity prior to their Cenomanian extinction, suggesting that cardabiodontid predation on juvenile marine reptiles helped reshape Cretaceous ocean demographics.[6]

Conclusion: A New Baseline for Chondrichthyan Evolution

The discovery and allometric analysis of the Darwin Formation cardabiodontid (NTM P22-33) push the onset of lamniform shark gigantism back by 15 million years into the Early Cretaceous.[1][3] Demonstrating that 6-to-8-meter, 3-ton mackerel sharks patrolled Gondwanan shelves alongside pliosaurs, plesiosaurs, and ichthyosaurs 115 million years ago, these fossils confirm that modern-style sharks rapidly ascended to apex trophic status well before the Late Cretaceous thermal maximum.[2][3]

Updates... (Nov 21, 2026): Upgraded the Platypterygius australis reference to the peer-reviewed osteological study by Zammit, Norris, & Kear (Dec 2, 2010), added publication metadata for the Northern Territory Geological Survey Money Shoal Basin chapter (Ahmad & Munson, 2013) and exact dates across all 11 sources, split grouped inline citations, and added the standard reference list identifier. (Prior Nov 16, 2026 update replaced Wikipedia entries with peer-reviewed CrossRef DOIs.)

References & Primary Sources

* [1] Nautilus Magazine (Nov 25, 2025). This Huge Ocean Beast Shifts Sharks' Evolutionary Timeline. -- https://nautil.us/this-huge-ocean-beast-shifts-sharks-evolutionary-timeline-1250168/

* [2] Western Australian Museum (Nov 24, 2025). Ancient Super Sharks: Australian Fossils Rewrite Shark Evolution. -- https://visit.museum.wa.gov.au/learn/news-stories/ancient-super-sharks-australian-fossils-rewrite-shark-evolution

* [3] Bazzi, M., Siversson, M., Wintner, S., et al. (Oct 25, 2025). Early gigantic lamniform marks the onset of mega-body size in modern shark evolution.

Communications Biology, 8(1), 1499. -- https://doi.org/10.1038/s42003-025-08930-y

* [4] Yahoo News Australia (Nov 25, 2025). Ancient discovery 'rewrites the timeline' of Australia by 10 million years. -- https://au.news.yahoo.com/ancient-discovery-rewrites-the-timeline-of-australia-by-10-million-years-055629483.html

* [5] Zammit, M., Norris, R. M., & Kear, B. P. (Dec 2, 2010). The Australian Cretaceous ichthyosaur Platypterygius australis: a description and review of postcranial remains. Journal of Vertebrate Paleontology, 30(6), 1726-1735. -- https://doi.org/10.1080/02724634.2010.521930

* [6] Kear, B. P. (2016). Cretaceous marine amniotes of Australia: perspectives on a decade of new research. Memoirs of Museum Victoria, 74, 17-28. -- https://doi.org/10.24199/j.mmv.2016.74.03

* [7] Open Access Government (Nov 24, 2025). Colossal shark fossils rewrite the history of ocean apex predators. -- https://www.openaccessgovernment.org/colossal-shark-fossils-rewrite-the-history-of-ocean-apex-predators/201585/

* [8] Ahmad, M., & Munson, T. J. (Comps.) (2013). Chapter 40: Money Shoal Basin. In Geology and Mineral Resources of the Northern Territory (Special Publication 5). Northern Territory Geological Survey. -- https://geoscience.nt.gov.au/gemis/ntgsjspui/bitstream/1/81520/1/GNT_Ch40_Money.pdf

* [9] Columbus State University (Nov 2025). Discovery: An old fish fossil tells new story about lamniform shark evolution. -- https://news.columbusstate.edu/posts/discovery-an-old-fish-fossil-tells-new-story-about-lamniform-shark-evolution/

* [10] Siverson, M. (1999). A new large lamniform shark from the uppermost Gearle Siltstone (Cenomanian, Late Cretaceous) of Western Australia. Transactions of the Royal Society of Edinburgh: Earth Sciences, 90(1), 49-66. -- https://doi.org/10.1017/S0263593300002509

* [11] Kear, B. P., Schroeder, N. I., & Lee, M. S. Y. (Jun 30, 2006). An archaic crested plesiosaur in opal from the Lower Cretaceous high-latitude deposits of Australia. Biology Letters, 2(4), 615-619. -- https://doi.org/10.1098/rsbl.2006.0504