Space & Astronomy | Aug 4, 2026 | 9 min read | By Bryan White

Beyond the Event Horizon: Tracking the Complete Feeding Cycle of a Black Hole

Space * Physics

Black Hole Accretion and the Outburst of Swift J1727.8 1613

Rather than acting as passive cosmic sinks that consume all infalling matter across their Schwarzschild radius, accreting black holes operate as inefficient regulatory engines[1]. Because gas captured from a companion star carries significant angular momentum, it spirals into a viscous, magnetized accretion disc heated to millions of Kelvin, radiating intensely in the X-ray band while redirecting a substantial fraction of the inflowing mass back into the interstellar medium via collimated relativistic jets and wide-angle equatorial disc winds[1][2][3].

The August 2023 eruption of the Galactic low-mass X-ray binary Swift J1727.8 1613 -- which rapidly reached seven times the X-ray flux of the Crab Nebula -- enabled a global multi-wavelength campaign tracking the evolution of an entire black hole feeding cycle[4][5][6]. Combining X-ray timing, radio interferometry, and 13 epochs of Very Large Telescope (VLT) X-Shooter optical spectroscopy, researchers discovered that dense, cool (

System Architecture and Hysteresis Across the Hardness-Intensity Diagram

Swift J1727.8 1613 comprises an 8-11 solar-mass black hole accreting via Roche-lobe overflow from an M3-M3.5 dwarf companion at a distance of 2.7-4.0 kiloparsecs[2][7]. Time-series optical photometry indicates an orbital period of 2.26 to 4.5 hours, while X-ray continuum fitting reveals a near-maximal dimensionless spin parameter (a* ~ 0.98) and a moderate disc inclination (~40 deg)[7][8].

| System Parameter | | Estimated Value | | Observational Method / Source |

| Primary Mass (Black Hole) | | 8 to 11 solar masses | | Hydrogen-alpha line width and FWHM calibration[7] |

| Donor Star Classification | | M3 to M3.5 Dwarf | | Quiescent titanium oxide (TiO) absorption bands[7] |

| Distance | | 2.7 to 4.0 kiloparsecs | | Diffuse interstellar bands and extinction modeling[7] |

| Orbital Period | | 2.26 to 4.5 hours | | Time-series optical photometry (double-humped modulation)[7] |

| Black Hole Spin Parameter | | 0.98 | | X-ray continuum spectral fitting[8] |

Throughout its 2023 outburst, Swift J1727.8 1613 traced a canonical q-shaped hysteresis loop across the X-ray hardness-intensity diagram[8]. In the initial Rising Hard State, the optically thick accretion disc remained truncated far from the innermost stable circular orbit (ISCO), while an inner advection-dominated hot

corona generated hard power-law X-rays via inverse-Compton scattering alongside a steady compact radio jet[8][9]. Coordinated NuSTAR, NICER, and Insight-HXMT observations detected an unusual non-thermal secondary hard component as low-energy count rates climbed toward 3,100 counts/s[8][10].

As the thin disc migrated inward and cooled the corona during the Hard-Intermediate and Soft-Intermediate transitions, Insight-HXMT recorded the simultaneous appearance of Type-B and Type-C quasi-periodic oscillations (QPOs) driven by Lense-Thirring relativistic precession, coinciding with a drop in broadband X-ray noise and the quenching of the compact jet via discrete ballistic radio ejections[11][12][13]. The system then entered the High-Soft State -- dominated by multi-temperature blackbody radiation extending to the ISCO and highly ionized iron X-ray absorption winds -- before mass depletion drove a reverse transition into the low-luminosity Dim-Hard State[5][7][9][14].

| Accretion State | | Inner Disc Geometry | | Dominant X-ray Emission | | Primary Outflow Signature |

| Rising Hard State | | Truncated far from the innermost stable circular orbit | | Hot Corona (Inverse-Compton scattering) | | Steady, continuous compact radio jet |

| Intermediate States | | Migrating inward, highly unstable | | Mixed Corona and Thermal Disc | | Discrete, relativistic ballistic jet ejections |

| High-Soft State | | Extends to the innermost stable circular orbit | | Thermal Accretion Disc (Multi-temperature Blackbody) | | Highly ionized X-ray disc winds; radio jet suppressed |

| Dim-Hard State | | Receding outward as accretion rate drops | | Rebuilding Corona | | Fading jet; emergence of broad, cool optical disc winds |

VLT X-Shooter Spectroscopy and the Late-Stage Cool Disc Wind

While X-ray telescopes probe the inner gravitational well, the VLT X-Shooter campaign tracked the outer disc across 13 optical and near-infrared epochs[1][6]. During the initial discrete jet flares (Epoch 1), hydrogen emission lines displayed strong red-velocity skews, relaxing during the soft state (Epochs 2-9) into narrower, symmetric double-peaked Keplerian profiles with elevated singly ionized helium (He II) flux -- demonstrating that intense flare irradiation illuminated outer disc radii while leaving the inner disc base stable[6][7].

The campaign's most unexpected finding emerged during the late Dim-Hard State (Epochs 10-11), when X-ray luminosity had fallen to ~1% of peak[1]. Under conventional models of black hole state transitions, thermally driven outflows are expected to subside rapidly once central radiation pressure and coronal illumination fade. Instead, the outer disc of Swift J1727.8 1613 produced deep, broad blueshifted higher-order Balmer hydrogen absorption lines alongside heavy suppression of the blue peak of He II emission[6][7]. Because absorption features only arise when cooler, lower-density gas intercepts the line of sight against a hotter continuum source, these blueshifted troughs provide unambiguous kinematic proof of a dense, expanding wind moving toward the observer[6][7].

Doppler kinematics and radiative transfer modeling confirm a dense, cool (

Physical Launch Mechanisms of Accretion Disc Winds

Launching dense matter out of a black hole's gravitational potential requires overcoming local escape velocity via three primary physical mechanisms[14]:

* Radiation Pressure: Direct photon momentum transfer via electron continuum scattering or atomic line driving dominates near or above the Eddington limit, but cannot independently power the heavy winds observed during Swift J1727.8 1613's sub-Eddington dim-hard state[14][17].

* Thermal (Compton) Driving: Central X-ray irradiation heats the flared outer disc atmosphere to millions of degrees; beyond the Compton radius, thermal gas velocities exceed local escape velocity, lifting outer-disc material into space[14][18][19].

* Magneto-Centrifugal (Blandford-Payne) Driving: Poloidal magnetic field lines anchored in the rotating disc and inclined at <60 deg fling ionized plasma outward via centrifugal force even well inside the Compton radius, operating alongside thermal expansion in hybrid magnetothermal winds[14].

| Driving Mechanism | | Primary Launch Force | | Favorable Accretion Conditions | | Typical Launch Region |

| Radiation Pressure | | Momentum transfer from photons to gas | | Super-Eddington luminosities | | Inner to mid-disc |

| Thermal (Compton) | | Gas pressure exceeding local escape velocity | | Strong X-ray irradiation | | Outer disc (beyond the Compton radius) |

| Magneto-Centrifugal | | Centrifugal force directed along magnetic field lines | | Strong, ordered poloidal magnetic fields | | Inner to outer disc |

Relativistic Jet Kinematics, Polarimetry, and Galactic Feedback

Very Long Baseline Array (VLBA) and Long Baseline Array (LBA) interferometry during the hard-intermediate state resolved a two-sided, asymmetrical north-south continuous radio jet spanning ~40 milliarcseconds -- the largest continuous jet ever imaged in a Galactic X-ray binary -- alongside a downstream shock knot 66.89 milliarcseconds south moving at 0.66 mas/hr[11][20]. Concurrently, optical polarimetry recorded a steady ~0.3% intrinsic polarization degree whose polarization angle rotated synchronously with discrete radio ejections due to electron scattering in the evolving disc wind, while IXPE X-ray polarimetry measured a sharp drop in coronal polarization during the soft state[7][21].

Just as supermassive black holes regulate galactic star formation via active galactic nucleus (AGN) feedback, the sustained mechanical energy injected by Swift J1727.8 1613's jets and late-stage winds inflates interstellar bubbles and ionizes surrounding gas clouds long after peak X-ray activity subsides[1][16][22]. Furthermore, because non-conservative mass loss strips substantial mass and orbital angular momentum from the binary before it can cross the event horizon, evolutionary models predicting black hole growth rates and gravitational-wave merger progenitors must explicitly incorporate persistent cool disc winds[4][23].

Black Holes as Inefficient Cosmic Engines

The multi-wavelength record of Swift J1727.8 1613's 2023 outburst establishes a complete observational synthesis of black hole accretion and expulsion. By linking inner-disc relativistic QPOs and record-breaking continuous radio jets with late-stage 800 km/s optical Balmer winds, this campaign proves that black holes continuously recycle a major fraction of their captured gas back into the cosmos, reshaping both binary star evolution and their galactic environments.

Works Cited

* University of Warwick (Jul 30, 2026). Black hole feeding frenzy ends in cosmic indigestion. -- https://warwick.ac.uk/news/pressreleases/black-hole-feeding/

* Remillard, R. A., & McClintock, J. E. (Sep 1, 2006). X-Ray Properties of Black-Hole Binaries. Annual Review of Astronomy and Astrophysics, 44, 49-92. -- https://doi.org/10.1146/annurev.astro.44.051905.092532

* Gizmodo (Aug 3, 2026). Even Black Holes Can Get Indigestion. -- https://gizmodo.com/even-black-holes-can-get-indigestion-unsurprisingly-their-farts-are-massive-expulsions-of-gas-2000794002

* AZoQuantum (Aug 3, 2026). Detailed Optical Records of Black Hole Eruption Cycle. -- https://www.azoquantum.com/News.aspx?newsID=11210

* Yu, W., Allak, S., Yang, Z.-X., Fan, X., Santangelo, A., & Xie, T.-h. (Jul 17, 2026). Stable X-ray reverberation lags in the black hole X-ray binary Swift J1727.8 1613. arXiv:2607.15729. -- https://arxiv.org/abs/2607.15729

* Solomons, K. S., & Castro Segura, N. (Apr 22, 2026). Outburst evolution of the black hole transient Swift J1727. PoS(DIM50th2025). -- https://doi.org/10.22323/1.493.0034

* Castro Segura, N., Solomons, K., Corral-Santana, J. M., Knigge, C., & Charles, P. A. (Jul 25, 2026). Optical outburst evolution of the transient black hole X-ray binary Swift J1727.8 1613: disc response to jet ejections and late-outburst emergence of powerful disc winds. Monthly Notices of the Royal Astronomical Society. -- https://doi.org/10.1093/mnras/stag1175

* Peng, J. Q., Zhang, S., Shui, Q. C., et al. (Jan 1, 2024). NICER, NuSTAR, and Insight-HXMT Views to the Newly Discovered Black Hole X-Ray Binary Swift J1727.8-1613. The Astrophysical Journal Letters. -- https://doi.org/10.3847/2041-8213/ad17ca

* Georganti, M., Knigge, C., et al. (Nov 11, 2025). Ultraviolet spectroscopy of the black hole X-ray binary MAXI J1820+070 across a state transition. Monthly Notices of the Royal Astronomical Society. -- https://doi.org/10.1093/mnras/staf1965

* Yu, W., Bu, Q.-C., Zhang, S.-N., et al. (Mar 19, 2024). Timing analysis of the newly discovered black hole candidate Swift J1727.8-1613 with Insight-HXMT. arXiv:2403.13127. -- https://arxiv.org/abs/2403.13127

* Wood, C. M., Miller-Jones, J. C. A., et al. (Aug 1, 2024). Swift J1727.8-1613 Has the Largest Resolved Continuous Jet Ever Seen in an X-Ray Binary. The Astrophysical Journal Letters. -- https://doi.org/10.3847/2041-8213/ad6572

* Mereminskiy, I., Lutovinov, A., Molkov, S., et al. (Jun 6, 2024). Hard X-rays and QPO in Swift J1727.8 1613: the rise and plateau of the 2023 outburst. Monthly Notices of the Royal Astronomical Society, 531(4), 4893-4899. -- https://doi.org/10.1093/mnras/stae1393

* Jin, P., M ndez, M., et al. (Oct 11, 2025). Black-hole X-ray binary Swift J1727.8 1613 shows simultaneous Type-B and Type-C QPOs. arXiv:2510.10353. -- https://arxiv.org/abs/2510.10353

* Waters, T., & Proga, D. (Sep 4, 2018). Magnetothermal disc winds in X-ray binaries: poloidal magnetic fields suppress thermal winds. Monthly Notices of the Royal Astronomical Society. -- https://doi.org/10.1093/mnras/sty2398

* Mu oz-Darias, T., Casares, J., et al. (May 9, 2016). Regulation of black-hole accretion by a disk wind during a violent outburst of V404 Cygni. Nature. -- https://doi.org/10.1038/nature17446

* Tech Explorist (Jul 31, 2026). Scientists discover black holes have a cosmic digestive system. -- https://www.techexplorist.com/black-holes-cosmic-digestive-system/103643/

* Mu oz-Darias, T., & D az Trigo, M. (Jan 8, 2026). Accretion disc winds in X-ray binaries. arXiv:2601.05319. -- https://arxiv.org/abs/2601.05319

* Higginbottom, N., Knigge, C., et al. (Jun 15, 2018). Radiation-hydrodynamic simulations of thermally driven disc winds in X-ray binaries: a direct comparison to GRO J1655 40. Monthly Notices of the Royal Astronomical Society. -- https://doi.org/10.1093/mnras/sty1599

* Higginbottom, N., Knigge, C., et al. (Jan 25, 2020). Thermal and radiation driving can produce observable disc winds in hard-state X-ray binaries. Monthly Notices of the Royal Astronomical Society. -- https://doi.org/10.1093/mnras/staa209

* Peng, J. Q., Zhang, S., et al. (Jan 9, 2025). A possible jet and corona configuration for Swift J1727.8-1613 during the hard state. Journal of High Energy Astrophysics. -- https://doi.org/10.1016/j.jheap.2025.01.003

* Liu, H. X., Xu, Y.-J., et al. (May 29, 2026). The broad-band X-ray spectral properties during the rising phases of the outburst of the new black hole X-ray binary candidate Swift J1727.8-1613. Monthly Notices of the Royal Astronomical Society. -- https://doi.org/10.1093/mnras/stag999

* Hu ko, F., Lacey, C. G., et al. (Jan 25, 2025). The effects of super-Eddington accretion and feedback on the growth of early supermassive black holes and galaxies. Monthly Notices of the Royal Astronomical Society. -- https://doi.org/10.1093/mnras/staf146

* ScienceDaily (Jul 31, 2026). Black holes may spit out nearly as much matter as they swallow. -- https://www.sciencedaily.com/releases/2026/07/260731034127.htm