Health & Medicine | Jun 3, 2026 | 10 min read | By Bryan White

Breaching the Barrier: What the 2026 Texas Screwworm Outbreak Means for US Agriculture

Climate Change * Public Health * Animals

1. Introduction: Breach of the U.S. Southern Border in Zavala County

On June 3, 2026, the Texas Animal Health Commission and USDA Animal and Plant Health Inspection Service (APHIS) confirmed an active New World screwworm (Cochliomyia hominivorax) infestation in a three-week-old beef calf near La Pryor in Zavala County, Texas -- roughly 50 miles north of the Rio Grande.[1][2] Verified by the National Veterinary Services Laboratories from umbilical lesion larvae, the case represents the first Texas livestock incursion since the 1960s eradication campaign.[1][2]

Following the 2023 collapse of Panama's Darien Gap barrier, C. hominivorax spread across Central America and Mexico, generating over 26,200 cumulative Mexican cases (with over 2,700 active infections monitored in mid-2026) and reaching a goat in Coahuila 25 miles from Texas on May 28, 2026.[4][6][7] Containing the breach requires combining classic epidemiology, area-wide Sterile Insect Technique (SIT) releases, and transgenic male-only biocontrol.[1][43]

2. Developmental Biology and Pathophysiology of Traumatic Myiasis

Unlike necrophagous calliphorids such as Cochliomyia macellaria, C. hominivorax is an obligate parasite of living warm-blooded vertebrates.[3][8] Adult females detect volatile compounds from open wounds, tick bites, shearing or branding abrasions, and neonatal umbilici, depositing shingle-like clusters of 200-400 eggs.[5][8] Crucially for genetic control, wild females mate only once during their 10-to-30-day lifespan, whereas males are polygamous.[5]

| Developmental Stage | | Approximate Duration | | Biological Characteristics and Host Interaction |

| Embryonic Stage (Egg) | | 10 to 21 hours | | Shingle-like egg clusters undergo rapid embryogenesis driven by mammalian body heat. |

| First-Instar Larva | | 1 to 2 days | | Emerging larvae use chitinous mouth hooks to burrow into live epithelial and subcutaneous tissues. |

| Second & Third-Instar Larva | | 5 to 7 days | | Rapid growth and intensive tissue consumption head-downward, forming a pocket of liquefactive necrosis. |

| Pupal Stage | | 7 to 54 days | | Mature third-instar larvae drop to the ground and pupate in topsoil; duration depends on soil temperature and moisture. |

| Adult Emergence | | 10 to 30 days | | Adults feed on nectar and wound fluids, reaching sexual maturity within three to five days. |

This holometabolous developmental biology causes severe traumatic myiasis.[8][12] Larvae tear into viable tissue with mouth hooks while secreting proteolytic enzymes

(proteases) that degrade host proteins.[8][9][13] Secondary bacterial infection triggers sepsis and toxemia, causing near-100% mortality in untreated livestock.[3][14] In humans, myiasis in chronic ulcers, nasal/aural cavities, or newborn umbilici poses a severe public health threat, underscored by a 2025 travel-associated case in Maryland.[4][10][14][15]

3. Incident Command Response and Macroeconomic Impact Modeling

Prior to 2026, the only recent U.S. incursion was the 2016-2017 Florida Keys outbreak in endangered Key deer (Odocoileus virginianus clavium), eradicated with 188 million sterile flies.[20][22] Following the June 3, 2026 Zavala County confirmation, Texas State Veterinarian Dr. Bud Dinges and USDA APHIS activated emergency containment around La Pryor.[1]

| Incident Response Protocol | | Specific Action Implemented in Zavala County (June 2026) |

| Quarantine Establishment | | 20-kilometer radius infested zone established around La Pryor, Texas. |

| Movement Restrictions | | Prohibition on moving warm-blooded animals outside the quarantine zone without veterinary inspection. |

| Sterile Fly Dispersal | | Ground release chambers and aerial drops dispersing ~4 million sterile flies per week. |

| Surveillance Expansion | | Expanded fly traps, border tick riders, and detector dogs intercepting wildlife. |

| Public Health Coordination | | Clinical and veterinary alerts for myiasis signs across southern Texas. |

With Texas holding over 12 million cattle and 655,000 sheep and goats amid record beef prices, USDA APHIS modeling projects severe economic damage if the outbreak spreads.[19][36]

| Economic Metric | | 2024 APHIS Projection | | Primary Driving Factors |

| Direct Producer Losses | | $732 Million (Annualized) | | Neonate mortality, adult weight loss, hide damage, veterinary treatment, and larvicides. |

| Operational Labor Expansion | | Integrated into Producer Losses | | Daily herd inspection labor, isolated herd management, and suspended shearing/castration. |

| Total State Economic Damage | | $1.8 Billion (Annualized) | | Supply chain bottlenecks, reduced slaughter throughput, auction closures, and export halts. |

| National Extrapolation Risk | | Up to $10.6 Billion | | Long-term damages if the pest re-establishes across the southern United States. |

4. Sterile Insect Technique and Next-Generation Transgenic Sexing Strains

Pioneered by Edward F. Knipling and Raymond C. Bushland and proven on Cura ao in 1954, traditional SIT rears larvae on spray-dried blood, egg, and milk diets before sterilizing five-to-six-day-old pupae with ~55 Gray of gamma radiation from

Cesium-137 or Cobalt-60.[5][34][40] However, co-releasing sterile females doubles diet costs, distracts sterile males from wild females, and requires higher female-sterilizing radiation doses that impair male fitness.[39][43][44]

To produce male-only cohorts, North Carolina State University researchers led by Dr. Max J. Scott, with USDA-ARS and COPEG, engineered transgenic Tet-off gene expression strains (DR6, DR7, and "NovoFly").[43][44][50] Using molecular genetics, early embryonic promoters (Lucilia cuprina nullo in DR6 or C. macellaria CG14427 in DR7) drive a tetracycline transactivator (tTA) cassette gated by the sex-specific introns and alternative RNA splicing of the blowfly transformer (tra) gene.[50][51]

With dietary doxycycline present during colony maintenance, tTA transcription is repressed and both sexes survive.[44][50] When field cohorts are reared without doxycycline, female-specific tra splicing triggers lethal tTA overexpression in female embryos, yielding 100% males that require lower radiation doses and provide secondary biosecurity in the wild ecosystem.[44][50][54]

| System Characteristic | | Traditional Sterile Insect Technique | | Transgenic Male-Only Technique (Tet-Off) |

| Release Population | | Mixed (Males and Females) | | Male Only |

| Dietary Efficiency | | Low (resources consumed by sterile females) | | High (females eliminated at embryonic stage) |

| Radiation Dosage | | High (dictated by female resistance levels) | | Optimized / Lower (tailored specifically to males) |

| Field Mating Efficiency | | Reduced (males expend energy courting sterile females) | | High (males seek out strictly wild females) |

| Secondary Biosecurity | | None | | High (escaped fertile males produce only male offspring in the wild) |

5. Conclusion: Domestic SIT Infrastructure and Strategic Policy

Backed by USDA Secretary Brooke Rollins' five-pronged strategy -- including construction of a 300-million-fly-per-week sterile facility in Edinburg, Texas, and the $100 million NWS Grand Challenge -- containing the 2026 Zavala County outbreak requires autonomous domestic SIT capacity and continuous genetic evolution in pest control.[19][31][57]

Works Cited

* USDA Confirms New World Screwworm Detection in Texas Calf, accessed June 3, 2026, https://westernagnetwork.com/usda-confirms-new-world-screwworm-detection-in-texas-calf

* New World screwworm confirmed in Zavala County calf, accessed June 3, 2026, https://www.ntxe-news.com/artman/publish/article_145883.shtml

* Understanding the New World Screwworm - Biology, Identification, and Prevention, accessed June 3, 2026, https://extension.psu.edu/understanding-the-new-world-screwworm-biology-identification-and-prevention

* The New World Screwworm in the United States: A Narrative Review ..., accessed June 3, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC12591281/

* New World Screwworm Ready Reference Guide Sterile Insect Response - usda aphis, accessed June 3, 2026, https://www.aphis.usda.gov/sites/default/files/nws_rrg_sterileinsectresponse.pdf

* Flesh-eating New World screwworm found in Texas calf, USDA says - CBS News, accessed June 3, 2026, https://www.cbsnews.com/news/flesh-eating-new-world-screwworm-found-texas-calf-usda/

* Flesh-eating screwworm detected 25 miles from U.S. border, USDA says - WCBI, accessed June 3, 2026, https://www.wcbi.com/flesh-eating-screwworm-detected-25-miles-from-u-s-border-usda-says/

* DPDx - New World Screwworm - CDC, accessed June 3, 2026, https://www.cdc.gov/dpdx/newworldscrewwormmyiasis/index.html

* New World Screwworm Prevention for Animals - usda aphis, accessed June 3, 2026, https://www.aphis.usda.gov/livestock-poultry-disease/cattle/ticks/screwworm

* Myiasis Caused by Old and New World Screwworms - The Center for Food Security and Public Health, accessed June 3, 2026, https://www.cfsph.iastate.edu/Factsheets/pdfs/screwworm_myiasis.pdf

* New World Screwworm - Integumentary System - Merck Veterinary Manual, accessed June 3, 2026, https://www.merckvetmanual.com/integumentary-system/flies/new-world-screwworm

* Screwworm flies as agents of wound myiasis - Food and Agriculture Organization of the United Nations, accessed June 3, 2026, https://www.fao.org/4/u4220t/u4220t07.htm

* About New World Screwworm - CDC, accessed June 3, 2026, https://www.cdc.gov/new-world-screwworm/about/index.html

* Myiasis - PMC - NIH, accessed June 3, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC3255963/

* Umbilical Myiasis by Cochliomyia hominivorax in an Infant in Colombia - PMC - NIH, accessed June 3, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC6987426/

* Case Report: Myiasis due to Cochliomyia hominivorax and Dermatobia hominis: Clinical and Pathological Differences between Two Species in Northern Peru - PMC, accessed June 3, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5928687/

* Anorectal Myiasis Caused by Cochliomyia hominivorax in a Septic Tank Worker - PMC - NIH, accessed June 3, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC12454922/

* Introduction * STOP Screwworms: Selections from the Screwworm Eradication Collection * - National Agricultural Library, accessed June 3, 2026, https://www.nal.usda.gov/exhibits/speccoll/exhibits/show/stop-screwworms--selections-fr/introduction

* NWS Economic Impact Report - USDA-Aphis, accessed June 3, 2026, https://www.aphis.usda.gov/sites/default/files/nws-historical-economic-impact.pdf

* Final Report for the APHIS Veterinary Services Response to the 2016-2017 Outbreak of New World Screwworm (NWS) in Florida, accessed June 3, 2026, https://www.aphis.usda.gov/sites/default/files/public-nws-usdaaphis-final-report.pdf

* Florida Key Deer Screwworm Final Report (Phase I) - Texas A&M NRI, accessed June

3, 2026, https://nri.tamu.edu/publications/research-reports/2016/florida-key-deer-screwworm-final-report-phase-i/

* Screwworm (Diptera: Calliphoridae) in the United States: Response to and Elimination of the 2016-2017 Outbreak in Florida - PubMed, accessed June 3, 2026, https://pubmed.ncbi.nlm.nih.gov/29722869/

* Screwworm infestation kills endangered deer | American Veterinary Medical Association, accessed June 3, 2026, https://www.avma.org/javma-news/2017-01-15/screwworm-infestation-kills-endangered-deer

* Screwworm Outbreak: Florida 2016 - 2017 - United States Animal Health Association, accessed June 3, 2026, https://usaha.org/upload/Meetings/2017/Presentations/1_3_Welch_Parasitic_Dis_2017.pdf

* Possible flesh-eating screwworm case in Texas after pest detected 25 miles from U.S. border, USDA says, accessed June 3, 2026, https://www.cbsnews.com/news/flesh-eating-screwworm-detected-25-miles-us-border-usda/

* With screwworm now closer than ever to Texas border, state ramps up efforts to fight the pest, accessed June 3, 2026, https://www.kvue.com/article/news/state/texas-news/screwworm-closest-us-state-federal-government-actions-slow-spread-pest/269-

* Flesh-eating New World screwworm found in Texas calf, USDA says, accessed June 3, 2026, https://www.youtube.com/watch?v=m2V7r4vHM1A

* USDA suspects New World screwworm fly has arrived in Texas | AP ..., accessed June 3, 2026, https://apnews.com/article/screwworm-flesh-eating-parasite-cattle-texas-2efc5ec69d9651b5c0bab4825eda4976

* New World Screwworm confirmed in South Texas calf, state and federal officials respond, accessed June 3, 2026, https://www.wfaa.com/article/news/state/texas-news/new-world-screwworm-confirmed-in-south-texas-calf-state-and-federal-official

* First US screwworm case confirmed in South Texas | kvue.com, accessed June 3, 2026, https://www.kvue.com/article/news/state/texas-news/new-world-screwworm-texas-reported-case/269-a0f1d261-b183-4968-81ec-aaa

* TAHC Supports Comprehensive New World Screwworm Eradication Actions Announced Today, accessed June 3, 2026, https://www.tahc.texas.gov/news/2025/2025-08-15_NWS-FederalSupport.pdf

* Texas gears up for New World Screwworm threat - Michigan Farm News, accessed June 3, 2026, https://www.michiganfarmnews.com/texas-gears-up-for-new-world-screwworm-threat

* June 25, 2025 Dr. Lewis R. "Bud" Dinges Executive Director and State Veterinarian Texas Animal Health Commission 2105 Kram, accessed June 3, 2026, https://gov.texas.gov/uploads/files/press/NWS_Response_Team_Directive_FINAL.pdf

* The development and application of the sterile insect technique (SIT) for New World screwworm eradication - Food and Agriculture Organization of the United Nations, accessed June 3, 2026, https://www.fao.org/4/u4220t/u4220T0j.htm

* History of screwworms in the U.S. | Veterinary Medicine Extension | Washington State University, accessed June 3, 2026,

https://vetextension.wsu.edu/2025/07/17/history-of-screwworms-in-the-u-s/

* US steps up response to screwworm threat | American Veterinary Medical Association, accessed June 3, 2026, https://www.avma.org/news/us-steps-response-screwworm-threat

* New World Screwworm and Livestock Border Closures: A Guide for New Mexico Livestock Producers - aces.nmsu.edu, accessed June 3, 2026, https://pubs.nmsu.edu/_z/Z131.pdf

* 1992: Knipling and Bushland - The World Food Prize, accessed June 3, 2026, https://www.worldfoodprize.org/en/laureates/19871999_laureates/1992_knipling_and_bushland/

* Sterile insect technique - Wikipedia, accessed June 3, 2026, https://en.wikipedia.org/wiki/Sterile_insect_technique

* The Sterile Insect Release Method and Other Genetic Control Strategies, accessed June 3, 2026, https://ipmworld.umn.edu/bartlett

* Eradicating New World Screwworm with Sterile Insect Technique - USDA-Aphis, accessed June 3, 2026, https://www.aphis.usda.gov/sites/default/files/factsheet-eradicating-nws-sit.pdf

* Disease name - WOAH, accessed June 3, 2026, https://www.woah.org/fileadmin/Home/eng/Our_scientific_expertise/reflabreports2009/2009_Reports/1_REPORTS/NEW_WORLD

* It's a Boy: Modified Male Flies Could More Efficiently Control Screwworm Population, accessed June 3, 2026, https://ges.research.ncsu.edu/2016/09/its-a-boy-modified-male-flies-could-more-efficiently-control-screwworm-population/

* NovoFly comment Max Scott - Regulations.gov, accessed June 3, 2026, https://downloads.regulations.gov/EPA-HQ-OPP-2026-0991-0016/attachment_1.pdf

* Male-only strains of the livestock pests, Cochliomyia hominivorax and Lucilia cuprina, accessed June 3, 2026, https://maxscottlab.wordpress.ncsu.edu/sample-page/male-only-strains-of-the-livestock-pests-cochliomyia-hominivorax-and-lucilia-

* Sterile Insect Technique (SIT) and Its Applications - PMC - NIH, accessed June 3, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC8304793/

* Screwworm Male-Only Lines: Past, Present and Future w/ Dr. Maxwell Scott - YouTube, accessed June 3, 2026, https://www.youtube.com/watch?v=ON960H5VFgI

* A transgenic male-only strain of the New World screwworm for an improved control program using the sterile insect technique - PubMed, accessed June 3, 2026, https://pubmed.ncbi.nlm.nih.gov/27576512/

* Achievement: New Technology - IGERT, accessed June 3, 2026, https://www.igert.org/spotlights/4614.html

* Development and evaluation of screwworm, Cochliomyia hominivorax, transgenic sexing strains with embryonic gene promoters for a genetic control program | bioRxiv, accessed June 3, 2026, https://www.biorxiv.org/content/10.1101/2025.05.01.651745

* A transgenic male-only strain of the New World screwworm for an improved control program using the sterile insect technique - PMC, accessed June 3, 2026,

https://pmc.ncbi.nlm.nih.gov/articles/PMC5004303/

* Development and evaluation of screwworm, Cochliomyia hominivorax (Diptera: Calliphoridae), transgenic sexing strains with embryonic gene promoters for a genetic control program | Journal of Economic Entomology | Oxford Academic, accessed June 3, 2026, https://academic.oup.com/jee/article/119/2/1457/8307731

* Doxycycline is a viable alternative to tetracycline for use in insect Tet-Off transgenic sexing systems, as assessed in the blowflies Cochliomyia hominivorax and Lucilia cuprina (Diptera: Calliphoridae) - PubMed, accessed June 3, 2026, https://pubmed.ncbi.nlm.nih.gov/38401129/

* COPEG funds New World screwworm project - Max Scott Lab, accessed June 3, 2026, https://maxscottlab.wordpress.ncsu.edu/2015/02/19/copeg-funds-new-world-screwworm-project/

* An early female lethal system of the New World screwworm, Cochliomyia hominivorax, for biotechnology-enhanced SIT - PMC, accessed June 3, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC7747452/

* A transgenic male-only strain of the New World screwworm for an improved control program using the sterile insect technique - Publication : USDA ARS, accessed June 3, 2026, https://www.ars.usda.gov/research/publications/publication/?seqNo115=328078

* Screwworm.gov | Unified Government Response To Protect the United States - usda aphis, accessed June 3, 2026, https://www.aphis.usda.gov/animalsanimal-health/livestock-and-poultry-disease/stop-screwworm