Natural Science | Oct 31, 2026 | 5 min read | By Bryan White

Cambridge Impact Simulations Challenge the Theory That Comets Kick-Started Life on Earth

Astrobiology * Abiogenesis * Comets

For more than six decades -- since Joan Or proposed in 1961 that volatile-rich comets could have delivered the carbon and nitrogen feedstocks needed to assemble the first nucleic acids -- planetary scientists have frequently cast comets as early Earth's primary prebiotic couriers. Space missions including the European Space Agency's Rosetta orbiter at Comet 67P/Churyumov-Gerasimenko have confirmed that cometary nuclei are rich in hydrogen cyanide (HCN), glycine, and phosphorus -bearing minerals. Yet a new quantitative study published on October 28, 2026, in Scientific Reports by astrophysicist Dr. Richard J. Anslow and colleagues at the University of Cambridge's Institute of Astronomy reveals a severe physical bottleneck: while comets carry abundant organic precursors through interplanetary space, the mechanics of atmospheric entry and hypervelocity impact destroy or disperse almost all of that cargo before it can accumulate in surface waters on an Earth-mass planet.

1. The Prebiotic Feedstock Bottleneck: Why Hydrogen Cyanide Requires Intact, Concentrated Delivery

In contemporary origin-of-life chemistry, hydrogen cyanide (HCN) occupies a central mechanistic role. Under ultraviolet irradiation in the presence of dissolvedbisulfite and copper or ironcyanide catalysts -- the cyanosulfidic protometabolic network elucidated by John Sutherland and colleagues -- concentrated aqueous HCN drives the divergent, high-yield synthesis of canonical ribonucleotides, amino acids, and lipid precursors. However, these bimolecular condensation reactions are strictly concentration-dependent: laboratory syntheses typically require local pond or evaporative lagoon concentrations of at least 1 to 100 millimolar (mM) HCN before competitive hydrolysis converts HCN into formamide and ammonium formate.

For a comet to jump-start cyanosulfidic chemistry on the Hadean Earth (approximately 4.3 to 3.8 billion years ago), two physical conditions must be satisfied simultaneously. First, the comet must traverse the atmosphere and strike exposed continental crust or a shallow subaerial basin rather than diluting its payload into the global Hadean ocean. Second, the impact velocity must be low enough that shock heating does not thermally dissociate the C N triple bond into equilibrium gases (N 2 , CO, CO 2 , and H 2 O).

| Impactor Class / Radius | | Entry Velocity (v entry) | | Atmospheric & Shock Regime | | Peak Shock Temp | | Intact HCN Pond Delivery |

| Small Comets (R < 100 m) | | 15 - 55 km/s | | Stratospheric pancake airburst & ablation | | 1,200 - 2,800 K | | Global atmospheric dilution (no local pond) |

| Typical Comets (R = 0.5 - 5 km) | | 20 - 70 km/s | | Unbraked hypervelocity cratering shock | | > 3,500 - 6,000 K | | < 0.01% survival (complete pyrolysis) |

| Low-Velocity Tail Comets (R ~ 0.2 - 1 km) | | 11.2 - 14.5 km/s | | Aerobraked

fragment train / oblique impact | | 900 - 1,500 K | | 0 copies in 54% of 100,000 Earth histories |

| Carbonaceous Asteroids (C-type) | | 11.2 - 17.0 km/s | | Higher tensile strength + lower orbital eccentricity | | 800 - 1,800 K | | Order-of-magnitude higher intact survival |

2. The Atmospheric Catch-22: Stratospheric Airbursts vs. Hypervelocity Shock Pyrolysis

To test whether cometary bombardment could reliably seed Hadean ponds, Anslow, Amy Bonsor, and Todd A. Thompson coupled a dynamical orbital population model of scattered-disk and Jupiter-family comets with a semi-analytical pancake fragmentation model of atmospheric entry and a shock- thermodynamics survival parameterization. Their calculations expose a fundamental physical catch-22 governed by Earth's gravitational escape velocity (v esc = 11.2 km/s) and the low structural strength of porous cometary ice (~10 [3] to 10 [4] Pa):

* Aerodynamic Disruption of Small Nuclei: Comets smaller than roughly 100 to 200 meters in radius experience ram pressures (P ram = atm v [2]) in the upper atmosphere that exceed their bulk tensile strength by several orders of magnitude. Instead of decelerating intact to terminal velocity, these nuclei flatten and detonate in high-altitude airbursts. Even if a fraction of their HCN survives the radiative fireball, it is dispersed across the stratosphere rather than concentrated into a single warm little pond.

* Shock Pyrolysis of Kilometre-Scale Nuclei: Comets larger than ~500 meters in radius possess enough column mass to punch through a 1- to 10-bar Hadean atmosphere with negligible aerodynamic deceleration. Because comets originate on high-eccentricity, high-inclination orbits from the outer Solar System , their typical Earth-crossing entry speeds cluster between 20 and 50 km/s. At impact velocities above ~15 km/s, the peak shock pressure exceeds 50 gigapascals (GPa) and post-shock temperatures climb above 3,000 to 5,000 K, thermally decomposing more than 99.99% of cometary HCN.

3. 100,000 Simulated Bombardment Histories: Why Cometary Seeding Is a Statistical Fluke

Because only the extreme low-velocity tail of the cometary impact distribution (v imp < 14 to 15 km/s) can deliver intact HCN without catastrophic vaporization, the Cambridge team ran 100,000 Monte Carlo realizations of early Earth's cometary bombardment history. They accounted for the total cometary mass flux during the Hadean eon alongside geological estimates that exposed subaerial land covered only 1% to 10% of early Earth's surface prior to widespread cratonic continent formation.

In 54% of the 100,000 simulated Earth histories, not a single comet successfully delivered an intact, concentrated payload of HCN onto subaerial land . Even in the remaining simulations where one or two low-velocity, oblique cometary impacts occurred over hundreds of millions of years, the authors note that relying on a solitary stochastic impact into an ephemeral crater lake represents a "one-off fluke" rather than a planetary mechanism capable of sustaining multi-step prebiotic evolution .

4. Implications for Endogenous Geochemistry, Asteroids, and Exoplanetary Biosignatures

Rather than leaving prebiotic chemistry without a feedstock source, the Cambridge findings sharpen focus on three far more robust pathways for Hadean and exoplanetary abiogenesis :

* Carbonaceous Chondrite Asteroids and Interplanetary Dust Particles (IDPs): Main-belt C-type asteroids (such as the parent bodies of the Murchison meteorite and asteroid 101955 Bennu, sampled by NASA's OSIRIS-REx mission) orbit on lower-eccentricity paths that intersect Earth at significantly lower average speeds (12 to 17 km/s) and possess higher mechanical strength than cometary snowballs, allowing carbonaceous meteorites and micrometeorites to deliver intact amino acids, nucleobases, and soluble phosphate at a steady background flux.

* Impact-Generated Reducing Atmospheres: Even when a large iron -rich asteroid or comet vaporizes completely on impact, its metallic iron core reduces oceanic water to molecular hydrogen (H 2), creating a transient H 2 -CH 4 -NH 3 atmosphere in which solar UV photolysis and lightning discharge synthesize HCN endogenously across the globe, followed by rainout into continental basins.

* Rocky Exoplanets in Compact Systems: In tightly packed multi-planet systems such as TRAPPIST-1 -- where higher orbital velocities around low-mass M-dwarf stars make intact cometary delivery even less probable -- Anslow and colleagues emphasize that endogenous geochemistry and low-velocity planetesimal accretion provide a much more reliable foundation for prebiotic chemistry than outer-system cometary scattering.

References & Primary Sources

* [1] Anslow, R. J., Bonsor, A., & Thompson, T. A. (2026). Cometary impacts an unlikely pathway towards the origins of life on Earth. Scientific Reports, 16, 31688. -- https://doi.org/10.1038/s41598-026-74214-y

* [2] University of Cambridge Institute of Astronomy (Oct 28, 2026). Comets unlikely to have kick-started life on Earth, study finds. -- https://www.cam.ac.uk/research/news

* [3] Patel, B. H., Percivalle, C., Ritson, D. J., Duffy, C. D., & Sutherland, J. D. (2015). Common origins of RNA, protein and lipid precursors in a cyanosulfidic protometabolism. Nature Chemistry, 7(4), 301-307. -- https://doi.org/10.1038/nchem.2202

* [4] Altwegg, K., Balsiger, H., Bar-Nun, A., et al. (2016). Prebiotic chemicals -- amino acid and phosphorus -- in the coma of comet 67P/Churyumov-Gerasimenko. Science Advances, 2(5), e1600285. -- https://doi.org/10.1126/sciadv.1600285