Climate & Earth Systems | Feb 21, 2026 | 8 min read | By Bryan White

Cracking the Pacific Puzzle: Why Part of the Ocean is Cooling While the World Warms

Environment * Climate Change * Physics

Introduction: The Model-Observation Divergence in the Pacific

For over a decade, a stark contradiction between observational records and coupled climate simulations has stood as a major unresolved problem in physical climatology. While global mean temperatures have risen steadily under anthropogenic greenhouse gas forcing, satellite-era reconstructions (1979-2014) document persistent multidecadal cooling across the eastern tropical Pacific cold tongue and the Pacific sector of the Southern Ocean.[1][2] Conversely, coarse-resolution Coupled Model Intercomparison Project Phase 6 (CMIP6) models almost universally simulate surface warming across these basins.[1][6]

Termed the "Pacific puzzle," this divergence impacts global climate sensitivity and regional hydroclimate projections because tropical Pacific sea surface temperature (SST) gradients act as a global pacemaker, and simulating eastern Pacific warming causes coarse ensembles to overestimate historical global surface warming.[1][2][7] Researchers at the Max Planck Institute for Meteorology recently resolved this discrepancy using a kilometer-scale coupled configuration of the ICOsahedral Non-hydrostatic (ICON) Earth System Model, demonstrating how explicitly resolved Southern Ocean mesoscale eddies, Andean topographic blocking, Peruvian stratocumulus clouds, and the equatorial Bjerknes feedback drive this interhemispheric cooling pathway.[1][2]

Historical Context and Competing Theoretical Frameworks

Between 1979 and 2014, ERSSTv5 and ERA5 datasets recorded a La Ni a-like trend despite rising atmospheric carbon dioxide: western Pacific warming paired with eastern Pacific cooling sharpened the zonal SST gradient, accelerated easterly trade winds, and shoaled the eastern thermocline, whereas CMIP6 models predicted an El Ni o-like weakening of upwelling.[2][4][6][9] Prior hypotheses struggled to bridge this gap:[5][9]

* Transient Ocean Thermostat: Upwelling of un-warmed deep water buffers the eastern Pacific against thermodynamic warming and strengthens the Walker circulation, but coarse models lack synchronous coupling and sufficient heat-flux damping.[8]

* Internal Variability (IPO): Attributes cooling to the Interdecadal Pacific Oscillation, yet the 35-year trend exceeds ensemble internal variability bounds.[9]

* Antarctic Meltwater & Aerosols: Antarctic ice sheet meltwater stratifies the Southern Ocean and expands sea ice, while aerosols reflect solar radiation; however, cooling persisted as aerosol emissions declined.[10][12]

* Land-Sea Heating Contrast: Continental warming under radiative forcing excites Rossby wave trains that strengthen subtropical highs.[5]

| Theoretical Framework | | Core Physical Mechanism | | Structural Limitation in Coarse CMIP6 Models |

| Transient Ocean Thermostat | | Upwelling of deep un-warmed water strengthens zonal gradient | | Asynchronous coupling and weak atmospheric heat flux damping[8] |

| Internal Variability (IPO) | | Multidecadal oscillation masks anthropogenic warming | | Observed trend duration exceeds modeled internal noise[9] |

| Meltwater / Aerosols / Land-Sea | | Freshwater stratification, aerosols, or Rossby waves | | Weak teleconnections and poorly parameterized stratocumulus feedbacks[5][10][12] |

The Exascale Paradigm Shift: ICON and the WarmWorld Initiative

Standard CMIP6 models (~100 km grid) rely on parameterizations for ocean eddies, convection, and steep terrain.[2][5] Using the unified ICON model, researchers ran a coupled HighResMIP simulation with a 10 km atmospheric grid (R2B8, 90 vertical levels) and a 5 km eddy-rich ocean grid (R2B9, 128 z* levels).[2][13][14][16][17]

| Model & WarmWorld Module | | Standard CMIP6 Ensembles | | Kilometer-Scale ICON Configuration (WarmWorld) |

| Grid & Physics Resolution | | ~100 km grid; parameterized eddies & convection | | 10 km Atmosphere / 5 km Ocean; explicit non-hydrostatic physics[2][17] |

| WarmWorld "Better" & "Faster" | | Legacy CPU Fortran solvers; cloud biases | | Turbulent energy schemes; OpenACC GPU refactoring (Gordon Bell Prize)[16][18][22] |

| WarmWorld "Easier" & "Smarter" | | Coarse file post-processing bottlenecks | | Petabyte Zarr/HEALPix/STAC workflows and in-situ memory analytics[19][21][23] |

Supported by Germany's WarmWorld initiative, NextGEMS, and Destination Earth on the Levante and JUWELS supercomputers, ICON reproduced the observed 0.56 K global mean surface warming over the 35-year study period alongside Southern and eastern Pacific cooling.[2][15][18][19][20]

The Four-Stage Interhemispheric Teleconnection Mechanism

The kilometer-scale simulation uncovered a four-stage interhemispheric pathway:[1][2]

* Southern Ocean Eddy Heat Export: Despite rising downward atmospheric heat flux, the Pacific sector of the Southern Ocean cools because 5 km resolved mesoscale eddies pump heat into the deep ocean across Antarctic Circumpolar Current (ACC) fronts and shift ACC fronts northward, expanding cold polar water outcropping.[2][11]

* Equatorward Advection and Topographic Blocking: Cold Southern Ocean SSTs induce a high-pressure anomaly that strengthens southeasterly winds along South America, carrying cooling northward via evaporative latent heat flux.[2] Explicit 10 km resolution of the steep Andes blocks warm continental air, preserving a coastal marine inversion and cold Ekman transport upwelling.[1][2]

* Peruvian Stratocumulus Cloud Amplification: Cold coastal waters and a strong inversion thicken the high-albedo Peruvian stratocumulus cloud deck, reflecting

shortwave sunlight to amplify cooling.[1][2][5]

* Equatorial Propagation via the Bjerknes Feedback: Southeastern Pacific cooling sharpens the equatorial east-west SST gradient, triggering the El Ni o-Southern Oscillation Bjerknes feedback.[5][24][25][27] Stronger easterly trade winds shoal the eastern thermocline, enhance cold upwelling, and propagate the cold tongue westward along the ITCZ.[2][5][11][26]

| Sequential Stage | | Geographic Domain | | Governing Physical Process | | Thermodynamic & Dynamical Outcome |

| 1. Eddy Export | | Southern Ocean | | Resolved 5 km eddies export heat across ACC fronts | | Surface cooling & northward ACC shift[2] |

| 2. Advection & Andes | | Southeastern Pacific | | High pressure strengthens trades; Andes block warm air | | Latent cooling advected north; Ekman upwelling[1][2] |

| 3. Cloud Feedback | | Peruvian Coast | | Strong inversion thickens stratocumulus deck | | Shortwave solar reflection amplifies cooling[1][2][5] |

| 4. Bjerknes Loop | | Equatorial Pacific | | Zonal gradient accelerates trades & shoals thermocline | | Westward-propagating La Ni a-like cold patch[2][5] |

Implications for Global Energy Budgets and Future Climate Projections

By capturing Southern Ocean eddy heat export and Peruvian cloud feedbacks, ICON corrects the global radiation budget biases of coarse CMIP6 ensembles.[2] Researchers must now determine whether this La Ni a-like cooling will persist -- sustaining western North American drought risk -- or whether greenhouse forcing will eventually destabilize stratocumulus decks and trigger rapid catch-up warming.[1][3][7][28]

Conclusion: Grounding Climate Projections in Explicit Physics

Resolving the Pacific puzzle proves that kilometer-scale representation of Southern Ocean eddies, Andean topography, stratocumulus clouds, and Bjerknes coupling is indispensable for next-generation climate projections.[1][2]

Updates... (Nov 21, 2026): Replaced the HTTP 403-blocked AMS portal URL in Ref. 12 with its verified CrossRef DOI (Dong et al., Oct 1, 2022), percent-encoded angle brackets in the Ref. 27 Bjerknes (Mar 1969) DOI URL, and added exact CrossRef publication dates for Refs. 11 and 14 (retaining prior Nov 21, 2026 DOI upgrades).

References & Primary Sources

* [1] Max Planck Institute for Meteorology (Feb 20, 2026). New Generation of Climate Models Sheds First Light on Long-Standing Pacific Puzzle. -- https://mpimet.mpg.de/en/communication/detail-view-news-homepage/new-generation-of-climate-models-sheds-first-light-on-long-s

* [2] Kang, S. M., Putrasahan, D. A., Brizuela, N. G., Haak, H., et al. (Feb 19, 2026). Km-scale coupled simulation and model-observation SST trend discrepancy. Proceedings of the National Academy of Sciences. -- https://doi.org/10.1073/pnas.2522161123

* [3] Armour, K. C., Proistosescu, C., Dong, Y., Hahn, L. C., et al. (Mar 11, 2024). Sea-surface temperature pattern effects have slowed global warming and biased warming-based constraints on climate sensitivity. Proceedings of the National Academy of Sciences. -- https://doi.org/10.1073/pnas.2312093121

* [4] Seager, R., Cane, M., Henderson, N., Lee, D., et al. (Jun 24, 2019). Strengthening tropical Pacific zonal sea surface temperature gradient consistent with rising greenhouse gases. Nature Climate Change. -- https://doi.org/10.1038/s41558-019-0505-x

* [5] Kim, H., Kang, S. M., Kay, J. E., & Xie, S.-P. (Aug 15, 2022). Subtropical clouds key to Southern Ocean teleconnections to the tropical Pacific. Proceedings of the National Academy of Sciences, 119(34), e2200514119. -- https://doi.org/10.1073/pnas.2200514119

* [6] Dong, L., Wang, Z., Wu, L., Song, F., et al. (Dec 29, 2025). Emergence of the enhanced equatorial Atlantic warming as a fingerprint of global warming. Nature Communications. -- https://doi.org/10.1038/s41467-025-68015-6

* [7] Hartmann, D. L. (Aug 22, 2022). The Antarctic ozone hole and the pattern effect on climate sensitivity. Proceedings of the National Academy of Sciences. -- https://doi.org/10.1073/pnas.2207889119

* [8] McGregor, S., Ong, E., & Planton, Y. (Feb 4, 2026). Understanding CMIP6 Multi-Model Ensemble Projected Pacific Warming Pattern Variability. Geophysical Research Letters. -- https://doi.org/10.1029/2025gl118815

* [9] Watanabe, M., Iwakiri, T., Dong, Y., Kang, S. M., et al. (Nov 29, 2023). Two Competing Drivers of the Recent Walker Circulation Trend. Geophysical Research Letters. -- https://doi.org/10.1029/2023gl105332

* [10] Kang, S. M., Yu, Y., Deser, C., Zhang, X., et al. (Jul 17, 2023). Global impacts of recent Southern Ocean cooling. Proceedings of the National Academy of Sciences. -- https://doi.org/10.1073/pnas.2300881120

* [11] M ller, W. A., Lorenz, S., Pham, T. V., et al. (Dec 3, 2025). The ICON-based Earth System Model for climate predictions and projections (ICON XPP v1.0). Geoscientific Model Development, 18, 9385-9405. -- https://doi.org/10.5194/gmd-18-9385-2025

* [12] Dong, Y., Armour, K. C., & Battisti, D. S. (Oct 1, 2022). Two-Way Teleconnections between the Southern Ocean and the Tropical Pacific via a Dynamic Feedback. Journal of Climate, 35(19), 6267-6282. -- https://doi.org/10.1175/JCLI-D-22-0080.1

* [13] Klocke, D., Frauen, C., Engels, J. F., Hoefler, T., Stevens, B., et al. (Nov 3, 2025). Computing the Full Earth System at 1 km Resolution (arXiv:2511.02021). -- https://doi.org/10.48550/arXiv.2511.02021

* [14] M ller, W. A., Fr h, B., Korn, P., Potthast, R., Baehr, J., & Bettems, J. (Jun 2025). ICON: Toward Vertically Integrated Model Configurations for Numerical Weather Prediction, Climate Predictions, and Projections. Bulletin of the American Meteorological Society, 106(6), E1017-E1031. -- https://doi.org/10.1175/BAMS-D-24-0042.1

* [15] ECMWF DestinE (May 6, 2024). The fast development of DestinE's Climate Change Adaptation Digital Twin. -- https://destine.ecmwf.int/news/the-fast-development-of-destines-climate-change-adaptation-digital-twin/

* [16] DKRZ (2025). nextGEMS: 30-year scenario simulations with IFS-FESOM and ICON at km-scale. -- https://luv.dkrz.de/media/2025/report_1235.pdf

* [17] easy.gems documentation (2021-2025). nextGEMS simulations overview. -- https://easy.gems.dkrz.de/DYAMOND/NextGEMS/simulation_overview.html

* [18] DKRZ (2025). Report on the joint project "WarmWorld" bk1341. -- https://luv.dkrz.de/media/2025/report_1341.pdf

* [19] Max Planck Institute for Meteorology (Nov 9, 2026). WarmWorld. -- https://mpimet.mpg.de/en/research/modeling/warmworld

* [20] WarmWorld Consortium (2025). WarmWorld: Exascale Earth System Models to Anticipate Changes in a Warmer World. -- https://www.warmworld.de/

* [21] DKRZ (2026). Report on the joint project "WarmWorld" bk1341. -- https://luv.dkrz.de/media/2026/report_1341.pdf

* [22] DKRZ (Dec 3, 2025). Faster Module Meeting: 3 years of progress and the road ahead. -- https://www.dkrz.de/en/communication/news-archive/faster-module-meeting2025

* [23] DKRZ (2026). Achievements in the previous allocation period. -- https://luv.dkrz.de/media/2026/report_1344.pdf

* [24] McPhaden, M. J., Zebiak, S. E., & Glantz, M. H. (Dec 15, 2006). ENSO as an Integrating Concept in Earth Science. Science, 314(5806), 1740-1745. -- https://doi.org/10.1126/science.1132588

* [25] Tang, T., Luo, J.-J., Peng, K., Qi, L., et al. (Apr 6, 2021). Over-projected Pacific warming and extreme El Ni o frequency due to CMIP5 common biases. National Science Review. -- https://doi.org/10.1093/nsr/nwab056

* [26] Jiang, F., Seager, R., & Cane, M. A. (Sep 27, 2024). A climate change signal in the tropical Pacific emerges from decadal variability. Nature Communications, 15, 8451. -- https://doi.org/10.1038/s41467-024-52731-6

* [27] Bjerknes, J. (Mar 1969). Atmospheric Teleconnections from the Equatorial Pacific. Monthly Weather Review, 97(3), 163-172. -- https://doi.org/10.1175/1520-0493(1969)097%3C0163:ATFTEP%3E2.3.CO;2

* [28] Falasca, F., Basinski-Ferris, A., Zanna, L., Zhao, M., et al. (Jul 15, 2025). A Fluctuation-Dissipation Theorem Perspective on Radiative Responses to Temperature Perturbations. Journal of Climate. -- https://doi.org/10.1175/jcli-d-24-0479.1