Climate & Earth Systems | Jun 5, 2026 | 12 min read | By Bryan White

Decoding the New World Screwworm: From Life Cycle to Eradication

Biology * Climate Change * Public Health

Discovery, Economic Burden, and Resurgence of the New World Screwworm

The New World screwworm, Cochliomyia hominivorax (Coquerel, 1858), is an obligate parasitic blowfly whose larvae feed exclusively on the living tissue of warm-blooded vertebrates, causing a rapidly progressive condition known as obligatory traumatic myiasis.[1] First described in 1858 by French naval surgeon Charles Coquerel following fatal nasal and sinus infestations among prisoners on Devil's Island in French Guiana, the species received the Latin epithet hominivorax ("man-eating").[2][4]

Historically distributed from the southern United States through Central America, the Caribbean, and South America to northern Argentina, the parasite inflicted $50 million to $100 million in annual mid-twentieth-century U.S. livestock losses.[2][6] Contemporary economic assessments indicate that widespread re-establishment in the United States could inflict billions of dollars in annual cattle, livestock, and wildlife losses.[6][7] Following the pioneering development of the Sterile Insect Technique (SIT) in the 1950s, the United States eradicated the pest by 1966, and multi-national campaigns progressively cleared Mexico and Central America, establishing a permanent biological barrier in Panama's Darien Gap by 2006.[2][7] However, beginning in 2023, a breach of this quarantine line triggered a rapid northward resurgence across Central America and Mexico, culminating in confirmed detections in South Texas in June 2026.[8] This renewed epidemiological threat underscores the need for a rigorous synthesis of the organism's genomics, morphology, thermal biology, pathology, and area-wide eradication logistics.

Taxonomy, Haplotype-Resolved Genomics, and Diagnostic Morphology

Classified within Diptera: Calliphoridae (subfamily Chrysomyinae), the genus Cochliomyia encompasses four American species -- C. hominivorax, C. macellaria, C. aldrichi, and C. minima -- forming a monophyletic sister clade to Compsomyiops.[1][10] Unlike its facultatively necrophagous congener C. macellaria (the secondary screwworm), C. hominivorax is obligately dependent on viable vertebrate tissue for larval development.[2]

Genetically, C. hominivorax possesses a diploid complement of 2n = 12 chromosomes, comprising five metacentric autosomal pairs and heteromorphic XY sex chromosomes in which a Y-linked male-determining factor governs sex regulation.[11] Recent trio-binning long-read sequencing and chromatin conformation scaffolding produced a haplotype-resolved 455.56 Mb haploid genome assembly (correcting earlier heterozygosity-inflated 534 Mb estimates), resolving a 6.96 Mb X chromosome, a 0.647 Mb Y contig, and over 98 percent conserved single-copy ortholog completeness.[12]

Population genetics surveys across South America reveal an average of 6.9 alleles per locus and high heterozygosity at flight-metabolism loci such as -glycerophosphate dehydrogenase, though isolated southern cohorts in Uruguay exhibit localized population bottleneck signatures.[13]

Rapid morphological differentiation between C. hominivorax and C. macellaria is essential for regulatory surveillance.[1] Adults measure 8-10 mm with a metallic blue-green sheen, reddish-orange compound eyes (holoptic in males for aerial mate interception, dichoptic in females), and three dark longitudinal thoracic stripes (vittae).[10] Mature third-instar larvae (15-17 mm) bear backward-facing cuticular spines resembling wood-screw threads and are definitively diagnosed by darkly pigmented dorsal tracheal trunks extending anteriorly across the ninth and tenth body segments.[1][14]

| Morphological/Ecological Trait | | Cochliomyia hominivorax (Primary Screwworm) | | Cochliomyia macellaria (Secondary Screwworm) |

| Larval Dietary Requirement | | Obligate parasite (consumes living, viable tissue) | | Facultative necrophage (consumes dead/decaying tissue) |

| Thoracic Stripes (Adult) | | Three distinct dark longitudinal vittae present | | Uniformly colored, stripes absent or highly indistinct |

| Basicosta Color (Adult Female) | | Dark brown to black | | Yellowish |

| Fronto-orbital Setulae (Adult) | | Darkly pigmented setulae | | Pale setulae |

| Larval Tracheal Trunks | | Darkly pigmented, faintly visible through body wall | | Clear or translucent, not pigmented |

| Larval Posterior Spiracles | | Incomplete peritreme, indistinct button | | Complete or distinct peritreme structures |

Table 1: Key morphological and ecological distinctions utilized in the laboratory identification of C. hominivorax and C. macellaria.[1][10]

Thermal Developmental Thresholds, Chemical Ecology, and Traumatic Myiasis

Lacking physiological diapause, C. hominivorax requires continuous year-round reproduction governed by ambient and soil temperatures between a lower developmental threshold of 14.5 deg C and an upper lethal limit near 43.5 deg C.[4] Egg-to-adult development completes in roughly 18 days at 29 deg C, 24 days at 22 deg C, and up to three months near minimum thermal thresholds.[4][15] Gravid females orient toward wound volatiles -- including ammonia and serous degradation products -- depositing tightly organized clutches of 200-500 eggs (up to 3,000 lifetime eggs) along wound margins, tick bites, or neonatal umbilical stumps.[2][10] Hatching within 12-24 hours, gregarious larvae tear living flesh with non-retractile oral hooks while secreting proteolytic enzymes, lipases, and ammonia that liquefy cellular proteins and alkalinize the wound microenvironment over 5-7 days.[10][15] Exiting third-instar larvae burrow into topsoil to pupate; soil temperatures below 7.7 deg C (46 deg F) for consecutive days or hard frosts are universally lethal to pupae, restricting permanent overwintering reservoirs to frost-free subtropics.[4][6]

Crucially for control, females are monandrous (mating once and storing sperm in spermathecae), whereas males are polygamous.[2][7] Tarsal mate recognition relies on contact pheromones composed of cuticular hydrocarbon (CHC) lipid profiles.[10][17] Because prolonged laboratory colonization can alter female CHC signatures and induce asymmetric mating discrimination by wild males, mixed-sex releases require substantially higher overflooding ratios than male-only releases.[17]

Unlike the facultative common green bottle fly (Lucilia sericata) employed in clinical maggot debridement therapy to clear necrotic slough without harming viable cells, C. hominivorax burrows up to 15 cm into vascularized musculature, cartilage, and cranial sinuses, threatening livestock and public health with fatal septicemia or hemorrhage within 7-10 days if untreated.[1][3][4][18] Veterinary management pairs larval extraction with systemic macrocyclic lactones, oral isoxazolines, and topical organophosphates or pyrethroids.[16][19]

| Active Ingredient | | Brand Name | | Application Type | | Target Species | | Indication / Mechanism |

| Ivermectin | | Ivomec | | Injectable Solution | | Cattle | | Systemic macrocyclic lactone; prevents myiasis when administered within 24 hours of birth or castration. |

| Doramectin | | Dectomax | | Injectable Solution | | Cattle, Swine, Sheep | | Systemic prevention and treatment; provides persistent plasma concentrations to eliminate feeding larvae. |

| Lotilaner | | Credelio | | Oral Chewable | | Dogs, Cats | | Systemic isoxazoline; rapidly enters bloodstream inducing 100% larval mortality within 24 hours. |

| Afoxolaner | | NexGard | | Oral Chewable | | Dogs | | Systemic isoxazoline; highly effective treatment of established myiasis. |

| Cypermethrin | | F10 Antiseptic Spray/Ointment | | Topical | | Livestock, Horses, Birds | | Synthetic pyrethroid combined with antiseptics for immediate contact mortality and wound disinfection. |

| Coumaphos/Propoxur | | Negasunt Powder | | Topical | | Livestock, Equids | | Organophosphate powder applied directly to the wound bed to kill larvae and prevent reinfestation. |

Table 2: Summary of pharmacological interventions and recent Emergency Use Authorizations for C. hominivorax myiasis.[16][19]

Area-Wide Control via the Sterile Insect Technique (SIT)

Conceived in 1937 by USDA entomologists Edward F. Knipling and Raymond C. Bushland and validated on Cura ao in 1954 (at 155 sterile males/km /week) as well as during the 1988-1991 Libyan eradication campaign (40 million sterile flies/week), SIT exploits female monandry by overflooding wild populations with irradiated sterile males.[5][7][20] Industrial rearing at facilities such as the COPEG plant in Pacora, Panama, evolved from whole-meat diets to cellulose- or polyacrylate-gelled formulations incorporating an 80:20 ratio of spray-dried bovine red blood cells (hemoglobin) and plasma, alongside cost-resilient chicken viscera/feather meal or soy alternatives.[21][22]

| Ingredient | | Standard Production Diet (%) | | Chicken By-Product Diet (%) | | Soy Alternative Diet (%) |

| Spray-dried RBCs | | 3.60 | | 3.60 | | 3.60 |

| Spray-dried Plasma | | 0.90 | | 2.40 | | 0.90 |

| Egg Powder | | 5.00 | | 0.00 | | 5.00 |

| Milk Powder | | 4.50 | | 0.00 | | 0.00 |

| Soy Powder | | 0.00 | | 0.00 | | 4.50 |

| Chicken By-products | | 0.00 | | 15.00 | | 0.00 |

| Cellulose Fiber | | 5.50 | | 5.00 | | 5.50 |

| Water | | 80.44 | | 73.94 | | 80.47 |

Table 3: Chemical formulations of artificial larval diets (% total mass) utilized in industrial screwworm mass rearing.[21][22]

Pupae are irradiated at 5.5-5.7 days of development -- after somatic organogenesis is largely complete -- using Cobalt-60 gamma or high-energy X-ray sources calibrated to 110 Gray (Gy), inducing >99% dominant lethal germline sterility without degrading emergence, flight propensity, or mating latency.[20][23][24] Modern dispersal replaces cardboard pupal boxes with automated chilled-adult release systems (maintaining emerged adults at 4-6 deg C) deployed via fixed-wing aircraft or drones.[16][25]

| System Component | | Specifications & Details |

| Operating Temperature | | 4 to 6 degrees Celsius (induces and maintains immobilization) |

| BSI Spreader Output | | ~60 flies per minute at 0.6 rpm setting |

| Birdview Cassette (Ver A.1) | | 20 cells, 464 cubic cm volume, 11.5 minute controlled release runtime |

| Birdview Cassette (Ver B) | | 40 cells, 774 cubic cm volume, Wi-Fi enabled ESP8266 microcontroller |

| Biological Impact | | Chilling and mechanical release process shows no significant negative impact on male flight propensity, mating competitiveness, or insemination rates.[25] |

Table 4: Technical specifications of automated chilled-adult SIT dispersal systems.[16][25]

Epidemiology of the 2023-2026 Resurgence and the Texas Quarantine

Driven by undocumented cattle movement across borders, surveillance gaps, and favorable warm weather anomalies, C. hominivorax breached the Darien Gap barrier in mid-2023.[4][6] Adult dispersal (up to 200 km per lifetime) and livestock transport propelled outbreaks across Costa Rica, Nicaragua, Honduras (where 76 human cases and fatalities were recorded in early 2026), El Salvador, Belize, and Guatemala, reaching Chiapas, Mexico in late 2024 and Coahuila by May 2026.[2][8][26]

On June 3, 2026, USDA APHIS and the Texas Animal Health Commission (TAHC) confirmed an autochthonous infestation in a three-week-old calf in Zavala County, Texas (~50 miles from the Mexican border), followed shortly by a secondary local detection.[8][27][28] Authorities immediately established "Infested Zone 01" -- a 20-kilometer quarantine perimeter across parts of Zavala and Uvalde counties prohibiting warm-blooded animal movement without veterinary inspection and prophylactic treatment -- while deploying Swormlure-4 surveillance traps and dispersing millions of chilled sterile flies per week over South Texas.[9][27][28]

Integrated Biosecurity Across International Borders

The 2023-2026 resurgence of Cochliomyia hominivorax from Panama into Texas underscores the speed at which an obligate parasite can exploit lapses in transboundary biosecurity. Re-establishing continental eradication requires integrating strict livestock movement quarantines, rapid veterinary administration of systemic isoxazolines and macrocyclic lactones, haplotype-informed colony management, and high-output 110 Gy chilled-adult SIT releases.[9][12][23]

Updates... (Nov 21, 2026): Updated U.S. outbreak status through November 18, 2026 -- noting confirmed animal detections reaching the high 100s across the American Southwest (spanning Texas, New Mexico, and Arizona), continued low general-public human risk with no locally acquired U.S. human cases reported by the CDC, and USDA's November 10 hiring drive (228 positions) and November 17 structural topping-off ceremony at Moore Air Base in Edinburg, Texas, to build domestic sterile-fly production capacity[8][9][28]; replaced Wikipedia and inaccessible/timeout URLs with verified primary literature (Wyss 2000, Welch 2016 CABI Compendium, Knipling 1955, and Texas Farm Bureau/TAHC reporting); corrected author/title metadata for Batista et al. (2009), e Souza et al. (2025), Thomas et al. (2018), and Mirieri et al. (2020); and standardized publication dates across all 28 references.

References & Primary Sources

* [1] Centers for Disease Control and Prevention (CDC) DPDx (Sep 16, 2025). New World Screwworm Myiasis. -- https://www.cdc.gov/dpdx/newworldscrewwormmyiasis/index.html

* [2] Wyss, J. H. (Dec 2000; online Jan 25, 2006). Screwworm Eradication in the Americas. Annals of the New York Academy of Sciences, 916(1), 186-193. -- https://doi.org/10.1111/j.1749-6632.2000.tb05289.x

* [3] Barros, G. P. de, & Bricarello, P. A. (2020). Myiasis by Cochliomyia hominivorax (Coquerel, 1858): A Neglected Zoonosis in Brazil. Open Journal of Veterinary Medicine, 10(6), 80-91. -- https://doi.org/10.4236/ojvm.2020.106007

* [4] Public Health Ontario (2026). Re-emergence of the New World Screwworm. -- https://www.publichealthontario.ca/-/media/Documents/N/26/new-world-screwworm-focus.pdf

* [5] International Atomic Energy Agency (IAEA) (1992). The New World Screwworm Eradication Programme: North Africa 1988-1992. -- https://www.iaea.org/sites/default/files/21/07/nafa-ipc-technical-report-screwworm-eradication-africa-1992.pdf

* [6] University of Arizona Cooperative Extension (Oct 2025). Preparing Livestock Producers for New World Screwworm Reemergence in the United States. --

https://extension.arizona.edu/publication/preparing-livestock-producers-new-world-screwworm-reemergence-united-states

* [7] M3 AgTech (Jun 8, 2025). Back to the Frontlines: Renewed Battle Against the New World Screwworm. -- https://m3agtech.com/2025/06/08/back-to-the-frontlines-renewed-battle-against-the-new-world-screwworm/

* [8] Centers for Disease Control and Prevention (CDC) (Nov 18, 2026). New World Screwworm Outbreak Situation Summary. -- https://www.cdc.gov/new-world-screwworm/situation-summary/index.html

* [9] Texas Animal Health Commission (TAHC) (2026). New World Screwworm Emergency Information and Movement Restrictions. -- https://www.tahc.texas.gov/emergency/nws.html

* [10] Marshall, S. A., Whitworth, T., & Roscoe, L. (2011). Blow flies of North America: Keys to the subfamilies and genera of Calliphoridae. Canadian Journal of Arthropod Identification. -- https://cjai.biologicalsurvey.ca/wp-content/uploads/2021/01/jwm_39.pdf

* [11] Batista, M. R. D., Ananina, G., Azeredo-Espin, A. M. L., & Klaczko, L. B. (Mar 24, 2009; print Jun 2009). Photographic map of the polytene chromosomes of Cochliomyia hominivorax. Medical and Veterinary Entomology, 23(s1), 92-97. -- https://doi.org/10.1111/j.1365-2915.2008.00775.x

* [12] Hickner, P. V., Sim, S. B., Luecke, D., Bailey, E., Saelao, P., & Yusseff-Vanegas, S. Z. (Feb 28, 2026; print May 6, 2026). Haplotype-resolved genome assemblies for the New World screwworm, Cochliomyia hominivorax (Diptera: Calliphoridae), using the trio binning approach. G3: Genes, Genomes, Genetics, 16(5), jkag053. -- https://doi.org/10.1093/g3journal/jkag053

* [13] e Souza, K. da S., de Paula, L. C. B., de Azeredo-Espin, A. M. L., & Torres, T. T. (Jan 20, 2025). Demographic and historical processes influencing Cochliomyia hominivorax (Diptera: Calliphoridae) population structure across South America. Parasites & Vectors, 18(1), 18. -- https://doi.org/10.1186/s13071-024-06622-w

* [14] Centers for Disease Control and Prevention (CDC) (2024). Lab Identification of New World Screwworm Bench Aid. -- https://www.cdc.gov/dpdx/resources/pdf/benchaids/NWS-Bench-Aid-2024-508.pdf

* [15] Welch, J. B. (Jan 27, 2016). Cochliomyia hominivorax infestation. CABI Compendium. -- https://doi.org/10.1079/cabicompendium.121349

* [16] Maiga, H., Zambrana, A. M., Somda, N. S. B., Mamai, W., Wallner, T., & Kotla, S. S. (Nov 27, 2025). A 3D-printed adult release system compatible with a drone for aerial deployment of Aedes aegypti and Glossina palpalis gambiensis. Parasites & Vectors, 19(1), 2. -- https://doi.org/10.1186/s13071-025-07146-7

* [17] Vreysen, M. J. B., Barclay, H. J., & Hendrichs, J. (May 1, 2006). Modeling of Preferential Mating in Areawide Control Programs That Integrate the Release of Strains of Sterile Males Only or Both Sexes. Annals of the Entomological Society of America, 99(3), 607-616. -- https://doi.org/10.1603/0013-8746(2006)99[607:mopmia]2.0.co;2

* [18] Sherman, R. A., Hall, M. J. R., & Thomas, S. (Jan 2000). Medicinal Maggots: An Ancient Remedy for Some Contemporary Afflictions. Annual Review of Entomology,

45(1), 55-81. -- https://doi.org/10.1146/annurev.ento.45.1.55

* [19] U.S. Food and Drug Administration (FDA) (Nov 17, 2026). New World Screwworm: Information for Veterinarians. -- https://www.fda.gov/animal-veterinary/safety-health/new-world-screwworm-information-veterinarians

* [20] Knipling, E. F. (Aug 1, 1955). Possibilities of Insect Control or Eradication Through the Use of Sexually Sterile Males. Journal of Economic Entomology, 48(4), 459-462. -- https://doi.org/10.1093/jee/48.4.459

* [21] Hickner, P. V., Sagel, A., Quintero, G., Vasquez, M., Tietjen, M., & Lohmeyer, K. H. (Nov 25, 2023; print Feb 12, 2024). An alternative chicken-based diet for mass-rearing screwworm flies. Journal of Economic Entomology, 117(1), 348-357. -- https://doi.org/10.1093/jee/toad219

* [22] Thomas, J. K., Fadul, G. J., Keller, G. P., & Chaudhury, M. F. (May 1, 2018). The Use of Dried Bovine Hemoglobin and Plasma for Mass Rearing New World Screwworm. Journal of Insect Science, 18(3). -- https://doi.org/10.1093/jisesa/iey052

* [23] Kabor , B. A., Nawaj, A., Maiga, H., Soukia, O., Pagabeleguem, S., & Ou draogo/Sanon, M. S. G. (Oct 17, 2023). X-rays are as effective as gamma-rays for the sterilization of Glossina palpalis gambiensis Vanderplank, 1911 (Diptera: Glossinidae) for use in the sterile insect technique. Scientific Reports, 13(1), 17633. -- https://doi.org/10.1038/s41598-023-44479-8

* [24] LaChance, L. E., & Bruns, S. B. (Feb 1963). Oogenesis and Radiosensitivity in Cochliomyia hominivorax (Diptera: Calliphoridae). The Biological Bulletin, 124(1), 65-83. -- https://doi.org/10.2307/1539569

* [25] Mirieri, C. K., Mutika, G. N., Bruno, J., Seck, M. T., Sall, B., & Parker, A. G. (Sep 28, 2020). A new automated chilled adult release system for the aerial distribution of sterile male tsetse flies. PLOS ONE, 15(9), e0232306. -- https://doi.org/10.1371/journal.pone.0232306

* [26] BeaconBio (2026). First New World screwworm myiasis death in 2026 in Honduras, among 76 human cases. -- https://beaconbio.org/en/report/?reportid=8d7d13c2-a451-4fab-a330-481fdbebc9db

* [27] Western Ag Network (Jun 3, 2026). USDA Confirms New World Screwworm Detection in Texas Calf. -- https://westernagnetwork.com/usda-confirms-new-world-screwworm-detection-in-texas-calf

* [28] Texas Farm Bureau (Nov 18, 2026). New World screwworm cases increase across Texas. -- https://texasfarmbureau.org/new-world-screwworm-cases-increase-across-texas/