Space & Astronomy | Nov 21, 2026 | 5 min read | By Bryan White

ESA's Hera Spacecraft Arrives at Didymos to Investigate NASA's Asteroid Impact Site

Space & Astronomy * Astrophysics

Four years after NASA's Double Asteroid Redirection Test (DART) deliberately crashed into the asteroid moonlet Dimorphos, the European Space Agency's (ESA) Hera spacecraft has arrived at the Didymos-Dimorphos binary system to conduct a close-range planetary defense investigation. Operating from 25 to 60 kilometers out during its early characterization phase, Hera has already returned monochromatic Asteroid Framing Camera (AFC) views of the previously unmapped hemisphere of the primary body, Didymos, revealing a boulder-strewn, heavily cratered landscape and a pronounced equatorial ridge. On November 25, 2026, ESA plans to unveil the first detailed post-impact close-up observations of Dimorphos itself, answering long-standing questions about whether the 2022 hypervelocity collision carved a discrete crater or globally reshaped the rubble-pile moonlet.

From Proof of Concept to Forensic Planetary Defense

On September 26, 2022, the 570-kilogram DART impactor struck the 151-meter moonlet Dimorphos at roughly 6.1 kilometers per second. Ground-based lightcurve and radar observations quickly confirmed that the collision shortened Dimorphos's orbital period around the 780-meter primary asteroid Didymos by approximately 33 minutes, dropping it from 11 hours and 55 minutes to roughly 11 hours and 22 minutes. Subsequent orbital tracking also revealed that the massive plume of ejected regolith subtly altered the binary system's heliocentric orbital period around the Sun by 0.15 seconds, corresponding to an orbital speed shift of about 11.7 micrometers per second.

While DART proved that a kinetic impactor can measurably deflect a hazardous near-Earth asteroid , Earth-based telescopes and the brief flyby of the Italian Space Agency's LICIACube could not resolve Dimorphos's mass, internal porosity, or post-impact topography. Without those physical parameters, planetary defense models cannot reliably scale kinetic impactor designs to larger or structurally different asteroids. Hera, launched in October 2024, was built specifically to close that observational gap by performing a multi-month "crash site investigation" of the binary system.

| Parameter / Metric | | Primary Body: Didymos | | Moonlet: Dimorphos |

| Approximate Diameter | | ~780 meters | | ~151 meters (pre-impact) |

| Pre-Impact Mutual Orbit | | Binary barycenter host | | 11 hours 55 minutes |

| Post-DART Orbital Shift | | System Sun orbit shifted by 0.15 s | | Shortened by ~33 minutes |

| Along-Track Velocity Change | | 11.7 um/s (heliocentric system) | | 2.70 +/- 0.10 mm/s (mutual orbit) |

| Momentum Enhancement (beta) | | N/A (target host) | | 2.2 to 4.9 (baseline ~3.61) |

Why Ejecta Recoil Multiplied DART's Impact Punch

The central physical mystery driving Hera's arrival is the magnitude of the momentum enhancement factor, denoted by the Greek letter beta. In a purely inelastic collision where an impactor embeds itself into a solid target with zero escaping debris, beta equals 1.0, meaning the asteroid receives only the direct linear momentum of the spacecraft. In a hypervelocity impact on a low-gravity rubble pile, however, shock waves excavate thousands of tons of boulders and fine dust backward into space, acting like a transient rocket exhaust.

Peer-reviewed analysis by the DART Investigation Team published in Nature found an instantaneous along-track orbital velocity reduction of 2.70 +/- 0.10 millimeters per second. Depending on Dimorphos's assumed bulk density (ranging between 1,500 and 3,300 kilograms per cubic meter), those orbital measurements yield a beta factor between 2.2 and 4.9, with equal-density models (2,400 kilograms per cubic meter) placing beta at 3.61. That means escaping ejecta delivered more than triple the momentum of the spacecraft itself. Yet density and mass remain degenerate in telescope-only observations; Hera must directly measure Dimorphos's gravitational pull on the spacecraft to pin down beta with high precision.

First Close-Up Views of Didymos and Upcoming CubeSat Sounding

Approaching from an illumination geometry opposite to DART's 2022 trajectory, Hera resolved Didymos and Dimorphos as distinct bodies on October 27, 2026, before releasing its first close-up AFC images from 60 kilometers away on November 5, 2026. Although Dimorphos sat largely in shadow in those initial frames, the imagery exposed Didymos's far hemisphere in sharp detail, showcasing a spinning-top profile forged by the Yarkovsky-O'Keefe-Radzievskii-Paddack (YORP) thermal torque effect, steep boulder fields, and impact craters that raise new questions about how DART's 2022 debris interacted with the primary body. Because Didymos rotates every 2.26 hours -- close to the structural spin limit for a cohesionless rubble pile -- material near its equatorial ridge sits in a low-effective-gravity regime where even modest ejecta impacts can mobilize surface regolith.

Over the coming months, Hera will combine its Asteroid Framing Cameras, the HyperScout H hyperspectral imager, a thermal infrared imager (TIRI), and a planetary laser altimeter (PALT) to construct global topographic and compositional maps. Simultaneously, precision radio-science tracking of Hera's trajectory perturbations during close flybys will directly weigh Dimorphos, breaking the long-standing degeneracy between bulk density and momentum enhancement. In early 2027, the spacecraft will deploy two 6U CubeSats -- Juventas and Milani -- to fly within a few kilometers of the surface. Juventas carries a low-frequency monostatic radar (JuRA) designed to peer inside Dimorphos for the first time, testing whether the moonlet is a loosely bound rubble pile held together primarily by self-gravity and weak van der Waals cohesive forces, while Milani maps S-type silicate mineralogy and sub-millimeter dust dynamics.

Transforming an Experiment Into a Repeatable Defense Standard

Hydrocode impact simulations have long struggled to predict whether a 6-kilometer-per-second strike on a weakly bound rubble pile produces a classic bowl-shaped crater or triggers global structural deformation that reshapes the entire body. By comparing pre-impact DART imagery with Hera's upcoming November 25, 2026 high-resolution Dimorphos views and 2027 gravity-field measurements, planetary scientists will directly calibrate those numerical models against real-world ground truth.

Establishing validated bounds on rubble-pile cohesion, porosity, and ejecta momentum amplification ensures that future deflection missions will not have to guess how a threatening near-Earth object will respond to kinetic redirection. As Hera transitions from initial characterization into close-proximity mapping and CubeSat deployment, the Didymos-Dimorphos system is serving as humanity's first fully instrumented laboratory for planetary defense mechanics and asteroid geophysics.

References & Primary Sources

* [1] European Space Agency (ESA) -- Hera Mission Official Portal: Planetary Defence Asteroid Investigation -- https://www.heramission.space/

* [2] NASA Science -- Double Asteroid Redirection Test (DART) Mission Overview and Orbital Results -- https://science.nasa.gov/mission/dart/

* [3] Cheng, A. F., et al. (2023). Momentum transfer from the DART mission kinetic impact on asteroid Dimorphos. Nature, 616, 457-460. -- https://doi.org/10.1038/s41586-023-05878-z