Space & Astronomy | Nov 11, 2026 | 5 min read | By Bryan White

ESA's Hera Spacecraft Reaches Dimorphos to Investigate Planetary Defense Impact Crater and Internal Rubble-Pile Architecture

Space & Astronomy * Orbital Mechanics * Planetary Science

More than four years after NASA's Double Asteroid Redirection Test (DART) spacecraft slammed into the near-Earth asteroid moonlet Dimorphos at 6.1 kilometers per second (13,600 mph), humanity's first dedicated forensic investigation of an intentional kinetic asteroid deflection has begun [1, 2] . The European Space Agency's (ESA) Hera spacecraft has successfully rendezvoused with the binary asteroid system (65803) Didymos , establishing an initial orbital station 25 to 30 kilometers (15 to 18 miles) away from the impacted moonlet [1, 5] . Over the coming weeks, Hera will close within 10 to 20 kilometers, deploying two specialized CubeSats -- Juventas and Milani -- to conduct low-frequency radar tomography, surface mineralogical mapping, and precise gravitational mass determinations [1, 5] . The mission aims to resolve a central question in planetary defense: whether the DART impact carved out a conventional bowl-shaped impact crater or globally deformed the internal structure of Dimorphos as a loosely bound rubble pile [1, 4] .

The DART Heritage: Validating Kinetic Impactor Physics

On September 26, 2022, NASA's 570-kilogram DART probe executed an autonomous optical terminal guidance maneuver, striking the 160-meter-diameter Dimorphos head-on [2, 3] . Ground-based radar networks and optical lightcurve monitoring subsequently confirmed that the collision reduced Dimorphos's orbital period around its 780-meter primary asteroid, Didymos, by 33.0 +/- 1.0 minutes (from 11 hours and 55 minutes down to 11 hours and 22 minutes) [2] .

This period reduction exceeded initial pre-impact expectations by more than a factor of three [2, 3] . In a purely inelastic collision -- where the kinetic impactor simply transfers its linear momentum (m p v p) and sticks to the target -- the theoretical period change would have been approximately 7 minutes [2, 3] . The additional 26 minutes of orbital deceleration was generated by the momentum enhancement factor (beta) , produced when millions of kilograms of pulverized rock and dust were blasted outward into space opposite the spacecraft's flight vector, acting as an impromptu rocket thruster [3, 4] . While terrestrial telescopes tracked the resulting 10,000-kilometer ejecta tail, measuring the precise mass of Dimorphos -- and thus pinning down the true value of beta -- requires close-proximity gravitational tracking by Hera [1, 3] .

Telemetry & Mission Parameters: Pre-Impact vs. Post-Impact System

Hera's multi-instrument payload -- including the Asteroid Framing Camera (AFC), the Thermal Infrared Imager (TIRI), and the Planetary Altimeter (PALT) -- will systematically compare the post-collision state of the binary system against pre-impact baseline models [1, 4, 5] .

| System Parameter | | Pre-Impact State (2022 Baseline) | | Post-DART State / Hera Target | | Planetary Defense Significance |

| Dimorphos Orbital Period | | 11 hours, 55 minutes (715 min) | | 11 hours, 22 minutes (682 min) | | Demonstrated reliable kinetic orbital deflection of an asteroid moonlet [2] . |

| Momentum Transfer Factor (beta) | | Theoretical range: 1.0-3.0 | | Estimated beta ~ 2.2-3.6 (Awaiting Hera) | | Quantifies the propulsive recoil efficiency of high-velocity ejecta [3, 4] . |

| Target Morphological State | | Oblate spheroid (rugby-ball profile) | | Triaxial reshape / flattened rubble pile | | Tests whether kinetic impacts produce localized craters or global deformation [4] . |

| Ejected Material Fraction | | 0% (intact pristine surface) | | ~1% of total asteroid mass ejected | | Dispersed an estimated 10 7 kg of rock into the interplanetary medium [3, 4] . |

| Internal Cohesion (τ c) | | Estimated <100 Pa (cohesionless) | | To be probed by Juventas radar | | Determines structural resistance to tidal disruption and kinetic deflection [1, 4] . |

Global Reshaping vs. Classical Crater Excavation

One of the most consequential findings from post-impact numerical shock-physics simulations -- published in Nature Astronomy by Raducan and colleagues -- is that Dimorphos likely suffered global structural reshaping rather than localized crater excavation [4] .

In solid rock bodies (such as basalt or iron- nickel meteorites), hypervelocity impacts are governed by the strength regime , where the yield strength of the rock confines shock damage to a hemispherical crater roughly 10 to 20 times the projectile's diameter [1, 4] . However, small near-Earth asteroids like Bennu, Ryugu, and Dimorphos are rubble piles : gravitational aggregations of boulders, pebbles, and fine dust held together by microscopic van der Waals forces and feeble self-gravity [1, 4] .

Because Dimorphos possesses an estimated cohesive strength of less than a few Pascals, the DART impact shock wave propagated across the entire interior of the body without encountering structural resistance [4] . Numerical models indicate that while roughly 1% of the asteroid's mass escaped into space as ejecta, approximately 8% of the moonlet's internal mass was redistributed , causing the entire body to settle into an altered triaxial ellipsoidal geometry [4] . Hera's close-range optical and thermal imaging will directly confirm whether the impact site is a distinct crater or an unrecognizable, resurfaced rubble field [1, 4] .

Sub-Spacecraft Exploration: Juventas and Milani CubeSats

To inspect Dimorphos without risking the primary mother ship, Hera carries two six-unit (6U) nanosatellites that will be released into independent orbits around the binary asteroid [1, 5] :

* Juventas: Equipped with the Juventas Radar (JuRa), a low-frequency monostatic

synthetic aperture radar, Juventas will perform humanity's first subsurface radar sounding of an asteroid interior [1, 5] . Operating at 60 MHz, JuRa can penetrate down to 100 meters, mapping internal void spaces, boulder packing fractions, and structural fault lines before attempting a gentle soft-landing on Dimorphos's surface [1, 5] .

* Milani: Named after celestial mechanician Andrea Milani, this CubeSat carries the ASPECT hyperspectral imager (operating from 500 to 1,500 nm) to map mineralogical alterations, space weathering , and surface volatile distribution across both Didymos and Dimorphos [1, 5] . Milani also carries the VISTA dust detector to monitor any lingering particulate clouds suspended in the binary orbital environment [1, 5] .

Calibrating the Kinetic Deflector Toolkit for Earth Protection

The ultimate objective of the joint NASA-ESA AIDA (Asteroid Impact and Deflection Assessment) collaboration is to turn kinetic deflection from an experimental proof-of-concept into a calibrated planetary defense technique [1, 3] . If astronomers discover a hazardous 100-to-300-meter asteroid on a collision course with Earth decades in advance, knowing the exact value of the momentum enhancement parameter beta is essential to calculate the required impactor mass and lead time needed to nudge the threat into a safe trajectory [1, 3] .

By measuring the precise mass of Dimorphos through radio Doppler tracking and mapping its internal cohesion, Hera will close the loop on the 2022 impact experiment [1, 2, 4] . The resulting datasets will provide planetary defense engineers with the ground-truth physical parameters required to safeguard Earth against future celestial impacts [1, 3, 5] .

References & Primary Sources

* [1] Michel, P., K ppers, M., Fitzsimmons, A. et al. (2025). The Hera Space Mission in the Context of Small Near-Earth Asteroid Missions in the Past, Present and Future. Space Science Reviews, 221, 70. -- https://doi.org/10.1007/s11214-025-01195-1

* [2] Thomas, C. A. et al. (2023). Orbital period change of Dimorphos due to the DART kinetic impact. Nature, 616(7957), 448-451. -- https://doi.org/10.1038/s41586-023-05805-2

* [3] Cheng, A. F. et al. (2023). Momentum transfer from the DART mission kinetic impact on asteroid Dimorphos. Nature, 616(7957), 457-460. -- https://doi.org/10.1038/s41586-023-05878-z

* [4] Raducan, S. D. et al. (2024). Physical properties of asteroid Dimorphos as derived from the DART impact. Nature Astronomy, 8, 445-455. -- https://doi.org/10.1038/s41550-024-02200-3

* [5] European Space Agency (2026). ESA's Hera Mission Begins Close Survey of Binary Asteroid Dimorphos. -- https://www.esa.int/Space_Safety/Hera