Climate & Earth Systems | Mar 10, 2026 | 9 min read | By Bryan White

Extreme Bifurcated Atmospheric Rivers Along the West Coast: Decoding the Predicted March 2026 Heatwave in California

Environment * Climate Change * Oceanography

The Continental Weather Dipole

During mid-March 2026, the North American West Coast experienced a pronounced latitudinal weather dipole characterized by an abrupt meridional split in temperature and precipitation.[1][9] Across the Pacific Northwest, a multi-day atmospheric river sequence delivered extreme coastal rainfall and multi-foot mountain snow accumulations.[2][9] Simultaneously, the southern branch of the dipole across California and the Southwest became locked beneath anomalous upper-level ridging, driving record-shattering early-spring heat, intense adiabatic drying, and episodic offshore Santa Ana winds.[1][22]

This synoptic divergence provides a revealing case study in large-scale meteorology and regional hydrology during the critical March transition -- the final window of Western snowpack accumulation before spring melt.[4] Both the precipitation deficits in California's Sierra Nevada and the extreme deluge across Washington and Oregon carried cascading consequences for agricultural water allocations and ecological restoration, notably across the Klamath River basin following the largest dam removal project in United States history, located directly along the boundary between these two regimes.[19][20]

Synoptic Climatology and Pacific Teleconnections

The mid-March 2026 dipole was driven by a high-amplitude, quasi-stationary Rossby wave configuration in the upper troposphere coinciding with a rapid transition in the El Ni o-Southern Oscillation (ENSO) -- specifically the decay of a cool-phase La Ni a.[5][8] Cooler-than-average equatorial Pacific sea surface temperatures during La Ni a typically displace the Pacific jet stream poleward, steering storm tracks into the Pacific Northwest while leaving the Southwest warm and dry; even as oceanic anomalies relaxed toward neutral in early 2026, the atmosphere exhibited a lagged wave response.[5][6][8] Simultaneously, Northern Hemisphere teleconnections shifted from the negative Arctic Oscillation blocking of January-February to a progressive positive Arctic Oscillation and negative Pacific-North American (PNA) pattern.[6]

Within this regime, a persistent high-pressure blocking ridge anchored over the Great Basin and offshore California deflected the westerly jet stream northward into a deep low-pressure trough off Washington and Oregon.[6][7][8] The resulting tight baroclinic gradient channeled deep tropical moisture into the Pacific Northwest while inducing large-scale subsidence across California.[9][11] As air descended through the ridge column, thermodynamic adiabatic compression warmed the air parcel according to the dry adiabatic lapse rate, evaporating cloud cover and maximizing solar insolation across California.[1][11]

The Northern Regime: Integrated Vapor Transport and Pacific Northwest Deluge

The northern half of the dipole is governed by narrow corridors of concentrated lower-tropospheric water vapor transport that account for up to 80% of West Coast flood damages ($1.1 billion annually).[9][10] Meteorologists classify these storms using the Ralph et al. (2019) scale (Categories 1-5), which pairs 3-hour mean maximum Integrated Vapor Transport (IVT, in kg m 1 s 1 across thresholds from 250 to 1,250 kg m 1 s 1) with event duration (upgrading or downgrading by one category for durations >48 hours or <24 hours).[12]

For the sequence beginning March 10, 2026, Center for Western Weather and Water Extremes (CW3E) West-WRF ensemble models indicated a >75% probability of Category 3 or stronger conditions across southwestern Washington and northwestern Oregon, with IVT remaining elevated for >72 hours.[2][3] At 45.5 deg N, 124 deg W (Tillamook County, Oregon), >90% of ensemble members projected at least Category 3 intensity and ~40% reached Category 4, supported by CW3E Atmospheric River Reconnaissance aircraft dropsonde campaigns.[2][3] While initial low freezing levels produced 2 to 4 feet of orographic snow above 3,000 feet in the Olympics and Cascades under winter storm warnings, a secondary "Pineapple Express" plume arriving around March 15 raised freezing levels sharply.[2][9][13] The resulting rain-on-snow regime released latent heat into fresh snowpack, driving rapid stream rises.[2][9]

| Geographic Region | | Forecasted Precipitation (7-Day Total) | | Forecasted Snow Accumulation | | Primary Associated Hazards |

| Olympic Mountains (WA) | | 7 to 15 inches | | Exceeding 48 inches | | Extreme avalanche danger, rapid rain-on-snow runoff |

| Washington Cascades | | 7 to 15 inches | | Exceeding 48 inches | | Blizzard conditions, mountain pass closures |

| Northern Oregon Coast Ranges | | 7 to 15 inches | | Minimal (Rain-dominated) | | Coastal riverine flooding, localized landslides |

| Willapa Hills / Lowlands (WA) | | 5 to 10 inches | | Minimal (Rain-dominated) | | Mainstem river flooding, mudslides |

The Southern Regime: Adiabatic Heating, Santa Ana Winds, and Sierra Snowpack Deficit

Beneath the descending branch of the Great Basin ridge, California and the Southwest experienced early-March forecast anomalies of 15 deg F to 25 deg F -- intensifying by mid-to-late March into heat-dome anomalies of 20 deg F to 30 deg F (11 deg C to 17 deg C) above mid-March climatology.[1][8][22] Baseline March averages in Sacramento (64.8 deg F), Yosemite Valley (56.7 deg F per NOAA NCEI 1991-2020 normals), and the Los Angeles Basin (68.0 deg F) gave way to forecast highs of 70-75 deg F, 79-82 deg F, and 85-90 deg F respectively before temperatures exceeded 100 deg F across portions of the Desert Southwest.[14][15][22] Concurrently, surface pressure gradients between the high-elevation Great Basin plateau and the Southern California coast favor episodic offshore Santa Ana winds through mountain passes, compressing adiabatically at ~5.5 deg F per 1,000 feet of descent to depress

relative humidity and elevate vapor pressure deficit and wildfire risk.[11][16]

| Location | | Historical March Average High | | Early-Event March 2026 Forecasted High | | Prevailing Synoptic & Mesoscale Conditions |

| Sacramento | | 64.8 deg F | | 70 to 75 deg F | | Clear skies, persistent subsidence warming |

| San Francisco | | 59.0 deg F | | 61 to 65 deg F | | Moderated coastal marine breeze, clear skies |

| Yosemite Valley | | 56.7 deg F | | 79 to 82 deg F | | Record-threatening heat, premature snowpack ablation |

| Los Angeles Basin | | 68.0 deg F | | 85 to 90 deg F | | Adiabatic compressional heating, offshore Santa Ana potential |

Hydrologically, severe March warmth threatened California's "frozen reservoir" -- the Sierra Nevada snowpack that supplies ~30% of annual state water and typically peaks near April 1.[4][17] Despite February storms that followed a five-week mid-winter drought, the California Department of Water Resources recorded just 28 inches of snow depth (11 inches snow water equivalent) at Phillips Station on February 27, 2026 -- roughly two-thirds of normal.[17][18] Although major reservoir carryover buffered liquid storage early in the season, premature March snowpack sublimation and melt -- compounded by low early-season Central Valley Project water allocations for contractors south of the Sacramento-San Joaquin Delta -- ultimately dropped statewide snowpack to 18%-20% of normal (with bare ground at Phillips Station) by April 1 and heightened agricultural reliance on groundwater pumping.[17][22][23]

Ecological Resilience Along the Dipole Boundary: The Klamath River Basin

Straddling the Oregon-California border directly along the inflection point of the March 2026 weather dipole lies the Klamath River basin, where the $500 million removal of four hydroelectric dams (J.C. Boyle, Copco 1, Copco 2, and Iron Gate) was completed in late 2024, reconnecting 257 miles of free-flowing river.[19][20] Prior to removal, stagnant warm impoundments fostered toxic cyanobacteria blooms, depressed dissolved oxygen, blocked historic spawning tributaries, and triggered the 2002 lower-river salmon die-off and 2006 commercial fishery closure.[19][20]

Post-dam monitoring using sonar, video weirs, eDNA, and telemetry recorded 7,700 anadromous fish (588 fish/day, ~96% Chinook salmon) passing the former Iron Gate Dam site between October and December 2024, reversing a decades-long trophic cascade across the ecosystem.[21] During spring heat events, free-flowing river connectivity allows cooler high-elevation storm runoff from the upper Klamath basin in southern Oregon to buffer downstream California reaches against thermal stress -- maintaining refugia and sediment transport corridors that would previously have been disrupted by warm reservoir impoundments.[2][20][21]

Conclusion: Managing Hydroclimate Divergence and Attribution Outcomes

The mid-March 2026 West Coast weather dipole demonstrated how decaying La Ni a teleconnections and amplified Great Basin ridging can bifurcate the Pacific storm track into simultaneous hydrometeorological extremes.[1][5] Subsequent post-event attribution by World Weather Attribution (March 20, 2026) and NOAA GOES West observations confirmed that the ensuing March heat dome shattered all-time monthly temperature records across California, Nevada, and Arizona -- reaching 20 deg F to 30 deg F (11 deg C to 17 deg C) above normal -- and was virtually impossible without human-induced climate change, which boosted event intensity by ~2.6 deg C and likelihood by roughly 800-fold relative to pre-industrial conditions (including +0.8 deg C and ~4-fold increased likelihood over the last decade alone).[1][22] Combined with an early late-February snowpack peak and rapid March ablation that left April 1 California snow water equivalent near 18%-20% of normal before ENSO transitioned later in 2026, the season underscored the urgency of heat-action planning, groundwater buffers, and free-flowing watershed restoration.[8][22][23]

Updates... (Nov 22, 2026): Updated tense and post-event outcomes through autumn 2026 to document the verified March 2026 Western North America heat dome (World Weather Attribution: ~2.6 deg C hotter and ~800 times more likely relative to pre-industrial baselines, including +0.8 deg C and ~4-fold over the past decade) and April 1 California snowpack collapse (~18%-20% of normal; bare ground at Phillips Station), aligned Yosemite Valley's historical March average high with NOAA NCEI 1991-2020 climate normals (56.7 deg F at station USC00049855), replaced undated/rolling and 404 travel/archive weather links with official NPS and NOAA NCEI 1991-2020 climate normals, corrected the NOAA NESDIS atmospheric river primer date (Dec 8, 2023) and USFWS Klamath report date (Feb 22, 2016), and standardized exact publication dates across Works Cited.

Works Cited

* NOAA NESDIS (2026, March 19). NOAA's GOES West Watches Record Heat Wave. https://www.nesdis.noaa.gov/news/noaas-goes-west-watches-record-heat-wave

* Center for Western Weather and Water Extremes (CW3E) (2026, March 9). CW3E AR Update: 9 March 2026 Outlook. https://cw3e.ucsd.edu/cw3e-ar-update-9-march-2026-outlook/

* Center for Western Weather and Water Extremes (CW3E) (2026, March 9). CW3E Atmospheric River Outlook: 9 March 2026 (PDF). https://cw3e.ucsd.edu/wp-content/uploads/2026/03/9Mar2026_Outlook/9Mar2026_Outlook.pdf

* California Department of Water Resources (2026, January 31). California Hydrology Update: Conditions as of January 31, 2026. https://cww.water.ca.gov/service/document/hydroreport

* Severe Weather Europe (2026, February 12). Spring 2026 Forecast: Why the Rapid Shift to El Ni o is Reshaping the United States and Canada Outlook. https://www.severe-weather.eu/long-range-2/spring-2026-forecast-la-nina-collapse-major-weather-shift-united-states-canada-fa/

* SnowBrains (2026, February 19). NOAA March 2026 Outlook: Could Miracle March Save

the Season?
https://snowbrains.com/noaa-march-2026-outlook-could-miracle-march-save-the-season/

* Palisades Tahoe (2026, March 8). Sunny March Pattern.
https://blog.palisadestahoe.com/weather/sunny-march-pattern/

* NOAA Climate Prediction Center (2026). ENSO: Recent Evolution, Current Status and Predictions.
https://www.cpc.ncep.noaa.gov/products/analysis_monitoring/lanina/enso_evolution-status-fcsts-web.pdf

* NOAA NESDIS (2023, December 8). Atmospheric Rivers Drench the Pacific Northwest.
https://www.nesdis.noaa.gov/news/atmospheric-rivers-drench-the-pacific-northwest

* U.S. Geological Survey (USGS) (2019). A scale to characterize the strength and impacts of atmospheric rivers. https://pubs.usgs.gov/publication/70202024

* UCLA Atmospheric and Oceanic Sciences (2026, March 5). Weather Synopsis - March 5, 2026. https://atmos.ucla.edu/weather-synopsis/weather-synopsis-march-5-2026/

* Ralph, F. Martin, Rutz, Jonathan J., Cordeira, Jason M., Dettinger, Michael, Anderson, Michael, Reynolds, David, Schick, Lawrence J., & Smallcomb, Chris (2019, February [online 2018, September 20]). A Scale to Characterize the Strength and Impacts of Atmospheric Rivers. Bulletin of the American Meteorological Society, 100(2), 269-289. https://doi.org/10.1175/BAMS-D-18-0023.1

* Ventusky (2026). Weather Warnings - North Cascades (48.97 deg N, 121.32 deg W).
https://www.ventusky.com/48.97;-121.32/warnings

* National Park Service (NPS) / NOAA 1991-2020 Climate Normals (2021). Yosemite National Park: Weather (Yosemite Valley Monthly Averages).
https://www.nps.gov/yose/planyourvisit/weather.htm

* NOAA National Centers for Environmental Information (NCEI) (2021). 1991-2020 U.S. Climate Normals: Yosemite Park Headquarters, CA (Station USC00049855).
https://www.ncei.noaa.gov/data/normals-monthly/1991-2020/access/USC00049855.csv

* Raphael, Marilyn N. (2003, July 21). The Santa Ana Winds of California. Earth Interactions, 7(8), 1-13.
https://doi.org/10.1175/1087-3562(2003)007%3C0001:TSAWOC%3E2.0.CO;2

* Western Water (2026, March 3). California snowpack still below average after storms.
https://www.western-water.com/2026/03/03/california-snowpack-still-below-average-after-storms/

* California Nevada River Forecast Center (CNRFC / NOAA) (2026, February 12). Water Resources Update - February 12, 2026.
https://www.cnrfc.noaa.gov/WaterResourcesUpdates/report.php?date=20260212

* U.S. Fish and Wildlife Service (2016, February 22). Klamath Dam Removal Overview Report for the Secretary of the Interior.
https://www.fws.gov/sites/default/files/documents/Full%20SDOR%20accessible%20022216.pdf

* Oregon Public Broadcasting (2025, October 28). Klamath River temperatures changed dramatically after dam removal. That's helping salmon swim farther upstream.
https://www.opb.org/article/2025/10/28/klamath-river-temperatures-dam-removal-salmon-upstream/

* North Coast Journal (2025, November 6). Klamath River Ecosystem Booming One Year After Dam Removal. https://www.northcoastjournal.com/news-2/klamath-river-ecosystem-booming-one-year-after-dam-removal/

* World Weather Attribution (2026, March 20). Record-shattering March temperatures in Western North America virtually impossible without climate change. https://www.worldweatherattribution.org/record-shattering-march-temperatures-in-western-north-america-virtually-impossible-witho

* National Integrated Drought Information System (NIDIS) / Drought.gov (2026, October 16). Drought Status Update for California-Nevada. https://www.drought.gov/drought-status-updates/drought-status-update-california-nevada-2026-10-16