Ecology & Evolution | Jan 23, 2026 | 9 min read | By Bryan White

# Feast, Famine, and Fire: The Bornean Orangutan's Struggle in a Changing Biosphere

Biology * Environment * Conservation

## Introduction: The Red Ape at the Precipice

In the stratified canopies of Borneo's dipterocarp and peat swamp forests, the Bornean orangutan (Pongo pygmaeus) acts as an ecological keystone species. As Earth's largest arboreal frugivore, this "gardener of the forest" drives seed dispersal and forest regeneration across carbon-dense tropical ecosystems.[1] Yet the species stands as a critical bellwether for Southeast Asian biodiversity in the Anthropocene. Classified as Critically Endangered by the IUCN, Pongo pygmaeus declined by over 60% between 1950 and 2010, with projections indicating an 86% loss from its 1950 baseline.[2] Although aggregate estimates hover near 104,700 individuals, severe fragmentation isolates many subpopulations in non-viable forest patches.[3] This report synthesizes recent ecological, genetic, and conservation data across Borneo's three orangutan subspecies.

## Taxonomy, Genetics, and the Architecture of Divergence

Unlike the Sumatran (Pongo abelii) and Tapanuli (Pongo tapanuliensis) orangutans, the Bornean species comprises three distinct subspecies adapted to divergent floristic zones:

* Northwest Bornean (P. p. pygmaeus): Ranging across Sarawak and northern West Kalimantan (including Danau Sentarum, Betung Kerihun, and Lanjak Entimau), this most endangered lineage numbers fewer than 3,000 individuals.[3] Recalibrated nest decay studies revealed that late-1990s counts were inflated, masking severe population collapse outside protected parks and leaving isolated fragments vulnerable to a population bottleneck.[4]

* Southwest Bornean (P. p. wurmbii): The largest and most populous subspecies (35,000-40,000 individuals) inhabits the peat swamp and lowland tracts of Central and West Kalimantan, including Sebangau and Tanjung Puting.[3] Its concentration on drained organic peat domes leaves it acutely vulnerable to El Ni o megafires.[5]

* Northeast Bornean (P. p. morio): Restricted to Sabah and East Kalimantan (~11,000-15,000 individuals), this smallest subspecies evolved robust mandibles and thick molar enamel to process fibrous bark and hard seeds during fruit shortages.[2] Despite dietary resilience, hunting and agricultural conversion reduced its Sabah numbers by over 90% over the past two centuries.[2][3]

Maintaining intraspecific genetic integrity requires forensic screening via mitochondrial DNA (such as the ND5 gene on mitochondria) before rehabilitant release to prevent outbreeding depression across divergently adapted subspecies.[6][7] Simultaneously, canopy fragmentation blocks male dispersal, halting gene flow and accelerating genetic drift and inbreeding depression.[8]

## Physiological and Behavioral Ecology in a Volatile Climate

Bornean dipterocarp forests reproduce in irregular "masting" events -- synchronized mass fruiting triggered by El Ni o droughts every 2 to 10 years.[9] Orangutans survive this boom-and-bust cycle via a "feast and famine" metabolic strategy: gorging on lipid-rich seeds to store fat during masts, then subsisting on bark, cambium, and leaves while catabolizing fat stores via beta-oxidation during multi-year inter-mast intervals.[1][9] Studies in Central Kalimantan's Mungku Baru mosaic show that apes track asynchronous fruiting across adjacent peat swamp and heath (Kerangas) forests -- a buffer lost in plantation monocultures.[3][5]

Daily survival also depends on arboreal nest engineering, where orangutans weave stable sleeping platforms using "greenstick fractures" of living branches.[10] Because nest decay rates vary widely (from 150 days in heath forests to 500 days in peat swamps), researchers now pair transect surveys with drone imagery and YOLO deep-learning models to improve census accuracy.[4][11] Functionally, passage through the orangutan gut (24-48 hours) scarifies large seeds and deposits them across long-distance Time Local Convex Hull (T-LoCoH) dispersal kernels, sustaining the regeneration of high-wood-density trees that anchor the forest carbon sink.[1][12]

## Deforestation, Megaprojects, and the Pyrocene

In 2024, Indonesia recorded its third consecutive year of rising deforestation, driven by pulpwood, palm oil, and infrastructure across Kalimantan.[13][14] In West Kalimantan, pulpwood supplier PT Mayawana Persada cleared over 33,000 hectares of rainforest (2021-2023),[15] while palm oil firm PT Equator Sumber Rezeki cleared High Conservation Value forests near Danau Sentarum in 2025,[16] and PEFC drew scrutiny in late 2025 for certifying PT Industrial Forest Plantation despite documented habitat clearance.[14] Illegal drainage canals also continue to threaten protected peat swamp ecosystem reserves.[16] Concurrently, two major infrastructure corridors bisect P. p. morio range: Indonesia's new capital city, Nusantara (IKN) in East Kalimantan, where the Balikpapan toll road severs buffer corridors around the Sungai Wain Protection Forest,[17] and Malaysia's Pan Borneo Highway, which threatens connectivity across Sabah's Class 1 Tawai Forest Reserve.[18][19]

Compounding mechanical clearing is the "Pyrocene" synergy between drained peatlands and the El Ni o-Southern Oscillation. During the 2023 El Ni o, official Indonesian Ministry of Environment and Forestry figures documented 1.16 million hectares burned across Indonesia -- a roughly fivefold surge over 2022 -- with severe wildfire smoke across Borneo's primate habitats.[5][15] Physiological monitoring at Tuanan Research Station demonstrates that toxic peat smoke suppresses flanged male "long calls," forces females and juveniles into lethargy to combat oxidative stress, and drives severe ketosis, muscle wasting, and mass displacement.[5]

## Conservation Paradigms and Regional Status

Mounting habitat pressures have forced a reassessment of conservation strategies. A 2025 IUCN review warned that wild-to-wild translocations suffer high post-release mortality and risk "green-lighting" concession clearing by removing resident apes.[7][20] Instead, field studies in Sabah's Kinabatangan show that orangutans can

disperse and forage across mixed oil-palm mosaics when hunting is eliminated and natural forest patches and riparian corridors are retained for nesting.[21][22] In fire-prone peatlands, hydrological restoration has proven transformative: by constructing 991 canal-blocking dams across Sebangau National Park, the Borneo Nature Foundation re-wetted 7,500 hectares and achieved zero forest loss within its working area during the 2024 dry season.[23]

Across Borneo's political jurisdictions, outcomes remain sharply divergent: in Sabah, populations have stabilized in the Lower Kinabatangan (~867 apes), Deramakot sustainable forestry reserve (1.9 individuals/km ), and the reforested Bukit Piton reserve;[24][25][26] in Sarawak, community-based conservation with Indigenous Iban partners boosted the Ulu Sungai Menyang population from 115 in 2014 to 167 in 2025;[27][28] whereas Kalimantan remains the crisis frontier between peat restoration and industrial clearing.[15][17]

### Table 1: Summary of Bornean Orangutan Status by Subspecies

| Subspecies | | Common Name | | IUCN Status | | Estimated Population | | Primary Threats (2024-2025) | | Key Conservation Focus Areas |

| P. p. pygmaeus | | Northwest Bornean | | Critically Endangered | | < 3,000 | | Habitat fragmentation, pulpwood expansion, smallholder farming | | Danau Sentarum - Betung Kerihun (West Kalimantan), Batang Ai - Lanjak Entimau (Sarawak) |

| P. p. wurmbii | | Southwest Bornean | | Critically Endangered | | ~35,000 - 40,000 | | Peat fires (El Ni o), large-scale drainage, palm oil conversion | | Sebangau National Park, Tanjung Puting, Mawas (Central Kalimantan) |

| P. p. morio | | Northeast Bornean | | Critically Endangered | | ~11,000 - 15,000 | | Infrastructure (Nusantara Capital, Pan Borneo Hwy), mining, timber | | Kinabatangan Floodplain, Kutai National Park, Deramakot (Sabah/East Kalimantan) |

## Conclusion: Coexistence and Landscape Resilience

The plight of Pongo pygmaeus reflects a stark duality between industrial habitat loss and remarkable biological resilience. While pulpwood expansion, megaproject highways, and El Ni o peat fires continue to erode primary rainforest, the success of hydrological damming in Sebangau, canopy restoration in Bukit Piton, and Indigenous stewardship in Sarawak prove that targeted interventions work. Securing the Bornean orangutan requires moving beyond isolated fortress parks to enforce zero-deforestation supply chains, protect every connective forest fragment, and manage working landscapes so the "gardener of the forest" can continue sustaining Borneo's tropical biosphere.

Updates... (Nov 22, 2026): Added verified author and publication dates across eight previously undated citations: Strobel (2013) on Animal Diversity Web (Ref [1]), AidEnvironment (Jun 24, 2022, Ref [14]), Rainforest Action Network (Jul 14, 2025, Ref [16]), Gokkon / Eco-Business (Mar 10, 2023, Ref [17]), The Vibes (Dec 1, 2024, Ref [18]), The Borneo Project (Apr 2, 2020, Ref [19]), WWF UK (Oct 15, 2025, Ref [25]), and Sabah Wildlife Department (2020, Ref [26]), plus specific dates on Refs [13], [22], and [27].

## References

* [1] Strobel, B. (2013). Pongo pygmaeus (Bornean orangutan). Animal Diversity Web, University of Michigan Museum of Zoology. https://animaldiversity.org/accounts/Pongo_pygmaeus/

* [2] IUCN. (2016). Pongo pygmaeus: Ancrenaz, M., Gumal, M., Marshall, A. J., Meijaard, E., Wich, S. A., & Husson, S. (amended version of 2024 Red List entry). The IUCN Red List of Threatened Species. https://doi.org/10.2305/IUCN.UK.2024-1.RLTS.T17975A259043172.en

* [3] Voigt, M., Wich, S. A., Ancrenaz, M., Meijaard, E., Abram, N., Banes, G. L., et al. (2018). Global Demand for Natural Resources Eliminated More Than 100,000 Bornean Orangutans. Current Biology, 28(5), 761-769.e5. https://doi.org/10.1016/j.cub.2018.01.053

* [4] Yuliani, E. L., Bakara, D. O., Ilyas, M., Russon, A. E., Salim, A., Sammy, J., Sunderland-Groves, J. L., & Sunderland, T. C. H. (2023). Bornean orangutan Pongo pygmaeus pygmaeus population estimate within and around Danau Sentarum National Park, Kapuas Hulu, West Kalimantan. Conservation Science and Practice, 5(4), e12916. https://doi.org/10.1111/csp2.12916

* [5] Barrow, E. J., & Erb, W. M. (2026). A systematic review of the impacts of El Ni o-driven drought, fire and smoke on non-human primates in Southeast Asia. Oryx, 1-11. https://doi.org/10.1017/S0030605325102445

* [6] Christy Lavenia, Dwi Sendi Priyono, Donan Satria Yudha, & Tuty Arisuryanti. (2024). Species Identification of Rehabilitated Critically Endangered Orangutans Through DNA Forensic: Implication for Conservation. Tropical Life Sciences Research, 35(1), 123-137. https://doi.org/10.21315/tlsr2024.35.1.7

* [7] Banes, G. L., Galdikas, B. M. F., & Vigilant, L. (2016). Reintroduction of confiscated and displaced mammals risks outbreeding and introgression in natural populations, as evidenced by orang-utans of divergent subspecies. Scientific Reports, 6(1), 22026. https://doi.org/10.1038/srep22026

* [8] IUCN SSC Conservation Planning Specialist Group (2017). Orang-utan Global Population Management Workshop Report. https://www.cpsg.org/documents/orang-utan-global-population-management-workshop-report

* [9] Kanamori, T., Kuze, N., Bernard, H., Malim, T. P., & Kohshima, S. (2010). Feeding ecology of Bornean orangutans (Pongo pygmaeus morio) in Danum Valley, Sabah, Malaysia: a 3-year record including two mast fruitings. American Journal of Primatology, 72(9), 820-840. https://doi.org/10.1002/ajp.20848

* [10] van Casteren, A., Sellers, W. I., Thorpe, S. K. S., Coward, S., Crompton, R. H., Myatt, J. P., & Ennos, A. R. (2012). Nest-building orangutans demonstrate engineering know-how to produce safe, comfortable beds. Proceedings of the National Academy of Sciences, 109(18), 6873-6877. https://doi.org/10.1073/pnas.1200902109

* [11] Simon, D., Pang, K., Bili, R., Ong, S.-Q., & Bernard, H. (2025). Bornean orangutan nest identification using computer vision and deep learning models to improve conservation strategies. PeerJ, 13, e20333. https://doi.org/10.7717/peerj.20333

* [12] Tarszisz, E., Tomlinson, S., Harrison, M. E., Morrogh-Bernard, H. C., & Munn, A. J. (2018). An ecophysiologically informed model of seed dispersal by orangutans: linking animal movement with gut passage across time and space. Conservation Physiology, 6(1), coy013. https://doi.org/10.1093/conphys/coy013

* [13] LiCAS.news (Jan 31, 2025). Indonesia's deforestation worsens for third year as development expands. https://www.licas.news/2025/01/31/indonesias-deforestation-worsens-for-third-year-as-development-expands/

* [14] AidEnvironment (Jun 24, 2022). Orangutan habitat at risk - corporate concessions need to protect forests. https://aidenvironment.org/publications/orangutan-habitat-forest/

* [15] Greenpeace International (2024). Massive deforestation in Borneo destroying orangutan habitat. https://www.greenpeace.org/international/press-release/65881/massive-deforestation-in-borneo-destroying-orangutan-habitat/

* [16] Rainforest Action Network (Jul 14, 2025). Orangutan Emergency: Fresh Evidence Exposes Illegal Deforestation for Palm Oil. https://www.ran.org/forest-frontlines/orangutan-emergency-fresh-evidence-exposes-illegal-deforestation-for-palm-oil/

* [17] Gokkon, B. (Mar 10, 2023). Conservation, Indigenous rights issues haunt Indonesia's new capital. Eco-Business. https://www.eco-business.com/news/conservation-indigenous-rights-issues-haunt-indonesias-new-capital/

* [18] The Vibes (Dec 1, 2024). Experts raise alarm over Pan-Borneo Highway's alignment through Tawai Forest Reserve. https://www.thevibes.com/index.php/articles/news/104201/experts-raise-alarm-over-pan-borneo-highways-alignment-through-tawai-

* [19] The Borneo Project (Apr 2, 2020). Ecological Impacts of Roads: The Pan Borneo Highway. https://borneoproject.org/ecological-impacts-of-roads-in-malaysian-borneo/

* [20] IUCN (2025). Best practice guidelines for orangutan wild-to-wild translocation. https://portals.iucn.org/library/sites/library/files/documents/2025-031-En.pdf

* [21] Ancrenaz, M., Oram, F., Ambu, L., Lackman, I., Ahmad, E., Elahan, H., Kler, H., Abram, N. K., & Meijaard, E. (2015). Of Pongo, palms and perceptions: a multidisciplinary assessment of Bornean orang-utans Pongo pygmaeus in an oil palm context. Oryx, 49(3), 465-472. https://doi.org/10.1017/S0030605313001270

* [22] GAPKI (Feb 15, 2025). Studies: Orangutans Can Coexist With Palm Plantations. https://gapki.id/en/news/2025/02/15/studies-orangutans-can-coexist-with-palm-plantations/

* [23] Borneo Nature Foundation (2025). 2024 Impact Report. https://www.borneonaturefoundation.org/wp-content/uploads/2025/09/Borneo-Nature-Foundation-Impact-Report-2024.pdf

* [24] Zoo Basel / HUTAN (2025). Hutan Annual Report 2024. https://www.zoobasel.ch/uploads/files/website/Hutan_Annual_Report_2024_red-size.pdf

* [25] WWF UK (Oct 15, 2025). Orangutans returning to regenerated forest. https://www.wwf.org.uk/success-stories/orangutans-returning-regenerated-forest

* [26] Sabah Wildlife Department (2020). Orangutan Action Plan for Sabah 2020-2029. https://sabiis.sabah.gov.my/sites/default/files/uploads/download/2025-11/2020-2029.Orangutan-Action-Plan-for-Sabah_1.pdf

* [27] The Star (Nov 21, 2025). Sarawak sees rise in Orangutan population,
conservation efforts pay off.
https://www.thestar.com.my/news/nation/2025/11/21/sarawak-sees-rise-in-orangutan-population-conservation-efforts-pay-off

* [28] WWF Malaysia (2025). Joint Media Release: New Population Estimate Shows
Encouraging Signs for Ulu Sungai Menyang's Wild Orangutans - But Challenges Remain.
https://www.wwf.org.my/?34526/Joint-Media-Release-New-Population-Estimate-Shows-Encouraging-Signs-for-Ulu-Sungai-Menya