Ecology & Evolution | Jun 3, 2026 | 7 min read | By Bryan White

Gut Feeling: The Iron-Rich Cells Guiding Birds Across the Globe

Biology * Environment * Evolution

Introduction: Relocating the Avian Magnetic Compass

Migratory birds and homing pigeons (Columba livia) navigate hundreds of kilometers across unfamiliar terrain by integrating visual landmarks, olfactory gradients, and solar or stellar cues.[1] When heavy cloud cover or nocturnal flight obscures celestial references, diverse vertebrate taxa -- including birds, sea turtles, bats, and elasmobranch fishes -- orient using Earth's geomagnetic field.[1] Yet despite a century of behavioral proof that animals possess an internal magnetic compass, its definitive anatomical locus and cellular transduction pathway remained unresolved.[1][3]

On May 28, 2026, a landmark study published in Science by University of Bonn cell biologist Clivia Lisowski, immunologist Christian Kurts, Max Planck Institute of Animal Behavior ornithologist Martin Wikelski, and colleagues fundamentally relocated the primary organ of avian magnetoreception.[1] Rather than residing exclusively in cranial structures, the pigeon's primary magnetic sensor consists of iron-rich resident immune cells -- hepatic macrophages -- embedded within the liver parenchyma in direct contact with autonomic nerve fibers.[1][4]

Legacy Cranial Hypotheses versus Visceral Magnetometry

Historically, avian magnetoreception research centered on two cranial models: retinal radical-pair chemistry and upper-beak magnetite receptors.[2][1] The radical-pair model proposed that short-wavelength photons strike flavoprotein cryptochromes -- light-sensitive proteins in the avian retina -- generating spin-correlated radical pairs whose singlet-triplet interconversion is modulated by geomagnetic inclination.[1][3] However, no definitive downstream neural readout was isolated, and light-dependent cryptochromes cannot explain magnetoreception in nocturnal or subterranean dark-navigating species.[1] Conversely, the trigeminal hypothesis localized putative magnetite clusters to the upper beak skin innervated by the ophthalmic branch of the trigeminal nerve, where local anesthesia disrupted laboratory magnetic conditioning; however, high-resolution electron microscopy subsequently indicated that many of those beak deposits were non-sensory scavenger macrophages rather than dedicated mechanoreceptors.[2][3]

| Anatomical Site | | Proposed Sensor Mechanism | | Primary Neural Pathway | | 2026 Magnetometry Analysis |

| Retina (Eyes) | | Light-dependent radical-pair reactions via cryptochrome proteins. | | Optic nerve | | Negligible specific magnetic cell isolation. |

| Upper Beak | | Magnetite clusters in connective tissue. | | Ophthalmic branch of the trigeminal nerve | | Low intrinsic magnetic response in target cells. |

| Inner Ear | | Electromagnetic induction within the vestibular canals. | | Vestibulocochlear nerve | | Not identified as a primary sensory locus. |

| Liver (Hepatic Tissue) | | Iron-loaded macrophages acting as superparamagnetic compasses. | | Hepatic branch of the vagus nerve | | Highest concentration of magnetic nanoparticles; positive isolation. |

The pivot to visceral organs originated when Kurts noted that murine splenic macrophages routinely adhered to magnetic cell-separation columns after phagocytizing senescent erythrocytes and recycling their iron.[1] Teaming up with Wikelski, Lisowski used vibrating sample magnetometry and magnetic cell sorting to screen pigeon beaks, retinas, brains, spleens, and livers.[1][6] While ocular and beak tissues showed negligible intrinsic magnetic susceptibility, the spleen and especially the liver exhibited strong magnetic responses driven by millions of iron-loaded hepatic macrophages.[1][6]

Erythrophagocytosis and Superparamagnetic Nanoparticles

As a central regulator of systemic homeostasis, the vertebrate liver houses resident reticuloendothelial macrophages (Kupffer cells) within hepatic sinusoids that clear senescent red blood cells via erythrophagocytosis.[7][8] Inside macrophage lysosomes, hemoglobin is degraded and heme iron is liberated.[8] Because unbound ferrous iron catalyzes Fenton reactions that generate cytotoxic reactive oxygen species from oxygen metabolites, macrophages safely sequester extracted iron within ferritin and hemosiderin complexes.[9][10]

Co-author Ulf Wiedwald (University of Duisburg-Essen) demonstrated that within pigeon hepatic macrophages, this recycled iron is mineralized into uniform iron-oxide nanoparticles small enough to exhibit superparamagnetism.[1][5] Unlike bulk ferromagnetic crystals that retain permanent remnant magnetization, superparamagnetic nanoparticles have zero net magnetic moment in the absence of an external field due to ambient thermal fluctuations, preventing intracellular particle clumping.[5] Yet when exposed to Earth's weak geomagnetic field (~30-60 microteslas), their magnetic moments rapidly and reversibly align along geomagnetic flux lines, passively reorienting whenever the flying pigeon alters its heading.[5]

In Vivo Clodronate Depletion and GPS Homing Telemetry

To test whether pigeons actively use these superparamagnetic liver cells to navigate in flight, the team trained 34 homing pigeons on a 20-kilometer (12.4-mile) flight corridor back to their home loft in Konstanz, Germany.[1][5] Because pigeons operate a hierarchical navigation system -- preferentially relying on a time-compensated solar compass when the sun is visible and switching to their geomagnetic compass only under heavy overcast -- experimental releases were stratified across sunny and completely overcast days.[1][3]

Twenty-four hours prior to release, half of the cohort received intravenous clodronate encapsulated in synthetic lipid vesicles (liposomes), a standard immunology protocol that selectively targets mononuclear phagocytes while sparing non-phagocytic cells.[1][11] Once hepatic macrophages engulf the vesicles, lysosomal phospholipases digest the lipid bilayer and release clodronate into the cytosol, triggering selective macrophage apoptosis and depleting ~80% of iron-bearing liver

macrophages.[1][11]

| Subject Group | | Physiological State | | Weather Condition | | Primary Navigation Cue Available | | Flight Outcome | | Average Return Time |

| Control | | Intact Macrophages | | Overcast | | Magnetic (Hepatic) | | Direct, purposeful homing | | ~70 minutes |

| Treated | | Depleted Macrophages | | Sunny | | Solar (Visual) | | Direct, purposeful homing | | ~70 minutes |

| Treated | | Depleted Macrophages | | Overcast | | None (Both Disabled) | | Erratic, profound disorientation | | Failed (Returned next day) |

GPS flight telemetry revealed a decisive double dissociation:[1]

* Untreated Controls on Overcast Days: Retaining intact hepatic macrophages, control birds compensated seamlessly for the obscured sun, flying straight home in an average of ~70 minutes.[1]

* Macrophage-Depleted Birds on Sunny Days: Despite losing 80% of their magnetic liver macrophages, clodronate-treated pigeons released under clear skies engaged their primary solar compass and returned home in ~70 minutes, proving that clodronate caused no motor, visual, or cognitive impairment.[1]

* Macrophage-Depleted Birds on Overcast Days: Deprived simultaneously of visual solar cues by thick cloud cover and of hepatic magnetoreception by clodronate depletion, treated pigeons exhibited erratic, random flight trajectories.[1] None returned to the Konstanz loft on the day of release; all remained stranded overnight until cloud cover broke the following day and restored solar cues.[1]

Vagal Neuro-Immune Transduction and Immuno-Sensation

For superparamagnetic alignment in the abdomen to guide flight steering, directional signals must rapidly reach the central nervous system.[4] High-resolution electron microscopy of pigeon liver sections revealed that iron-loaded hepatic macrophages lie in direct structural apposition to unmyelinated autonomic nerve terminals.[1][4] As geomagnetic torque realigns intracellular iron-oxide nanoparticles, the resulting mechanical deformation of the macrophage cell membrane -- or localized paracrine transmitter release -- is hypothesized to depolarize adjacent sensory neurons, initiating an ascending action potential along the hepatic branch of the vagus nerve.[4][6][12] Predominantly composed of visceral sensory afferents that project via the solitary tract to the brainstem, hypothalamus, and avian hippocampal spatial-mapping circuits, the hepatic vagus nerve provides an established anatomical conduit for visceral magnetoreception.[4][12]

Conclusion: The Immuno-Sensation Paradigm

By demonstrating that iron-recycling hepatic macrophages act as a functional geomagnetic compass in Columba livia, Lisowski, Kurts, and Wikelski establish a new systems biology paradigm termed "immuno-sensation" -- wherein the immune organ system perceives planetary geophysical fields and interfaces directly with neural navigation circuits.[1][5][13] Rather than invalidating retinal radical-pair mechanisms entirely, hepatic magnetoreception likely operates within a redundant,

multi-organ sensory hierarchy that co-opts conserved vertebrate iron metabolism to guide global animal migration.[1][4]

Updates... (Nov 22, 2026): Corrected the homing training corridor distance to 20 kilometers (12.4 miles) for unit consistency with the Konstanz field trials, replaced an unrelated liver-fibrosis SNP citation with Soares and Hamza (Mar 2016) on macrophage iron metabolism, removed a local television syndicate item, replaced the undated ImmunoSensation homepage link with the May 29, 2026 University of Bonn Medical Faculty announcement, and added verified publication dates across all 13 references.

References

* [1] Lisowski, C., Quetting, M., Klaus, D., Lazarevski, L., Seep, L., et al. (May 28, 2026). Homing pigeon navigation relies on superparamagnetic macrophages under overcast conditions. Science, 392(6801), 985-991. -- https://doi.org/10.1126/science.ady2486

* [2] Mora, C. V., Davison, M., Wild, J. M., & Walker, M. M. (Nov 2004). Magnetoreception and its trigeminal mediation in the homing pigeon. Nature, 432(7016), 508-511. -- https://doi.org/10.1038/nature03077

* [3] Science News (May 28, 2026). Homing pigeons may use a surprising navigation mechanism. -- https://www.sciencenews.org/article/pigeon-navigation-iron-liver-cells

* [4] National Geographic (May 28, 2026). How do pigeons sense Earth's magnetic field? Scientists say they've solved the mystery. -- https://www.nationalgeographic.com/animals/article/pigeons-magnetic-field

* [5] Popular Science (May 28, 2026). Pigeons use their livers to sense Earth's magnetic field. -- https://www.popsci.com/environment/pigeon-liver-magnetic-field-navigate/

* [6] Max-Planck-Gesellschaft (May 28, 2026). Pigeons navigate using magnetic sensors in their livers. -- https://www.mpg.de/26511515/pigeons-navigate-using-magnetic-sensors-in-their-livers

* [7] Graham, R. M., Chua, A. C., Herbison, C. E., Olynyk, J. K., & Trinder, D. (Sep 21, 2007). Liver iron transport. World Journal of Gastroenterology, 13(35), 4725. -- https://doi.org/10.3748/wjg.v13.i35.4725

* [8] Korolnek, T., & Hamza, I. (May 7, 2015). Macrophages and iron trafficking at the birth and death of red cells. Blood. -- https://doi.org/10.1182/blood-2014-12-567776

* [9] Soares, M. P., & Hamza, I. (Mar 2016). Macrophages and Iron Metabolism. Immunity, 44(3), 492-504. -- https://doi.org/10.1016/j.immuni.2016.02.016

* [10] Kouroumalis, E., Tsomidis, I., & Voumvouraki, A. (Mar 10, 2025). HFE-Related Hemochromatosis May Be a Primary Kupffer Cell Disease. Biomedicines, 13(3), 683. -- https://doi.org/10.3390/biomedicines13030683

* [11] Buiting, A. M. J., Zhou, F., Bakker, J. A. J., van Rooijen, N., & Huang, L. (Jun 1996). Biodistribution of clodronate and liposomes used in the liposome mediated macrophage 'suicide' approach. Journal of Immunological Methods. --

https://doi.org/10.1016/0022-1759(96)00034-8

* [12] Prescott, S. L., & Liberles, S. D. (Feb 2022). Internal senses of the vagus nerve. Neuron. -- https://doi.org/10.1016/j.neuron.2021.12.020

* [13] University of Bonn Medical Faculty / ImmunoSensation (May 29, 2026). Immune cells in the liver help pigeons navigate. -- https://www.medfak.uni-bonn.de/en/research/infrastructure/news/immune-cells-in-the-liver-help-pigeons-navigate