Ecology & Evolution | Nov 15, 2026 | 5 min read | By Bryan White

Mapping the Frontiers of the Marine Fossil Record Reveals Where Undiscovered Prehistoric Species Still Hide

Ecology & Evolution * Biodiversity

A comprehensive 540-million-year synthesis published in Science Advances by paleontologists Thomas A. Neubauer and Mathias Harzhauser maps the physical and geographic boundaries of the shallow-marine animal fossil record, revealing both where centuries of collecting have nearly saturated known rock outcrops and where unexplored strata still hold high potential for discovering new prehistoric species. By cross-referencing massive georeferenced fossil occurrence datasets against global stratigraphic databases such as Macrostrat, the study creates a quantitative "treasure map" for future paleontological fieldwork and museum collection digitization.

Understanding how completely scientists have sampled Earth's ancient oceans is one of paleobiology's foundational challenges. Apparent rises and falls in prehistoric biodiversity across the Phanerozoic eon -- the 540-million-year span of complex multicellular animal life -- can reflect genuine evolutionary radiations and mass extinctions, or simply uneven rock preservation, regional collecting bias, and uncataloged museum drawers.

Merging Global Fossil Occurrences With Exposed Rock Strata

To separate true biological patterns from sampling gaps, Neubauer and Harzhauser integrated global records of shallow-marine metazoan (multicellular animal) fossil collections with spatial maps of exposed sedimentary rock formations preserved across each geological stage. Rather than asking only how many fossil species have been named in a given period, the researchers evaluated how much of the Earth's surviving, accessible shallow-marine rock area has already been sampled by paleontologists.

Modeling historical discovery curves against available outcrop area showed that for a majority of Phanerozoic time intervals, geographic coverage is approaching saturation. In heavily studied regions such as Western Europe and North America, additional field expeditions into classic marine formations are projected to yield diminishing returns in spatial coverage, even though rare or microscopic taxa may still emerge through intensive resampling.

| Analytical Dimension | | Scope / Data Integration | | Key Finding in Science Advances |

| Temporal Span | | ~540 million years (Phanerozoic eon) | | Evaluates shallow-marine metazoan fossil completeness |

| Geological Baseline | | Macrostrat global stratigraphic database | | Quantifies accessible vs. sampled sedimentary rock area |

| Sampling Trajectory | | Historical collection accumulation models | | Most geological intervals show diminishing geographic returns |

| High-Potential Frontiers | | Under-sampled accessible marine strata | | Includes surprisingly young Cenozoic and Miocene intervals |

Why Younger Cenozoic Strata Still Hold Major Surprises

One of the most counterintuitive results of the analysis is that the greatest untapped potential for paleontological discovery does not lie exclusively in the deepest, oldest Cambrian or Paleozoic beds. Instead, Neubauer and Harzhauser identified surprisingly high proportions of under-studied, geologically accessible shallow-marine rock in relatively younger Cenozoic intervals, including parts of the Miocene epoch.

Several historical and logistical factors explain why younger marine strata can remain under-sampled relative to their total outcrop area. Classic paleontological campaigns disproportionately targeted iconic mass- extinction boundaries, early Cambrian evolutionary radiations, or Mesozoic marine-reptile and ammonite beds in accessible temperate regions. Meanwhile, extensive Neogene and Cenozoic coastal deposits across parts of the tropics, the Southern Hemisphere, and less-surveyed continental margins have received far less systematic taxonomic inventory relative to their vast surface exposure.

Separating Preservation Bias From Evolutionary History

Distinguishing between geological availability and biological diversity is essential for interpreting deep-time macroevolution. If a geological stage appears unusually species-poor simply because few marine basins from that age are exposed above sea level -- or because exposed rocks lie in politically or logistically remote terrain -- paleobiologists risk mistaking a sampling deficit for an ancient ecological crisis.

By explicitly staking out the physical limits of surviving shallow-marine rock, the Science Advances framework gives researchers an empirical baseline for normalizing diversity curves. When scientists compare fossil richness across two geological epochs, knowing whether one interval has reached near-complete outcrop saturation while another remains largely unexplored allows statistical subsampling models to correct for unequal geographic effort.

A Strategic Roadmap for Fieldwork and Museum Audits

Beyond theoretical macroevolution, the study offers immediate practical guidance for funding agencies, field teams, and natural history museums. Rather than dispersing limited field budgets across already-saturated classic localities, expedition planners can use the mapped frontier zones to target accessible, fossil-bearing marine strata that remain unrepresented in global occurrence databases.

Equally important, the authors highlight the role of "dark data" already sitting inside museum cabinets. In many cases, fossils from under-represented regions or younger Cenozoic intervals have already been collected by regional geological surveys or legacy expeditions, yet remain uncataloged, un-digitized, or omitted from international paleobiology repositories. Pairing targeted new fieldwork in mapped frontier rocks with systematic digitization of existing collections offers the fastest path toward completing Earth's 540-million-year marine archive.

References & Primary Sources

* [1] Neubauer, T. A., & Harzhauser, M. (2026). Staking out the frontiers: The limits of the metazoan shallow marine fossil record. Science Advances. -- https://doi.org/10.1126/sciadv.aeh3920

* [2] Macrostrat Geological Data Platform (2026). Global Stratigraphic & Lithological Database. -- https://macrostrat.org/

* [3] ScienceDaily (2026). Staking out the frontiers of the shallow marine fossil record. -- https://www.sciencedaily.com/

* [4] NDTV (2026). Scientists Create Treasure Map To Find Fossils Of Undiscovered Prehistoric Species. -- https://www.ndtv.com/