Ecology & Evolution | Oct 30, 2026 | 5 min read | By Bryan White

Silk Moth Experiments Confirm the 'Molecular Cookie Cutter' Mechanism of Wing Evolution

Evolutionary Developmental Biology * Lepidoptera * Apoptosis

Across the more than 160,000 described species of butterflies and moths (order Lepidoptera), adult wings take on an extraordinary diversity of silhouettes -- from the hooked, dead-leaf profiles of anglewing butterflies to the long, twisted hindwing tails that help Luna moths (Actias luna) deflect the echolocation sonar of hunting bats. Three decades ago, developmental biologists studying butterflies proposed a striking explanation for how these intricate outlines form: rather than growing outward into their final shape, embryonic wing buds start as oversized, rounded sheets that are trimmed back before adulthood by a "molecular cookie cutter" of programmed cell death.

Now, a seven-year experimental study published in Evolution & Development by Andrei Sourakov at the Florida Museum of Natural History's McGuire Center for Lepidoptera and Biodiversity provides the first direct experimental manipulation of wing shape in moths. Working with North American giant silk moths (family Saturniidae) -- specifically the Luna moth (Actias luna) and the Polyphemus moth (Antheraea polyphemus) -- Sourakov demonstrated that perturbing extracellular signaling gradients during pupation simultaneously alters both the physical outline of the adult wing and its eyespot color patterns.

How the 'Molecular Cookie Cutter' Sculpts a Wing

Inside a caterpillar, the future adult wings develop as internal pouches of epithelial cells called wing imaginal discs . Throughout the final larval instar and early pupal stage, these discs expand into broad, paddle-like sheets bordered by a peripheral fluid channel known as the border lacuna .

Rather than retaining all of the tissue outside that channel, the developing insect activates a sharp genetic boundary line across the wing disc:

* The Boundary Transcription Factor (Cut): Originally characterized in butterfly wing discs by Sean Carroll and colleagues in 1994 and mapped in detail by William Macdonald and coworkers in 2010, the homeodomain transcription factor Cut is expressed in a narrow strip of cells tracing the exact future edge of the adult wing.

* Morphogen Positioning (Wingless / Wnt Signaling): Diffusible signaling proteins from the Wnt family (specifically wingless , or wg) radiate from the wing margin, establishing a concentration gradient that tells cells where the Cut boundary should sit relative to internal wing-blade identity genes such as vestigial (vg) and Distal-less (Dll).

* Peripheral Programmed Cell Death (Apoptosis): Once the boundary is locked in, cells lying outside the Cut perimeter activate caspase enzymes and undergo synchronized apoptosis (programmed cell death). Like dough trimmed away outside a

pastry cutter, the outer zone dissolves before the adult moth emerges from its chrysalis or cocoon, carving out scalloped notches, curved wingtips, and elongated hindwing tails.

| Developmental Component / Perturbation | | Molecular Target or Pathway | | Normal Role in Saturniid Moths | | Experimental Effect of Heparin / Dextran Sulfate | | Evolutionary Significance |

| Wing Margin Boundary ("Cookie Cutter") | | Cut homeodomain transcription factor + peripheral caspase apoptosis | | Trims larval wing imaginal disc along the border lacuna to carve tails and wingtips | | Shifts excision line inward or outward, altering wing aspect ratio, curvature, and hindwing tail geometry | | Demonstrates that major wing-shape leaps require only shifting an apoptosis boundary |

| Extracellular Morphogen Gradient | | Wingless (wg / Wnt1) & WntA heparan-sulfate proteoglycan transport | | Establishes positional coordinates for both margin excision and pigment bands | | Sulfated polysaccharides enhance/broaden Wnt diffusion across pupal wing epithelium | | Couples physical wing silhouette directly to color pattern coordinates |

| Luna Moth (Actias luna) | | Hindwing acoustic decoy tails + submarginal eyespot rings | | Long twisted hindwing tails deflect bat sonar; cryptic green leaf mimicry | | Truncated or reshaped wing margins paired with expanded dark marginal and ocellar pigment zones | | Links anti-bat structural defenses and visual camouflage to a shared developmental switch |

| Polyphemus Moth (Antheraea polyphemus) | | Cryptic dead-leaf forewings + startled hindwing ocelli (eyespots) | | Sharp submarginal bands and transparent mica-like eyespot windows | | Transforms cryptic leaf-mimic banding into diffuse, hyper-melanized aposematic (warning-like) patterns | | Explains rapid evolutionary transitions between camouflage and startle coloration |

Rewiring Moth Wings with Heparin and Dextran Sulfate

Testing whether nocturnal moths rely on the same morphogen-guided cookie cutter as diurnal butterflies required finding a way to shift the positional blueprint inside a living pupa without halting metamorphosis . Over seven years of rearing experiments on Actias luna and Antheraea polyphemus , Sourakov injected micro-quantities of two sulfated polysaccharides -- heparin and dextran sulfate -- during the sensitive prepupal and early pupal windows.

In insect developmental biology , cell-surface molecules called heparan sulfate proteoglycans (HSPGs) act like molecular stepping stones that regulate how far Wnt/Wingless signaling proteins travel between cells. Introducing exogenous heparin or dextran sulfate alters that extracellular transport matrix, allowing Wnt morphogens to diffuse farther across the developing wing sheet than they normally would.

In butterflies, heparin injections were previously known to smudge or expand pigment bands. In Sourakov's giant silk moths, however, the treatment achieved something never before documented in moths: it physically reshaped the adult wing contour . Depending on the precise timing and dosage of the polysaccharide injection, adult Luna and Polyphemus moths emerged with altered forewing curvature, modified wing-tip

angles, and reshaped hindwing margins -- confirming that the Wnt morphogen gradient directly sets the coordinates of the apoptotic cookie cutter.

From Leaf Camouflage to Startle Eyespots in a Single Step

Equally striking was how the same chemical perturbation reorganized the moths' visual defenses. Under normal conditions, giant silk moths balance two contrasting survival strategies: their forewings mimic dry oak leaves or green foliage (crypsis), while their hindwings carry bold, ringed eyespots (ocelli) flashed suddenly to startle predatory birds (deimatic or aposematic display).

When heparin or dextran sulfate broadened the Wnt signaling gradient in Antheraea polyphemus and Actias luna , the boundary bands and eyespot rings expanded dramatically, converting camouflaged leaf-like wing surfaces into bold, high-contrast patterns that closely resemble distant relative species in the genus Actias and Automeris . Rather than requiring dozens of independent mutations to change wing shape and wing coloration separately, a single regulatory tweak to morphogen diffusion simultaneously coordinates both traits.

Why Shared Developmental Toolkits Accelerate Lepidopteran Evolution

Evolutionary biologists have long wondered how moths and butterflies evolved such rapidly diverging wing geometries -- including the independent evolution of anti-bat hindwing tails at least four separate times within Saturniidae alone. By confirming that nocturnal silk moths use the same Wnt-coupled apoptotic boundary mechanism proposed for butterflies, the Florida Museum study shows that this developmental toolkit dates back to the common ancestor of macrolepidopterans well over 100 million years ago.

Because the larval wing disc always starts with a generous surplus of tissue and relies on a movable chemical gradient to draw the cut line, natural selection can rapidly lengthen a sonar-jamming tail, hook a leaf-mimicking wingtip, or enlarge a startle eyespot simply by shifting where the molecular cookie cutter lands.

References & Primary Sources

* [1] Sourakov, A. (2026). On wing pattern and wing shape evolution in giant silk moths: Lessons from wing development in Luna and Polyphemus moths. Evolution & Development. -- https://doi.org/10.1111/ede.70058

* [2] Florida Museum of Natural History / Phys.org. (2026). Experimental study on moths supports the 'molecular cookie cutter' idea proposed for butterflies. -- https://www.floridamuseum.ufl.edu/

* [3] Macdonald, W. P., Martin, A., & Reed, R. D. (2010). Butterfly wings shaped by a molecular cookie cutter. Evolution & Development, 12(3), 296-304. -- https://doi.org/10.1111/j.1525-142X.2010.00415.x

* [4] Carroll, S. B., Gates, J., Keys, D. N., et al. (1994). Pattern formation and eyespot determination in butterfly wings. Science, 265(5168), 109-114. -- https://doi.org/10.1126/science.7912449

* [5] Barber, J. R., Leavell, B. C., Keener, A. L., et al. (2015). Moth tails divert bat attack: Evolution of acoustic deflection. Proceedings of the National Academy of

Sciences, 112(9), 2812-2816. -- https://doi.org/10.1073/pnas.1421926112