Space & Astronomy | Jan 7, 2026 | 6 min read | By Bryan White

Observational Evidence for Relativistic Frame-Dragging in Black Hole System AT2020afhd

Space * Astronomy * Physics

Introduction: Relativistic Frame-Dragging in the Strong-Field Regime

In Albert Einstein's General Theory of Relativity, gravitation arises from the geometric curvature of four-dimensional spacetime rather than a classical Newtonian force. Among the most consequential predictions of the Kerr metric for rotating masses is relativistic frame-dragging -- the Lense-Thirring effect -- first derived by Josef Lense and Hans Thirring in 1918.[1] Rather than rotating passively in static space, a spinning massive body drags local inertial frames along its azimuthal rotation vector, imparting a gravitomagnetic torque on nearby orbiting matter.[1]

While weak-field frame-dragging around Earth was confirmed by NASA's Gravity Probe B and laser-ranged geodetic satellites, directly isolating Lense-Thirring precession in the strong gravitational field near a supermassive black hole (SMBH) event horizon has remained a major observational challenge. High-cadence multi-wavelength observations of the tidal disruption event (TDE) AT2020afhd have now provided the first definitive detection of synchronized disk-jet Lense-Thirring coprecession, establishing a direct dynamical probe of black hole spin and relativistic accretion geometry.[1][2]

Relativistic Mechanics of Tidal Disruption Events

A tidal disruption event occurs when a star's periastron passage carries it inside the tidal radius of a central SMBH, where differential gravitational gradients across the stellar diameter exceed the star's self-gravity. Approximately half of the shredded stellar debris becomes gravitationally unbound and escapes at hyperbolic speeds, while the bound fraction falls back toward the black hole to circularize into a superheated, optically thick accretion disk.[1]

Because the disrupted star approaches on a random trajectory relative to the black hole's equatorial plane, the nascent accretion disk is typically inclined at an angle with respect to the SMBH spin axis. Within a few gravitational radii ($R_g = GM/c^2$), the gravitomagnetic Lense-Thirring torque acts on the tilted fluid rings. When internal hydrodynamic viscosity (or magnetic coupling) communicates faster than the differential precession rate -- a regime known as the Bardeen-Petterson or rigid-tilt limit -- the inner accretion disk precesses coherently as a solid body around the black hole spin vector.[1] If the inner disk launches a magnetically collimated relativistic jet via the Blandford-Znajek or Blandford-Payne mechanism, the jet axis must precess in lockstep with the disk, sweeping both thermal X-ray emission and non-thermal radio synchrotron radiation across the observer's line of sight.[1][4]

Multi-Wavelength Campaign and the 19.6-Day Coprecession Signal

First detected in 2020 by the Zwicky Transient Facility (ZTF) in the nucleus of the galaxy LEDA 145386 (~400 million light-years away), AT2020afhd initially followed a standard TDE luminosity decay curve over three years.[2] In January 2024, however, optical surveys recorded a dramatic rebrightening event, prompting an intensive international campaign led by the National Astronomical Observatories of the Chinese Academy of Sciences (NAOC) and Cardiff University:[1][2]

* High-Energy X-Ray Spectroscopy and Timing: The Neil Gehrels Swift Observatory, NICER aboard the International Space Station, and XMM-Newton sampled thermal continuum radiation from the innermost accretion disk (~102-103 km scale).[1]

* Centimeter-Wave Radio Interferometry: The Karl G. Jansky Very Large Array (VLA) and e-MERLIN tracked non-thermal synchrotron emission from the collimated relativistic outflow.[1]

* Optical Photometry: Continuous cadence from the Xinglong 2.16-meter and Lijiang 2.4-meter telescopes constrained outer-envelope reprocessing.[1]

Approximately 215 days after the January 2024 optical rebrightening, the X-ray flux began executing high-amplitude quasi-periodic oscillations (QPOs), modulating by an order of magnitude on a strict 19.6-day period.[1][2] Crucially, VLA and e-MERLIN radio light curves exhibited synchronized 19.6-day modulations with zero statistically significant phase lag relative to the X-ray peaks.[1] Because X-rays trace the inner accretion funnel while radio waves originate far downstream in the jet, zero-lag synchronization proves that the inner accretion disk and relativistic jet are rigidly coupled and coprecessing as a single physical structure under Lense-Thirring torque.[1]

| Evolutionary Epoch | | Observational Window | | Multi-Wavelength Signature | | Astrophysical Interpretation |

| Initial Tidal Flare | | 2020 | | Optical transient discovered by ZTF in LEDA 145386. | | Tidal disruption and initial fallback of stellar debris.[2] |

| Primary Decay Phase | | 2020 - Late 2023 | | Monotonic decline in optical/UV bolometric output. | | Viscous drainage of the primary debris stream.[2] |

| Secondary Rebrightening | | January 2024 (Day 0) | | Rapid optical brightening; onset of high-cadence campaign. | | Delayed circularization shock or secondary mass-transfer episode.[1][2] |

| Ultrafast Outflow (UFO) Epoch | | Days 74 - 215 | | Blueshifted X-ray absorption lines decelerating from 0.19c to 0.0097c. | | Launch and rapid deceleration of metal-enriched radiation-driven disk winds.[3] |

| Rigid Coprecession Epoch | | Day 215+ | | Synchronized 19.6-day, 10x amplitude QPOs in X-ray and radio bands. | | Global Lense-Thirring coprecession of the inner disk and relativistic jet.[1] |

Ultrafast Outflow Deceleration and Disk State Transitions

High-resolution X-ray spectroscopy prior to the precession phase also captured the complete lifecycle of Ultrafast Outflows (UFOs) -- relativistic accretion disk winds

never previously tracked from birth to termination in a TDE.[3] Emerging at Day 74 post-rebrightening and peaking in column density between Days 172 and 194, these winds vanished abruptly at Day 215 just as the 19.6-day coprecession signal stabilized.[3]

During a 10-day interval, the outflow velocity decelerated precipitously from 0.19c (19% of light speed) down to 0.0097c (<1% of light speed), accompanied by an inverse correlation between outflow velocity and gas ionization parameter.[3] This rapid braking indicates either a widening of the wind launching cone or progressive enrichment of the circumnuclear gas with high-opacity heavy elements such as iron and oxygen synthesized in the disrupted stellar core.[3] The cessation of the UFO at Day 215 marks a structural state transition in which the radiatively inflated disk settled into a compact, geometrically thick torus capable of sustaining rigid-body Lense-Thirring precession.[1][3]

Constraining Black Hole Spin and Comparative TDE Demographics

In general relativistic hydrodynamics, the rigid-body Lense-Thirring precession period depends strictly on the black hole mass (MBH), the dimensionless Kerr spin parameter (a*), and the outer radius of the precessing disk ring. Using host-galaxy scaling relations for LEDA 145386 to fix MBH at several million solar masses, the observed 19.6-day period constrains the central SMBH in AT2020afhd to a low-to-moderate spin parameter of a* ~ 0.1-0.2.[1] A maximally spinning Kerr black hole (a* -> 1) would induce precession on timescales of only a few days; thus, the 19.6-day cadence independently confirms a slowly rotating SMBH probably grown via chaotic, multi-directional accretion episodes rather than coherent prolonged disk spin-up.[1]

| System Parameter | | AT2020afhd | | AT2020ocn | | Swift J1644+57 |

| Modulation Period | | 19.6 Days[1][2] | | ~15 Days[1] | | ~2.7 Days[1] |

| Observational Bands | | Synchronized X-ray & Radio (Zero Lag) | | Soft X-ray Only (No Radio QPO) | | Hard X-ray & Radio (Blazar-like alignment) |

| Inferred Kerr Spin (a*) | | Low (0.1 - 0.2)[1] | | Weakly constrained | | Moderate to High |

| Physical Mechanism | | Global Rigid Disk-Jet Lense-Thirring Coprecession | | Inner Accretion Disk Precession | | Jet Nutation / Internal Shock Instability |

Conclusion: A New Relativistic Gyroscope for Time-Domain Astronomy

By capturing zero-lag 19.6-day X-ray and radio oscillations alongside transient ultrafast disk winds, observations of AT2020afhd confirm that rotating supermassive black holes drag local spacetime and torque misaligned accretion disk-jet systems into coherent precession. As wide-field time-domain facilities such as the Vera C. Rubin Observatory Legacy Survey of Space and Time (LSST) expand the census of tidal disruption events into the thousands, multi-wavelength coprecession timing will serve as a precision relativistic gyroscope to map the spin distribution of quiescent black holes across cosmic history.

Updates... (Nov 22, 2026): Enriched all four primary journal DOIs and Chinese

Academy of Sciences release entries with verified exact CrossRef and institutional publication dates (including updating Yuan et al., ApJ 996(1), to Jan 1, 2026).

References & Primary Sources

* [1] Wang, Y., Lin, Z., Wu, L., Lei, W.-H., Wei, S., et al. (Dec 10, 2025). Detection of disk-jet coprecession in a tidal disruption event. Science Advances, 11(50), eady9068. -- https://doi.org/10.1126/sciadv.ady9068

* [2] Chinese Academy of Sciences (Dec 12, 2025). Scientists Discover First Clear Observation of Disk-Jet Co-precession in Tidal Disruption Event. -- https://english.cas.cn/newsroom/research_news/phys/202512/t20251212_1136062.shtml

* [3] Lin, Z., Wang, Y., Bu, D.-F., Mao, J., & Liu, J. (Aug 10, 2025). Delayed Launch of Ultrafast Outflows in the Tidal Disruption Event AT2020afhd. The Astrophysical Journal Letters, 989(1), L9. -- https://doi.org/10.3847/2041-8213/adf205

* [4] Yuan, H.-Y., Wu, H.-Z., & Lei, W.-H. (Jan 1, 2026). Propagation of a Precessing Jet in the Envelope of Tidal Disruption Events. The Astrophysical Journal, 996(1), 42. -- https://doi.org/10.3847/1538-4357/ae1eed