Health & Medicine | Jan 8, 2026 | 8 min read | By Bryan White

Panzootic Bird Flu: A Comprehensive Analysis of the H5N1 Influenza Crisis in the United States (2024-2026)

Biology * Public Health * Epidemiology

Introduction: The Multi-Species Evolution of Clade 2.3.4.4b

Since its initial identification in Southern China in 1996, Highly Pathogenic Avian Influenza (HPAI) A(H5N1) has undergone continuous evolutionary expansion. Throughout 2024-2026, the dominant clade 2.3.4.4b confronting the United States has transformed from a seasonal avian pathogen into a year-round, multi-species panzootic. Exploiting the ecological interfaces where intensive livestock operations intersect migratory flyways, H5N1 now poses a tripartite threat: a persistent epizootic across U.S. dairy cattle (reaching 22 affected states by November 2026), continuous attrition of commercial and backyard poultry exceeding 166 million depopulated birds since 2022, and entrenched spillover across terrestrial scavengers and coastal marine mammals -- permanently altering the North American viral ecosystem.[1][12][13]

Simultaneously, human zoonotic infections have continued to accumulate, reaching 74 confirmed U.S. human cases by November 2026 (up from 71 cases and two fatalities recorded through early January 2026), demonstrating that clade 2.3.4.4b is repeatedly probing the mammalian species barrier.[14] Examining the convergent polymerase mutations of genotypes B3.13 and D1.1, the emergence of the H5N5 reassortant, and the contrasting mechanics of agricultural versus wild transmission clarifies why containing H5N1 requires a coordinated One Health firewall.

Viral Architecture and Genomic Divergence: Genotypes B3.13, D1.1, and H5N5

As orthomyxoviruses possessing an eight-segment negative-sense RNA genome, Influenza A viruses evolve rapidly through both point mutation and genetic reassortment inside co-infected host cells. Viral tropism is governed by two surface proteins: Hemagglutinin (HA), which binds host sialic acid receptors, and Neuraminidase (NA), which cleaves sialic acid linkages to release nascent virions. While human seasonal influenza binds alpha-2,6 sialic acid receptors concentrated in the upper respiratory tract, the H5 hemagglutinin of clade 2.3.4.4b preferentially binds avian-type alpha-2,3 receptors abundant in avian intestinal tracts, bovine mammary alveoli, and the human conjunctiva and lower lung.

Between 2024 and 2026, genomic surveillance tracked three distinct epidemiological lineages across North America:

* Genotype B3.13 (The Bovine Specialist): Originating from a late 2023 reassortment event, B3.13 drove the multi-state U.S. dairy cattle outbreak. Its hallmark adaptation is PB2 M631L within the viral polymerase complex, which enables the avian polymerase to replicate RNA inside the cooler mammalian nucleus by binding the shorter mammalian host cofactor ANP32A.[3]

* Genotype D1.1 (The Wild Bird Challenger): Representing independent de novo spillovers from migratory waterfowl into dairy herds in Nevada, Arizona, and Wisconsin during 2025, D1.1 carries the PB2 D701N mutation, which enhances nuclear import of the viral polymerase via host importin-alpha.[4][5][10]

* The H5N5 Reassortant: Identified in November 2025 in a fatal Washington State human infection linked to backyard poultry, this reassortant pairs the clade 2.3.4.4b H5 hemagglutinin with a North American N5 neuraminidase.[2][6]

| Feature | | Genotype B3.13 (Bovine Strain) | | Genotype D1.1 (Wild Bird Strain) | | H5N5 Reassortant |

| Emergence Context | | Wild bird spillover (Late 2023) | | Independent spillovers (2025) | | Pacific Flyway reassortment (Nov 2025) |

| Primary Reservoir | | U.S. Dairy Cattle | | Migratory Waterfowl / Spillover Herds | | Wild Birds / Backyard Poultry |

| Key Polymerase Mutation | | PB2 M631L (Mammalian ANP32A binding) | | PB2 D701N (Importin-alpha nuclear import) | | Variable / Under genomic characterization |

| Bovine Clinical Presentation | | Mastitis, thickened milk, sharp yield drop | | Often subclinical / asymptomatic in cattle | | Not established in dairy cattle |

| Human Clinical Impact | | Predominantly conjunctivitis / mild illness | | Severe pneumonia; 1 Fatality (Louisiana) | | Severe pneumonia; 1 Fatality (Washington) |

The Mammalian Breach: Dairy Herds, Poultry Fortresses, and Marine Pinnipeds

The spread of H5N1 across U.S. dairy cattle -- spanning 19 states by early 2026 and expanding to 22 states including Oregon and Washington by late 2026 -- upended conventional respiratory models of influenza.[1][14] Because the virus exhibits intense tropism for bovine mammary epithelium, milk titers exceed respiratory secretions by orders of magnitude, turning unsterilized milking machine teat cups into high-efficiency mechanical fomites alongside secondary exposure in dry cows.[1][7] Critically, the December 2025 detection of genotype D1.1 in an asymptomatic Wisconsin dairy herd via the USDA's National Milk Testing Strategy (NMTS) proved that subclinical bulk-tank surveillance is essential.[8] While commercial High-Temperature Short-Time (HTST) pasteurization completely inactivates infectious H5N1, raw unpasteurized milk remains a direct public health hazard.[9]

Meanwhile, migratory flyways -- particularly the Pacific, Central, and Mississippi corridors -- continue to seed wetlands, where virions persist in cold water and drive trophic scavenger cycles across raptors, raccoons, and harbor seals in Washington's Puget Sound.[11][12][13] In commercial poultry, where bio-exclusion failures allow dust or fomite entry into 50,000-bird barns, outbreaks have forced the culling of over 166 million birds since 2022 (predominantly egg-laying flocks, including 76,210 birds in the first week of January 2026 alone), while peri-urban backyard flocks consistently precede commercial county outbreaks by roughly 9 days.[11][16]

Clinical Reality and Comparative Spread Dynamics

Among the 71 confirmed U.S. human cases cataloged by January 2026 -- 41 linked to dairy cattle, 24 to poultry culling, and 6 to backyard or undetermined exposures -- clinical outcomes bifurcated sharply by genotype and exposure route, with cumulative U.S. detections reaching 74 cases by November 2026 (including a late-2026 fatal dairy worker infection in Washington State alongside the earlier Washington H5N5 and Louisiana D1.1 fatalities).[6][14][15] Dairy workers exposed to B3.13 milk splashes have predominantly developed conjunctivitis and mild upper respiratory symptoms, whereas fatal lower-respiratory infections underscored how intra-host viral adaptation or novel reassortants can elevate clinical severity.[6][15]

| Jurisdiction / Cohort (Early-2026 Baseline & Late-2026 Updates) | | Human Case Burden | | Primary Exposure Context & Outcome |

| California & Michigan | | High (>20 in CA; Moderate in MI) | | Occupational dairy parlor exposure (Genotype B3.13; conjunctivitis). |

| Colorado | | High (>10 cases) | | Commercial poultry depopulation cluster events. |

| Washington State | | Moderate (Includes late-2025 H5N5 & late-2026 dairy fatalities) | | Backyard poultry (H5N5) and occupational dairy farm exposures. |

| Louisiana | | Low (1 Case; 1 Death) | | Backyard poultry / wild bird exposure; fatal D1.1 infection with intra-host evolution. |

| Missouri | | Low (1 Case) | | Hospitalized sentinel detection with no confirmed animal contact. |

| U.S. Cumulative Total | | 74 Confirmed Cases (as of Nov 2026; 71 by Jan 2026) | | Predominantly agricultural occupational zoonosis (Human R0 < 1). |

From an epidemiological modeling perspective, H5N1 exploits two complementary transmission regimes defined by their Basic Reproduction Number (R0):

* Wild Ecological Spread (Variable Intensity / Unlimited Extent): In wild bird populations, R0 fluctuates seasonally with migratory stopover density while cold-water environmental persistence maintains an abiotic reservoir in wetlands. Trophic cascades carry the virus from waterfowl into raptors, terrestrial mesocarnivores, and densely hauled-out marine pinnipeds.[11][12][13] While heterogeneous immunity prevents uniform flock mortality in wild ducks, geographic spread across the Pacific, Central, Mississippi, and Atlantic flyways is functionally unbounded.

* Agricultural Network Spread (Explosive Intensity / Anthropogenic Extent): Inside commercial poultry barns housing 20,000 to 50,000 genetically uniform birds, a single biosecurity breach via windborne dust or contaminated footwear drives R0 above 10, necessitating rapid depopulation.[11][16] In dairy operations, unsterilized milking clusters sustain high intra-herd transmission, while interstate shipment of subclinical heifers and shared feed or veterinary transport trucks connect distant agricultural counties into a small-world transmission network.[1][7]

Where open dairy wastewater lagoons attract migratory waterfowl along major flyways,

these two regimes fuse into a self-sustaining amplification loop: wild reservoirs continuously introduce novel genotypes like D1.1, and intensive livestock infrastructure amplifies mammalian exposure.[4][8]

Conclusion: Maintaining the One Health Firewall

Through 2026, HPAI A(H5N1) clade 2.3.4.4b has entrenched itself across North American wildlife, poultry, and dairy livestock. Although the virus has not acquired sustained human-to-human aerosol transmission -- keeping human R0 below 1 -- convergent mammalian polymerase adaptations (PB2 M631L in B3.13 and PB2 D701N in D1.1) alongside severe human infections demonstrate ongoing evolutionary plasticity.[3][4][6][15] Preventing a pandemic transition requires maintaining an uncompromising One Health firewall: mandatory national bulk-milk silo testing, strict milking-parlor disinfection, personal protective equipment for agricultural workers, and real-time genomic surveillance across the wildlife-livestock interface.

Updates... (Nov 16, 2026): By November 2026, U.S. dairy cattle H5N1 detections expanded to 22 states (including initial detections in Oregon and Washington in October-November 2026), while cumulative CDC-confirmed U.S. human cases rose to 74 (including a fatal Washington State dairy worker case), with no sustained person-to-person transmission detected.

Works Cited

* Caserta, L. C., Frye, E. A., Butt, S. L., Laverack, M., Nooruzzaman, M., Covaleda, L. M., et al. (Jul 25, 2024). Spillover of highly pathogenic avian influenza H5N1 virus to dairy cattle. Nature, 634, 669-676. https://doi.org/10.1038/s41586-024-07849-4

* Whatcom County Health and Community Services (Nov 20, 2025). 11/20/25 Provider Alert: Case of H5N5 Influenza Confirmed in Washington. https://providers.whatcomcounty.org/2025/11/11-19-25-provider-alert-case-of-h5n5-influenza-confirmed-in-washington/

* Dholakia, V., Quantrill, J. L., Richardson, S. A. S., Pankaew, B., Brown, R. E., Yang, J., et al. (Jan 16, 2026). Polymerase mutations underlie early adaptation of H5N1 influenza virus to dairy cattle and other mammals. Nature Communications. https://doi.org/10.1038/s41467-026-68306-6

* USDA Animal and Plant Health Inspection Service (APHIS) (2025). The Occurrence of Another Highly Pathogenic Avian Influenza (HPAI) Spillover from Wild Birds into Dairy Cattle. https://www.aphis.usda.gov/sites/default/files/dairy-cattle-hpai-tech-brief.pdf

* Virological.org (Feb 21, 2025). Timing and molecular characterisation of the transmission to cattle of H5N1 influenza A virus genotype D1.1, clade 2.3.4.4b. https://virological.org/t/timing-and-molecular-characterisation-of-the-transmission-to-cattle-of-h5n1-influenza-a-virus-genotype-d1-1

* World Health Organization (WHO) (Nov 2025). Avian Influenza A(H5N5) - United States of America. https://www.who.int/emergencies/disease-outbreak-news/item/2025-DON590

* CIDRAP (Mar 13, 2025). H5N1 dairy cow study finds sustained milk production drop, extensive transmission across herd.

https://www.cidrap.umn.edu/avian-influenza-bird-flu/h5n1-dairy-cow-study-finds-sustained-milk-production-drop-extensive

* USDA APHIS (Dec 14, 2025). USDA Confirms Highly Pathogenic Avian Influenza in a Dairy Herd in Wisconsin. https://www.aphis.usda.gov/news/agency-announcements/usda-confirms-highly-pathogenic-avian-influenza-dairy-herd-wisconsin

* Public Health Ontario (2024). Rapid Review: Survivability of Influenza A(H5N1) in Milk. https://www.publichealthontario.ca/-/media/Documents/A/24/ah5n1-survivability-influenza-milk.pdf?rev=69786e6f11fd402c8aba52

* Crespo-Bellido, A., Trov o, N. S., Puryear, W., et al. (Dec 23, 2025). Emergence of D1.1 reassortant H5N1 avian influenza viruses in North America. bioRxiv. https://doi.org/10.64898/2025.12.19.695329

* Damodaran, L., Jaeger, A., & Moncla, L. H. (Dec 20, 2024). Intensive transmission in wild, migratory birds drove rapid geographic dissemination and repeated spillovers of H5N1 into agriculture in North America. bioRxiv. https://doi.org/10.1101/2024.12.16.628739

* CIDRAP (Sep 11, 2023). Tests confirm H5N1 avian flu in Washington seals. https://www.cidrap.umn.edu/avian-influenza-bird-flu/tests-confirm-h5n1-avian-flu-washington-seals

* University of California (Jun 6, 2024). Bird flu is increasingly adapting to mammals. https://www.universityofcalifornia.edu/news/bird-flu-increasingly-adapting-mammals

* U.S. Centers for Disease Control and Prevention (CDC) (Nov 20, 2026). H5 Bird Flu: Current Situation. https://www.cdc.gov/bird-flu/situation-summary/index.html

* U.S. Centers for Disease Control and Prevention (CDC) (Dec 26, 2024; updated Feb 28, 2025). Genetic Sequences of Highly Pathogenic Avian Influenza A(H5N1) Viruses Identified in a Person in Louisiana. https://www.cdc.gov/bird-flu/spotlights/h5n1-response-12232024.html

* Farm Credit Administration (Mar 2025). Update on Highly Pathogenic Avian Influenza. https://www.fca.gov/template-fca/about/2025MarchUpdateonHPAI.pdf