Space & Astronomy | Jun 21, 2026 | 8 min read | By Bryan White

Rethinking Space Weather: Inside the DAPHNE Mission's Bottom-Up Approach

Space * Astronomy * Solar System

Introduction: Paradigm Shift in Space Weather Forecasting

Space weather forecasting has traditionally relied upon a top-down paradigm, focusing overwhelmingly on the Sun as the primary driver of disturbances in the near-Earth environment. Solar flares, coronal mass ejections, and the steady outpouring of the solar wind have long been treated as the principal catalysts for geomagnetic storms, auroral activity, and satellite disruptions[1]. However, recent advances in atmospheric physics demonstrate a bidirectional relationship across the atmosphere of Earth: the upper atmosphere is not merely a passive receptor of solar radiation, but a dynamic transition zone shaped equally by terrestrial weather systems propagating upward from below[3].

To resolve this observational gap, the National Aeronautics and Space Administration (NASA) selected the Dynamic Atmosphere-Ionosphere Explorer (DAPHNE) mission for Phase B development in June 2026[3]. Led by the Laboratory for Atmospheric and Space Physics (LASP) at the University of Colorado Boulder, DAPHNE will collect coordinated multi-point measurements of the neutral thermosphere and its coupling with the ionized plasma of the ionosphere[3]. By quantifying how lower-atmospheric dynamics modulate upper-atmospheric variability, DAPHNE aims to refine predictive space weather models that safeguard orbital infrastructure and navigation networks[4].

The Thermosphere-Ionosphere System and Bottom-Up Forcing

The altitude regime between 90 and 300 kilometers forms the critical interface between the lower atmosphere and space, encompassing two overlapping domains: the neutral thermosphere and the ionized ionosphere[3]. Intense solar extreme ultraviolet radiation dissociates molecular nitrogen (N_2) and molecular oxygen (O_2) into atomic oxygen (O) while stripping electrons to generate a conductive plasma shell[7]. Whereas the neutral thermosphere is governed by fluid dynamics, pressure gradients, and the Coriolis effect[2], ionospheric charged particles are constrained by Earth's geomagnetic field[9]. Constant collisional friction transfers momentum and energy between neutral gases and ions[4]. Yet because this middle thermosphere lies above balloon altitudes while exerting severe aerodynamic drag on low-altitude spacecraft, it remains a poorly sampled "thermospheric gap"[5][14].

DAPHNE specifically targets two physical mechanisms driving bottom-up forcing across this boundary: atmospheric gravity waves and the ionospheric wind dynamo[4][9]. Atmospheric gravity waves are buoyancy-driven oscillations excited by flow over topography or deep convective storms that transport energy and momentum from the troposphere into the thermosphere[11][12]. As these waves ascend through exponentially decreasing atmospheric density, their amplitudes grow to conserve kinetic energy until non-linear instabilities cause them to break[11]. Wave breaking

deposits substantial momentum into background high-altitude winds and triggers secondary waves that manifest as Traveling Atmospheric Disturbances and Traveling Ionospheric Disturbances[11][13].

Concurrently, the ionospheric wind dynamo converts neutral wind kinetic energy into electromagnetic fields[9]. As neutral winds collide with ionospheric plasma across geomagnetic field lines, heavier ions are displaced relative to magnetized electrons, driving Pedersen and Hall currents[9]. Spatial gradients in ionospheric conductivity cause charge accumulation and polarization electric fields, which interact with the geomagnetic field to drive horizontal and vertical E-cross-B plasma drifts[9][10]. Without global measurements of the neutral winds powering this dynamo, space weather models cannot separate terrestrial wave forcing from magnetospheric storm inputs[3].

Mission Architecture and Dual-Satellite Stereoscopy

DAPHNE fulfills the Dynamical Neutral Atmosphere-Ionosphere Coupling (DYNAMIC) mission priority established in the 2013 and 2024 Decadal Surveys for Solar and Space Physics[5]. Managed by NASA's Goddard Space Flight Center under the Solar Terrestrial Probes program, DAPHNE operates within a $250 million Fiscal Year 2023 cost cap (excluding launch) for a target launch no earlier than 2029, joining CU Boulder, the Naval Research Laboratory, BAE Systems, UC Berkeley Space Sciences Laboratory, and the University of Central Florida[3][5].

Single-satellite low-Earth-orbit missions suffer from spatial-temporal ambiguity: an observer cannot distinguish whether a measured shift in wind or temperature reflects a stationary geographic structure or a transient propagating wave[4]. DAPHNE resolves this ambiguity by flying two identical spacecraft in a trailing "string-of-pearls" formation within a shared orbital plane[4][15]. Sampling the same atmospheric volume across a controlled time lag enables direct calculation of gravity wave phase speeds, propagation vectors, and vertical momentum fluxes[4].

| Mission Parameter | | Specification |

| Mission Lead | | Aimee Merkel (LASP, CU Boulder) |

| Program Office | | NASA Goddard Space Flight Center (Solar Terrestrial Probes) |

| Flight System | | Two identical satellites in a string-of-pearls formation |

| Target Launch Date | | No earlier than 2029 |

| Cost Cap | | $250 million (FY23 dollars, excluding launch) |

| Primary Scientific Goal | | Differentiating lower-atmosphere forcing from solar forcing in the ionosphere-thermosphere system |

Remote-Sensing Instrumentation Suite

Each DAPHNE spacecraft carries three high-heritage remote-sensing instruments to simultaneously profile neutral winds, temperatures, and composition[5][16].

The Michelson Interferometer for Global High-resolution Thermospheric Imaging (MIGHTI), adapted from NASA's ICON mission, measures horizontal wind vectors and temperatures using Doppler Asymmetric Spatial Heterodyne spectroscopy[8][16]. Rather

than relying on moving mirrors, MIGHTI uses fixed, low-groove-density echelle diffraction gratings (64 grooves/mm) to image Doppler shifts in atomic oxygen airglow emissions at 630.0 nm (red line) and 557.7 nm (green line)[8]. Two orthogonal views offset at 45 deg and 135 deg from the spacecraft velocity vector reconstruct full zonal and meridional wind profiles, while near-infrared sampling of the O2 A-band around 762 nm yields altitude-resolved kinetic temperatures[8][16].

The Far Ultraviolet Imager (FUVI) is a compact 2U nadir-viewing refractive camera utilizing a thermally controlled barium fluoride lens and bandpass filters between 135.6 nm and 168 nm[16][17]. FUVI maps dayside N2 Lyman-Birge-Hopfield bands and nightside 135.6 nm O emissions, resolving density ripples induced by gravity waves with horizontal wavelengths from 100 to over 1,000 kilometers via tomographic inversion[12][16].

Complementing these imagers, the Spectrograph Limb Imager for Thermosphere Ionosphere (PLATO) observes the trailing limb in the ultraviolet to retrieve daytime temperature profiles and the atomic oxygen to molecular nitrogen (O/N2) composition ratio -- a primary diagnostic of the thermodynamic state of the upper atmosphere[7][16]. Thermal expansion during storm or wave-heating events drives the upwelling of N2-rich air, accelerating electron recombination and depleting ionospheric plasma density[7].

| Instrument | | Full Name | | Measurement Technique | | Primary Scientific Targets |

| MIGHTI | | Michelson Interferometer for Global High-resolution Thermospheric Imaging | | Doppler Asymmetric Spatial Heterodyne Spectroscopy | | Altitude profiles of horizontal vector winds (zonal/meridional) and temperatures |

| FUVI | | Far Ultraviolet Imager | | Nadir-viewing refractive ultraviolet imaging | | 2D spatial maps of gravity waves, neutral density variations, and auroral emissions |

| PLATO | | Spectrograph Limb Imager for Thermosphere Ionosphere | | Ultraviolet limb imaging spectrography | | Altitude profiles of daytime O/N2 composition ratio and kinetic temperature |

Constellation Synergy and Operational Impacts

DAPHNE's scientific return will be amplified through coordinated operations with NASA's six-satellite Geospace Dynamics Constellation (GDC) at ~380 kilometers altitude[14][18]. Following earlier budget uncertainty, Congress reinstated ~$100 million for GDC in the final Fiscal Year 2026 NASA appropriations bill to sustain development alongside the Solar Terrestrial Probes queue[20][21]. While GDC performs in-situ measurements of top-down magnetospheric plasma and electromagnetic energy inputs, DAPHNE remotely senses the underlying neutral gas dynamics, closing the upper-atmospheric energy and momentum budget[6][7][18].

Closing this observational gap directly addresses critical operational hazards. Sudden thermospheric heating and expansion produce severe satellite drag spikes, as demonstrated in 2022 when an unpredicted geomagnetic storm deorbited forty newly launched commercial satellites[4][6]. Furthermore, neutral wind dynamos and breaking gravity waves seed ionospheric plasma bubbles that scatter radio and GPS signals --

degrading aviation links, maritime navigation, and centimeter-level precision agriculture workflows such as automated fertilization[1][3][19].

Conclusion: Toward Predictive Upper-Atmospheric Modeling

The selection of the DAPHNE mission for Phase B development marks a decisive shift toward whole-atmosphere heliophysics. By pairing dual-satellite stereoscopic limb and nadir imaging with top-down magnetospheric observatories, DAPHNE will disentangle terrestrial meteorological forcing from solar drivers across the thermospheric gap, transforming space weather forecasting into a rigorous, physically coupled predictive discipline[3][7].

Updates... (Nov 19, 2026): Replaced the secondary bioscience.com.pk news summary with the official LASP University of Colorado Boulder DAPHNE mission page, added full author attributions to the 2024 and 2025 AGU DYNAMIC/DAPHNE abstracts, split grouped inline reference tags, and added numbered reference identifiers.

References & Primary Sources

* [1] Space Weather - NASA Science, https://science.nasa.gov/heliophysics/focus-areas/space-weather/

* [2] NASA Heliophysics, https://science.nasa.gov/heliophysics/

* [3] NASA Mission to Study Space Weather Impacts of Earth's Atmosphere, https://www.nasa.gov/news-release/nasa-mission-to-study-space-weather-impacts-of-earths-atmosphere/

* [4] Laboratory for Atmospheric and Space Physics (LASP), University of Colorado Boulder (2026). DAPHNE: Dynamic Atmosphere-Ionosphere Explorer, https://lasp.colorado.edu/missions/daphne/

* [5] NASA selects LASP mission to study how Earth and space weather connect, University of Colorado Boulder, https://www.colorado.edu/today/2026/06/18/nasa-selects-lasp-mission-study-how-earth-and-space-weather-connect

* [6] 40 Years in the Making: UT Dallas' Next NASA Mission (GDC), https://nsm.utdallas.edu/nasa-gdc-mission/

* [7] Harvey, V. L., et al. (2025). The DYNAMIC-X Mission, AGU Fall Meeting Abstract, https://agu.confex.com/agu/agu25/meetingapp.cgi/Paper/1974071

* [8] Englert, C. R., et al. (2017). High-efficiency echelle gratings for MIGHTI, the spatial heterodyne interferometers for the ICON mission. Applied Optics. https://doi.org/10.1364/ao.56.002090

* [9] Ionospheric Dynamo Process - NICT, https://www.nict.go.jp/publication/shuppan/kihou-journal/journal-vol56no1_2_3_4/journal-vol56no1-4_020307.pdf

* [10] Blanc, M., & Richmond, A. (1980). The ionospheric disturbance dynamo. Journal of Geophysical Research: Space Physics. https://doi.org/10.1029/ja085ia04p01669

* [11] Kumar, S., et al. (2023). Characterization of Gravity Waves in Three Dimensions in the Daytime Thermosphere Using Combined Optical and Radio Measurements and Estimation of Horizontal Neutral Winds. Journal of Geophysical Research: Space Physics. https://doi.org/10.1029/2022ja030954

* [12] Tomographic retrieval of thermosphere gravity waves from the Far Ultraviolet Imager (FUVI) on the future DYNAGLO CubeSat mission | CEDAR, https://cedarscience.org/poster-2024/tomographic-retrieval-thermosphere-gravity-waves-far-ultraviolet-imager-fuvi-future

* [13] Watching the thermosphere respond to disturbances: tracking large-scale thermospheric gravity waves with GOLD | High Altitude Observatory, https://www2.hao.ucar.edu/watching-thermosphere-respond-disturbances-tracking-large-scale-thermospheric-gravity-waves-gold

* [14] DYNAMIC Announcements - Science Office for Mission Assessments (SOMA), NASA, https://soma.larc.nasa.gov/stp/dynamic/announcements.html

* [15] Merkel, A. W., Harding, B. J., Curry, S., et al. (2025). The DAPHNE DYNAMIC mission - AGU Fall Meeting Abstract, https://agu.confex.com/agu/agu25/meetingapp.cgi/Paper/1989416

* [16] McClintock, W. E., Merkel, A. W., Budzien, S., et al. (2024). Science Instrumentation for the DAPHNE DYNAMIC Mission Concept, AGU Fall Meeting Abstract, https://agu.confex.com/agu/agu24/meetingapp.cgi/Paper/1618447

* [17] FUVI - Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, https://lasp.colorado.edu/instruments/fuvi/

* [18] Oberheide, J., et al. (2023). Resolving the tidal weather of the thermosphere using GDC. Frontiers in Astronomy and Space Sciences. https://doi.org/10.3389/fspas.2023.1282261

* [19] GDC STDT Final Report | NASA Science, https://science.nasa.gov/wp-content/uploads/2023/04/Oct_2_4b_GDCSTDTReporttoHPAC_Ridley_Jaynes__20191002.pdf

* [20] AAS Summary of NASA FY 2026 Budget, NASA Watch, https://nasawatch.com/2026/01/16/aas-summary-of-nasa-fy-2026-budget/

* [21] NASA Heliophysics Needs Major Budget Increases, Decadal Survey Argues, AIP FYI, https://www.aip.org/fyi/nasa-heliophysics-needs-major-budget-increases-decadal-survey-argues