Ecology & Evolution | Jan 13, 2026 | 9 min read | By Bryan White

Strength Without Meat: How the Mammalian Body Adapts Regardless of Diet

Biology * Public Health * Food Science

Introduction: The Meat-Protein Paradigm and the 83-Participant Cohort Study

For decades, sports nutrition and exercise physiology have been anchored by the hypothesis that animal-derived proteins are biochemically indispensable for optimal skeletal muscle remodeling and strength acquisition. Because meat, eggs, and dairy possess amino acid profiles closely homologous to human skeletal muscle and exhibit high Digestible Indispensable Amino Acid Scores (DIAAS), plant-based diets were long assumed to induce "anabolic resistance" due to lower leucine density, limiting essential amino acids (such as lysine in grains or methionine in legumes), and plant anti-nutritional factors like phytates and tannins.[1] At the same time, concerns over climate change and epidemiological evidence linking plant-based nutrition to lower cancer and cardiometabolic risk have accelerated the adoption of vegan diets among athletes.[2][3]

A controlled clinical intervention by C rcamo-Regla et al. published in Nutrition (online late 2025; print 2026, PII: S0899900725003648) directly tested whether free-living dietary patterns constrain strength adaptations.[4] Tracking 83 habitual omnivores and vegans (each maintaining their diet for at least six months prior) across a 16-week resistance training or control protocol, the study demonstrated that resistance training significantly improved muscular strength regardless of diet type.[4] By synthesizing these ecological findings with nutritional biochemistry and protein-clamped trials by Hevia-Larra n et al. (2021) and Monteyne et al. (2023),[5][14][15] this report examines how mechanical tension, neuromuscular recruitment, and metabolic redundancy override acute differences in dietary protein kinetics.

Cellular Mechanisms of Skeletal Muscle Adaptation

Skeletal muscle remodeling is governed by the Stimulus-Recovery-Adaptation (SRA) cycle and the principles of bioenergetics and cellular metabolism. When external resistance imposes mechanical tension on muscle fibers, mechanosensors embedded in the sarcolemma (cell membrane) -- such as focal adhesion kinases -- transduce physical force into intracellular biochemical signals.[6] Satellite cells proliferate and fuse to damaged myofibrils, donating nuclei to support parallel deposition of the contractile myofilaments actin and myosin (hypertrophy), while the central nervous system enhances motor-unit synchronization and firing frequency (strength).[5][6] Because strength acquisition during the first 6 to 16 weeks of training is heavily mediated by neural drive and motor coordination, the 83-participant cohort's equal strength gains confirm that neuromuscular adaptation operates independently of animal versus plant protein origin.[4]

At the molecular level, myofibrillar protein synthesis (MPS) is regulated by the Mechanistic Target of Rapamycin Complex 1 (mTORC1), which integrates four upstream inputs: mechanical loading, endocrine growth factors (IGF-1 and insulin), cellular energy status via the AMP-to-adenosine triphosphate (ATP) ratio (AMPK pathway), and intracellular amino acid availability.[7][8] Activated mTORC1 phosphorylates S6K1 and 4E-BP1 to initiate mRNA translation at the ribosome.[7] Although the "Leucine Threshold Hypothesis" notes that leucine binds Sestrin2 to activate mTORC1 -- requiring ~2.0 g per meal in young adults and ~3.0 g in older adults[9][10][12] -- chronic mechanical tension sensitizes the muscle fiber to circulating amino acids and drives ribosome biogenesis, expanding total translational capacity over a 24-hour window.[6][11]

Protein Kinetics, Microbiome Synergy, and Comparative Clinical Trials

While isolated whey protein produces a rapid hyperaminoacidemic peak compared to fiber-bound plant matrices,[1][13] multi-week training trials indicate that cumulative 24-hour amino acid exposure ("Area Under the Curve") and hepatic amino acid pooling compensate for slower digestion kinetics.[1][12] Habitual vegans consuming complementary grains, legumes, nuts, and seeds across the day maintain a complete endogenous amino acid pool.[1] Furthermore, insights from microbiology and the human microbiome indicate that fiber-rich plant diets enrich short-chain fatty acid (SCFA)-producing bacteria such as Prevotella, whose butyrate production strengthens gut barrier integrity and dampens systemic inflammation -- reducing catabolic ubiquitin-proteasome muscle proteolysis while improving cardiometabolic markers.[3]

| Study Parameter | | Hevia-Larra n et al. (2021)[14] | | Monteyne et al. (2023)[5][15] | | C rcamo-Regla et al. (2026) Nutrition[4] |

| Intervention Duration | | 12 Weeks | | 10 Weeks | | 16 Weeks |

| Sample Size & Cohort | | 38 young men (19 vegan, 19 omnivore) | | 22 healthy young adults | | 83 habitual vegans & omnivores (>6 mo tenure) |

| Dietary Protocol | | Protein-clamped at 1.6 g/kg/day (soy vs. whey) | | High-protein whole-food / mycoprotein (~1.8-2.0 g/kg/day) | | Unclamped habitual free-living diet |

| Strength Adaptation | | Equivalent 1RM leg press gains (+97 kg vs. +117 kg, NS) | | Equivalent muscular strength progression | | Significant, equivalent strength gains across diets |

| Hypertrophy / Lean Mass | | Equivalent (+1.2 kg leg lean mass; Type I/II CSA) | | Equivalent (+3.1 kg vegan vs. +2.6 kg omnivore; equal MPS) | | No significant musculoskeletal mass change in either group |

| Primary Mechanistic Takeaway | | Protein quantity (1.6 g/kg) eliminates source differences. | | Whole-food fungal/plant matrices match animal MPS rates. | | Habitual diets support equal strength; mass requires caloric/protein surplus. |

Comparing the 16-week cohort trial against prior controlled trials clarifies the distinction between strength and hypertrophy:[4][14][18]

* Protein-Clamped Equivalence (Hevia-Larra n 2021 & Monteyne 2023): When total daily protein is matched at optimal anabolic thresholds (1.6 g/kg/day) via soy isolate[14] or whole-food fungal mycoprotein,[5][15][16] vegans and omnivores achieve identical daily myofibrillar protein synthesis rates, muscle fiber cross-sectional area (CSA) expansion, and lean mass accretion.

* The "Strength Without Mass" Paradox: In the 16-week free-living trial, both habitual vegans and habitual omnivores gained significant strength, yet neither group exhibited statistically significant gains in total musculoskeletal mass.[4] Because unguided habitual diets typically provide 0.8-1.2 g/kg/day of protein without a dedicated caloric surplus, both dietary groups had ample substrate for neurological strength adaptation but remained below the quantitative threshold required for maximal structural hypertrophy.[4][12]

Practical Guidelines and Methodological Limitations

Synthesizing the 16-week free-living trial with protein-clamped hypertrophy studies yields clear, evidence-based guidelines across athletic and clinical populations:[1][17][19]

* Strength & Recreational Athletics: Habitual plant-based and omnivorous diets support equal strength progression over 16 weeks without rigid protein timing, provided caloric intake and progressive mechanical overload are maintained.[4][19]

* Maximal Hypertrophy (Bodybuilding): To drive significant lean tissue accretion alongside strength, athletes on either diet must intentionally elevate total protein intake to 1.6 g/kg/day using protein-dense whole foods (tofu, tempeh, seitan, mycoprotein) or complementary pea/rice/soy isolates.[14][18][20]

* Healthy Aging & Sarcopenia Prevention: Older adults adopting plant-forward diets for cardiovascular longevity can preserve functional strength through consistent resistance training, while targeting higher per-meal leucine intakes (~2.5-3.0 g) to counter age-related anabolic resistance.[10][17]

Several methodological constraints of the free-living trial warrant future investigation, including granular macronutrient tracking to pinpoint exact g/kg/day intakes,[4] potential motivational differences among habitual vegans, and multi-year longitudinal tracking of micronutrients such as vitamin B12, iron, and zinc.[3][4]

Conclusion: Mechanical Effort as the Great Equalizer

The 16-week investigation of 83 habitual vegans and omnivores refutes the longstanding assumption that animal protein is required for strength adaptation.[4] Together with controlled trials demonstrating identical myofibrillar protein synthesis when daily protein targets are met,[5][14] the evidence underscores the physiological redundancy of the mammalian musculoskeletal system: while total protein quantity governs structural tissue accretion, progressive mechanical tension remains the primary driver of human strength regardless of dietary origin.

Updates... (Nov 21, 2026): Attributed Layman (2024) in Frontiers in Nutrition and Naclerio & Seijo (2019) via verified CrossRef DOIs; replaced advocacy and redundant secondary summaries with peer-reviewed meta-analyses (Morton et al. 2018, Reid-McCann et al. 2025, and Callegari et al. October 2026); added specific

publication dates across all 20 references; and standardized reference list markup.

References & Primary Sources

* [1] Ashtary-Larky, D. (Jul 2025). Are plant-based and omnivorous diets the same for muscle hypertrophy? A narrative review of possible challenges of plant-based diets in resistance-trained athletes. Nutrition, 135, 112742. -- https://doi.org/10.1016/j.nut.2025.112742

* [2] Dinu, M., Abbate, R., Gensini, G. F., Casini, A., & Sofi, F. (Jun 13, 2017; print Nov 22, 2017). Vegetarian, vegan diets and multiple health outcomes: A systematic review with meta-analysis of observational studies. Critical Reviews in Food Science and Nutrition, 57(17), 3640-3649. -- https://doi.org/10.1080/10408398.2016.1138447

* [3] Tomova, A., Bukovsky, I., Rembert, E., Yonas, W., Alwarith, J., Barnard, N. D., & Kahleova, H. (Apr 17, 2019). The Effects of Vegetarian and Vegan Diets on Gut Microbiota. Frontiers in Nutrition, 6, 47. -- https://doi.org/10.3389/fnut.2019.00047

* [4] C rcamo-Regla, R., Alcaide-Leyva, J. M., Zapata-Lamana, R., Martorell-Pons, M., Carrasco-Mar n, F., & Molina-Recio, G. (Aug 2026). Efficacy of a resistance training program on muscle mass and muscle strength in adults following a vegan versus omnivorous diet: A nonrandomized four-arm parallel clinical trial. Nutrition, 148, 113047. -- https://doi.org/10.1016/j.nut.2025.113047

* [5] Monteyne, A. J., Coelho, M. O. C., Murton, A. J., Abdelrahman, D. R., Blackwell, J. R., Koscien, C. P., et al. (Jun 2023). Vegan and Omnivorous High Protein Diets Support Comparable Daily Myofibrillar Protein Synthesis Rates and Skeletal Muscle Hypertrophy in Young Adults. The Journal of Nutrition, 153(6), 1680-1695. -- https://doi.org/10.1016/j.tjnut.2023.02.023

* [6] Schiaffino, S., Reggiani, C., Akimoto, T., & Blaauw, B. (Mar 2, 2021). Molecular Mechanisms of Skeletal Muscle Hypertrophy. Journal of Neuromuscular Diseases, 8(2), 169-183. -- https://doi.org/10.3233/JND-200568

* [7] Yang, M., Lu, Y., Piao, W., & Jin, H. (Jun 8, 2022). The Translational Regulation in mTOR Pathway. Biomolecules, 12(6), 802. -- https://doi.org/10.3390/biom12060802

* [8] Bodine, S. C. (Nov 11, 2022). The role of mTORC1 in the regulation of skeletal muscle mass. Faculty Reviews, 11, 32. -- https://doi.org/10.12703/r/11-32

* [9] Zhang, X., Kapoor, D., Jeong, S.-J., Fappi, A., Stitham, J., Shabrish, V., et al. (Feb 19, 2024). Identification of a leucine-mediated threshold effect governing macrophage mTOR signalling and cardiovascular risk. Nature Metabolism, 6(2), 359-377. -- https://doi.org/10.1038/s42255-024-00984-2

* [10] Witard, O. C., Wardle, S. L., Macnaughton, L. S., Hodgson, A. B., & Tipton, K. D. (Mar 23, 2016). Protein Considerations for Optimising Skeletal Muscle Mass in Healthy Young and Older Adults. Nutrients, 8(4), 181. -- https://doi.org/10.3390/nu8040181

* [11] Nader, G. A., McLoughlin, T. J., & Esser, K. A. (Dec 2005). mTOR function in

skeletal muscle hypertrophy: increased ribosomal RNA via cell cycle regulators. American Journal of Physiology-Cell Physiology, 289(6), C1457-C1465. -- https://doi.org/10.1152/ajpcell.00165.2005

* [12] Layman, D. K. (May 3, 2024). Impacts of protein quantity and distribution on body composition. Frontiers in Nutrition, 11, 1388986. -- https://doi.org/10.3389/fnut.2024.1388986

* [13] Naclerio, F., & Seijo, M. (Aug 29, 2019). Whey protein supplementation and muscle mass: current perspectives. Nutrition and Dietary Supplements, 11, 37-48. -- https://doi.org/10.2147/NDS.S166195

* [14] Hevia-Larra n, V., Gualano, B., Longobardi, I., Gil, S., Fernandes, A. L., Costa, L. A. R., et al. (Feb 18, 2021; print Jun 2021). High-Protein Plant-Based Diet Versus a Protein-Matched Omnivorous Diet to Support Resistance Training Adaptations: A Comparison Between Habitual Vegans and Omnivores. Sports Medicine, 51(6), 1317-1330. -- https://doi.org/10.1007/s40279-021-01434-9

* [15] University of Exeter News (Mar 30, 2023). Vegan protein supports muscle building as effectively as animal protein, according to study. -- https://news.exeter.ac.uk/faculty-of-health-and-life-sciences/vegan-protein-supports-muscle-building-as-effectively-as-animal-protei

* [16] Reid-McCann, R. J., Brennan, S. F., Ward, N. A., Logan, D., McKinley, M. C., & McEvoy, C. T. (Jan 15, 2025; print Jul 1, 2025). Effect of Plant Versus Animal Protein on Muscle Mass, Strength, Physical Performance, and Sarcopenia: A Systematic Review and Meta-analysis of Randomized Controlled Trials. Nutrition Reviews, 83(7), e1581-e1603. -- https://doi.org/10.1093/nutrit/nuae200

* [17] Morton, R. W., Murphy, K. T., McKellar, S. R., Schoenfeld, B. J., Henselmans, M., Helms, E., et al. (Jul 11, 2017; print Mar 2018). A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. British Journal of Sports Medicine, 52(6), 376-384. -- https://doi.org/10.1136/bjsports-2017-097608

* [18] Santini, M. H., Erwig Leit o, A., Mazzolani, B. C., Smaira, F. I., de Souza, M. S. C., & Santamaria, A. (Oct 8, 2025; print Dec 31, 2025). Similar effects between animal-based and plant-based protein blend as complementary dietary protein on muscle adaptations to resistance training: findings from a randomized clinical trial. Journal of the International Society of Sports Nutrition, 22(1), 2568047. -- https://doi.org/10.1080/15502783.2025.2568047

* [19] Isenmann, E., Eggers, L., Havers, T., Schalla, J., Lesch, A., & Geisler, S. (Jan 19, 2023). Change to a Plant-Based Diet Has No Effect on Strength Performance in Trained Persons in the First 8 Weeks -- A 16-Week Controlled Pilot Study. International Journal of Environmental Research and Public Health, 20(3), 1856. -- https://doi.org/10.3390/ijerph20031856

* [20] Callegari, I. O. M., Silva, G. M., & Oliveira, A. G. (Oct 29, 2026). Is plant-based protein supplementation as effective as whey protein? A systematic review and meta-analysis in untrained adults undergoing resistance training. Journal of the International Society of Sports Nutrition, 23(1), 2752317. -- https://doi.org/10.1080/15502783.2026.2752317