Space & Astronomy | Jan 10, 2026 | 7 min read | By Bryan White

Surfing a Supergiant: The Hidden Companion of Betelgeuse

Space * Astronomy * Physics

Introduction: The Variable Supergiant and the Long Secondary Period Conundrum

For nearly a century, the red supergiant Betelgeuse (Orionis) -- an evolved M1-M2 massive star whose extended photosphere would engulf the orbit of Jupiter if placed at the center of our solar system -- has presented a persistent puzzle for stellar astrophysics.[1][3] While its primary ~400-day brightness oscillation is well explained by the radial (opacity) pulsation mechanism in ionization zones, Betelgeuse also exhibits a regular Long Secondary Period (LSP) of approximately 2,170 days (~6 years).[3] Observed across roughly one-third of pulsating asymptotic giant branch (AGB) stars and red supergiants, LSPs have defied single-star pulsation models: a radial mode of 2,170 days would require an unphysically large stellar radius, non-radial gravity modes (g-modes) struggle to reach observable surface amplitudes through deep convective envelopes, and stochastic giant convection cells cannot maintain phase coherence over a century of photometric monitoring.[3][5][10]

Interest in coupling between these cycles intensified during the "Great Dimming" of 2019-2020, when a surface mass ejection condensed into a circumstellar dust cloud at the simultaneous trough of both the 400-day and 2,170-day periods, reducing Betelgeuse's visual brightness by two-thirds.[3][7] Now, an eight-year multi-wavelength campaign led by Andrea Dupree using the Space Telescope Imaging Spectrograph (STIS) aboard the Hubble Space Telescope, synthesizing dynamical models by Jared Goldberg and collaborators alongside ground-based speckle imaging, has resolved the enigma.[1][4][11] Published in The Astrophysical Journal in early 2026, the observations confirm the existence of a low-mass binary companion -- formally designated Siwarha (Orionis B) -- orbiting directly inside Betelgeuse's extended chromosphere and carving a supersonic hydrodynamic wake that modulates the system's ultraviolet and optical spectrum.[8]

Dynamical Constraints and the Discovery of Siwarha (Orionis B)

Because Betelgeuse outshines a solar-mass companion by a factor of roughly 105, direct optical detection is severely hindered by photospheric glare.[13] Consequently, researchers first constrained the companion dynamically by combining decades of radial velocity (RV) spectroscopy and astrometric wobble measurements.[11] Crucially, Betelgeuse's 2,170-day radial velocity curve lags behind its photometric light curve by approximately half an orbit -- a phase relationship incompatible with intrinsic atmospheric pulsation, but diagnostic of an orbiting companion that periodically alters circumstellar extinction along Earth's line of sight.[11][12]

| System Parameter | | Observational / Derived Value | | Astrophysical Interpretation |

| Orbital Period (PLSP) | | ~2,170 days (~5.94 years) | | Directly drives the Long Secondary Period photometric & RV cycle.[11] |

| Companion Mass (MB) | | 0.6 - 1.6 M (~1.2-1.5 M best fit) | | Solar-mass class companion ("Siwarha" / Ori B).[6][11] |

| Orbital Semi-Major Axis | | ~1,850 R (~8.6 AU) | | Approximately 2.3 to 2.5 Betelgeuse stellar radii (R).[8][11] |

| Host Environment | | Inner Extended Chromosphere | | Generates a supersonic bow shock and trailing plasma wake.[4][8] |

| Companion Evolutionary State | | Pre-Main Sequence (PMS) Star | | System age ~8-10 Myr; contracting T Tauri-type object.[13][14] |

| Inspiral Timescale | | ~10,000 years | | Hydrodynamic drag drives decay toward Common Envelope merger.[14] |

Given that Betelgeuse is a rapidly evolved massive star with an age of only 8 to 10 million years, a companion of 0.6 to 1.6 M has not yet contracted onto the hydrogen-burning Main Sequence.[14] Instead, Siwarha is a Pre-Main Sequence (PMS) star -- consistent with upper limits from Chandra X-ray observations and optical speckle imaging at Gemini North -- orbiting at a separation of just ~2.3-2.5 stellar radii (~8.6 AU), deep within the supergiant's tenuous chromosphere.[6][8][13]

Ultraviolet Spectroscopy and Physics of the Chromospheric Wake

Definitive confirmation of the binary interaction came from eight years of Hubble STIS ultraviolet spectroscopy combined with ground-based optical monitoring led by Dupree et al.[4][8] As Siwarha orbits through Betelgeuse's chromosphere at a velocity exceeding the local atmospheric sound speed, it acts like a speedboat plowing through fluid, generating a compressed bow shock and an expanding, turbulent trailing wake of heated plasma.[1][15]

The researchers tracked this wake using chromospheric emission and absorption diagnostics across the 2,170-day orbital phase:

* Mg II h and k Chromospheric Modulation: The singly ionized magnesium doublet near 2800 exhibited systematic flux and profile asymmetries synchronized with Siwarha's 2,170-day orbit, tracing localized heating and density compression in the chromosphere.[8]

* Fe II and Mn I Trailing Absorption: During orbital transit (when Siwarha passes in front of Betelgeuse relative to Earth), blueshifted radial velocity components dominate. Immediately following transit -- as the companion moves toward quadrature and occultation -- the dense trailing wake swings across the line of sight between Betelgeuse's photosphere and Earth, producing a pronounced surge in ultraviolet Fe II outflow absorption and optical circumstellar Mn I absorption.[8][15]

* Dust Destruction ("Snowplow" Channeling): Rather than dragging an obscuring dust cloud, Siwarha's hot shock wave and radiation field vaporize or clear fragile alumina and silicate dust grains along its orbital path.[2][12] Because circumstellar dust preferentially scatters short-wavelength light, carving a dust-depleted channel through the envelope modulates blue-band light transmission

over the 2,170-day orbit.[12]

Implications for Common Envelope Evolution and the 2027 Elongation

The confirmation of Siwarha reshapes models of terminal massive stellar evolution and binary-driven mass loss.[1] Acting as a gravitational and hydrodynamic paddle inside the outer envelope, the companion injects kinetic energy and angular momentum into Betelgeuse's convective layers. During the 2019-2020 Great Dimming, orbital reconstruction indicates that wake turbulence and tidal perturbation from Siwarha helped destabilize a surface convective plume, precipitating the dust-forming mass ejection.[3][14] Moreover, because Siwarha experiences continuous hydrodynamic and gravitational drag inside the chromosphere, its orbit is decaying; within roughly 10,000 years, the companion will plunge into Betelgeuse's envelope, initiating a Common Envelope merger that will accelerate mass stripping prior to core-collapse supernova.[14]

Conclusion: Decoding the Ghost in Orion

By uniting radial velocity phase analysis with Hubble STIS detections of a trailing Fe II and Mg II plasma wake, astronomers have demonstrated that Betelgeuse's century-old 2,170-day Long Secondary Period is the orbital signature of its hidden companion, Siwarha (Ori B).[8][11] This finding validates the binary origin of Wood Sequence LSPs across evolved red giants while providing a nearby laboratory for pre-common-envelope hydrodynamics.[5][9] With orbital ephemerides predicting that Siwarha will reach its next maximum angular elongation from Betelgeuse in August 2027, upcoming high-contrast coronagraphic and interferometric campaigns will have an optimal window to directly characterize this companion surfing the atmosphere of Orion's supergiant.[6][8]

Updates... (Nov 22, 2026): Enriched all 15 citations -- including NASA Science, Simons Foundation, ScienceAlert, NOIRLab, Physics World, AAS Nova, CMU, arXiv, and primary ApJ/Nature/MNRAS DOIs -- with exact verified publication dates.

References

* [1] NASA Science (Jan. 5, 2026; updated Jan. 6, 2026). NASA Hubble Helps Detect 'Wake' of Betelgeuse's Elusive Companion Star. -- https://science.nasa.gov/missions/hubble/nasa-hubble-helps-detect-wake-of-betelgeuses-elusive-companion-star/

* [2] Simons Foundation (Oct. 21, 2024). Betelgeuse Betelgeuse? Bright Star Betelgeuse Likely Has a 'Betelbuddy' Stellar Companion. -- https://www.simonsfoundation.org/2024/10/21/betelgeuse-betelgeuse-bright-star-betelgeuse-likely-has-a-betelbuddy-stellar-compani

* [3] Montarg s, M., Cannon, E., Lagadec, E., et al. (June 16, 2021). A dusty veil shading Betelgeuse during its Great Dimming. Nature, 594(7863), 365-368. -- https://doi.org/10.1038/s41586-021-03546-8

* [4] ScienceAlert (Jan. 7, 2026). Betelgeuse Is Definitely Not Alone, 8-Year Study Confirms. -- https://www.sciencealert.com/betelgeuse-is-definitely-not-alone-8-year-study-confirms

* [5] Soszy ski, I., et al. (Apr. 6, 2022). Eppur si muove: On the Origin of Long Secondary Periods in Red Giant Stars (arXiv:2204.02832). --

https://arxiv.org/abs/2204.02832

* [6] NOIRLab (July 21, 2025). Gemini North Discovers Long-Predicted Stellar Companion of Betelgeuse. -- https://noirlab.edu/public/news/noirlab2523/

* [7] Physics World (Nov. 6, 2024). 'Buddy star' could explain Betelgeuse's varying brightness. -- https://physicsworld.com/a/buddy-star-could-explain-betelgeuses-varying-brightness/

* [8] Dupree, A. K., Cristofari, P. I., MacLeod, M., et al. (Feb. 3, 2026). Betelgeuse: Detection of the Expanding Wake of the Companion Star. The Astrophysical Journal, 998, 50. -- https://doi.org/10.3847/1538-4357/ae2ed5

* [9] AAS Nova (Apr. 26, 2021). Solving the Mystery of Varying Red Giants. -- https://aasnova.org/2021/04/26/solving-the-mystery-of-varying-red-giants/

* [10] Kiss, L. L., Szab , G. M., & Bedding, T. R. (Nov. 11, 2006). Variability in red supergiant stars: pulsations, long secondary periods and convection noise. Monthly Notices of the Royal Astronomical Society, 372(4), 1721-1734. -- https://doi.org/10.1111/j.1365-2966.2006.10973.x

* [11] Goldberg, J. A., Joyce, M., & Moln r, L. (Nov. 29, 2024). A Buddy for Betelgeuse: Binarity as the Origin of the Long Secondary Period in Orionis. The Astrophysical Journal, 977, 35. -- https://doi.org/10.3847/1538-4357/ad87f4

* [12] AAS Nova (Dec. 10, 2024). Hiding in Plain Sight: Betelgeuse's Binary Buddy. -- https://aasnova.org/2024/12/10/hiding-in-plain-sight-betelgeuses-binary-buddy/

* [13] Carnegie Mellon University (Oct. 8, 2025). X-Ray Study Reveals New Details About Betelgeuse's Elusive Companion Star. -- https://www.cmu.edu/mcs/news-events/2025/1008-betelgeuse-chandra

* [14] MacLeod, M., et al. (Jan. 5, 2026). The Betelgeuse Enigma: The Betelbuddy Hypothesis (arXiv:2601.02012). -- https://arxiv.org/abs/2601.02012

* [15] NASA Science (Jan. 5, 2026; updated Jan. 6, 2026). Betelgeuse: Effect of Companion Star Wake. -- https://science.nasa.gov/asset/hubble/betelgeuse-effect-of-companion-star-wake/