Ecology & Evolution | Jan 8, 2026 | 7 min read | By Bryan White

The Northern Cradle: Re-evaluating the Birthplace of Modern Hominins

Evolution * Paleontology * Archaeology

Introduction: The "Muddle in the Middle" and the Pan-African Shift

The paleoanthropological record of human evolution across the Middle Pleistocene -- spanning roughly 780,000 to 125,000 years ago -- has long been characterized as the "muddle in the middle" due to a scarcity of securely dated fossils bridging archaic Homo erectus and the derived lineages of Homo sapiens and Homo neanderthalensis. In January 2026, a landmark study published in Nature describing hominin fossils from the Thomas Quarry I complex in Casablanca, Morocco, resolved a critical segment of this transition.[1]

For decades, the "East Side Story" hypothesis treated the East African Rift Valley as the primary evolutionary engine of the genus Homo, relegating North Africa to a peripheral cul-de-sac. While the 2017 redating of the Jebel Irhoud crania to 315,000 years ago demonstrated that early Homo sapiens inhabited the Maghreb far earlier than previously assumed, the deeper evolutionary roots of those North African populations remained unknown.[3] Dated with high-resolution magnetostratigraphy to 773,000 +/- 4,000 years ago, the newly analyzed Thomas Quarry I specimens (recovered from the Grotte Hominid s) reveal a mosaic population standing near the base of the lineage leading to modern humans.[1] Combining archaic mandibular robustness with derived dental morphology, these fossils establish an African sister group to European Homo antecessor and substantiate a Pan-African model of human origins.[1][2]

Stratigraphy and Magnetostratigraphic Chronometry of Casablanca

The preservation of the Casablanca hominin sequence stems from the Quaternary coastal geology of the Moroccan Atlantic littoral.[2] Driven by Milankovitch orbital forcing, Pleistocene glacial-interglacial oscillations repeatedly shifted global sea levels. During interglacial periods when polar ice sheets melted, marine high-stands carved sea caves and wave-cut platforms into coastal bedrock; during glacial regressions, prevailing winds swept exposed shelf sands inland into carbonate-cemented eolianite dune ridges.[2] Coupled with gradual tectonic uplift, this cycle constructed a stepped sequence of marine and continental formations, including the Oulad Hamida Formation that hosts the Grotte Hominid s at Thomas Quarry I.[1][2]

Following the initial discovery of a partial mandible at the quarry by amateur collector Philippe Beriro in 1969, systematic excavations under the Franco-Moroccan Program "Pr histoire de Casablanca" -- co-directed by Jean-Paul Raynal and Abderrahim Mohib -- transformed the coastal quarries into a high-resolution paleoenvironmental archive.[1][2] During the 2008-2009 field seasons, researchers recovered a critical suite of adult and juvenile mandibles, isolated dentition, and post-cranial vertebrae from the cemented calcarenite matrix of the cave.[1][2]

To establish an exact chronological anchor, geologists applied high-density magnetostratigraphy to hundreds of oriented sediment cubes across the undisturbed cave infillings. Because Earth's dipole field shields the surface from solar radiation while periodically reversing polarity, ferromagnetic minerals such as magnetite lock in the prevailing planetary magnetic orientation during lithification. The fossil-bearing stratum at the Grotte Hominid s directly straddles the global reversal between the reversed-polarity Matuyama Chron (which began ~2.58 million years ago) and the normal-polarity Brunhes Chron.[1][5] Pinpointing the Matuyama-Brunhes Boundary (MBB) directly within the hominin horizon dates the assemblage to 773,000 +/- 4,000 years ago -- an extraordinary level of chronometric precision that permits direct synchronization with contemporaneous European sites such as Gran Dolina at Atapuerca.[1][2]

Mosaic Morphology: Archaic Mandibles and Derived Dentition

Led by Jean-Jacques Hublin at the Max Planck Institute for Evolutionary Anthropology, comparative morphometric analysis of the Thomas Quarry I hominins -- including adult mandibles ThI-GH-1 and ThI-GH-10717, juvenile mandible ThI-GH-10978, and eight associated cervical and thoracic vertebrae -- revealed a pronounced mosaic evolutionary pattern.[1][2]

* Mandibular and Axial Architecture: The mandibles retain thick, heavily buttressed corpora designed for high masticatory loads and completely lack a projecting mental eminence (true chin), an autapomorphy restricted to Homo sapiens.[1] Similarly, the small cervical and thoracic vertebrae associated with ThI-GH-10717 resemble the archaic post-cranial proportions of Homo erectus.[1][2] Yet the dental arcade already exhibits a derived parabolic curvature rather than the primitive U-shaped geometry of early Homo.[1]

* Enamel-Dentine Junction (EDJ) Signals: Using micro-computed tomography (micro-CT) to reconstruct the genetically conservative Enamel-Dentine Junction, researchers identified crown proportions and cusp topologies shared with Homo sapiens and Neanderthals, clearly diverging from Homo erectus.[1][2] Crucially, the Thomas Quarry teeth lack specialized Neanderthal apomorphies such as molar taurodontism and continuous mid-trigonid crests, positioning the population near the root of the Homo sapiens lineage prior to Neanderthal divergence.[1][2]

| Anatomical Trait | | Thomas Quarry I (773 ka) | | Homo antecessor (800 ka) | | Homo erectus (~1.0 Ma) | | Homo sapiens (Jebel Irhoud, 315 ka) |

| Mandibular Corpus | | High Robustness | | Moderate to High | | Very High Robustness | | Moderate |

| Mental Eminence (Chin) | | Absent | | Absent | | Absent | | Incipient / Variable |

| Dental Arcade Shape | | Parabolic (Derived) | | Parabolic | | U-Shaped / Sub-rectangular | | Parabolic |

| Molar Taurodontism | | Absent | | Absent | | Absent | | Absent |

| Mid-Trigonid Crest | | Absent | | Variable / Incipient | | Absent | | Absent |

| Vertebral Morphology | | Small / Erectus-like | | Modern-like | | Archaic / Robust | | Gracile / Modern |

Acheulean Techno-Economics and Green Sahara Paleoecology

The Thomas Quarry I sequence documents over half a million years of cultural and behavioral continuity within the North African Acheulean technocomplex, beginning with 1.3-million-year-old quartzite Large Cutting Tools (LCTs) in lower Unit L1 and continuing through refined bifacial handaxes, picks, and transverse-edged cleavers in the 773,000-year-old hominin horizon.[6][7] Detailed techno-economic reconstructions confirm structured raw-material selection of coastal quartzite cobbles and multi-stage bifacial flake reduction geared toward heavy megafaunal carcass disarticulation and butchery.[7]

Paleoecological evidence demonstrates that these hominins occupied a humid savanna-woodland ecosystem during periodic "Green Sahara" monsoon intervals.[1] Faunal remains from the Grotte Hominid s include Elephas, hippopotamuses, rhinoceroses, equids, gazelles, and the giant East African grazing gelada Theropithecus oswaldi -- confirming active trans-Saharan biogeographic corridors.[4][8] Taphonomic inspection of a hominin femoral shaft bearing hyena tooth marks further indicates intense competition with large cave carnivores for shelter and prey.[9]

| Site | | Location | | Chronology | | Key Assemblage | | Evolutionary Significance |

| Ain Hanech | | Algeria | | ~1.8 Ma | | Oldowan Lithics | | Earliest documented hominin presence in North Africa. |

| Thomas Quarry I (Unit L1) | | Morocco | | 1.3 Ma | | Acheulean LCTs | | Earliest Acheulean bifacial technology in the Maghreb. |

| Tighennif (Ternifine) | | Algeria | | ~1.0 Ma | | H. erectus Mandibles | | Classic North African early Pleistocene Homo erectus. |

| Thomas Quarry I (GH) | | Morocco | | 773 ka | | Mandibles, Teeth, Vertebrae | | Basal to H. sapiens; African sister lineage to H. antecessor. |

| Jebel Irhoud | | Morocco | | 315 ka | | Crania, Mandibles, Lithics | | Earliest known pan-African Homo sapiens fossils. |

Conclusion: A Braided Stream Across the Maghreb

Because ancient genomic molecular clocks place the evolutionary divergence between the modern human lineage and the Neanderthal-Denisovan clade between 550,000 and 765,000 years ago, the 773,000-year-old Thomas Quarry I fossils sample the exact biological horizon immediately preceding this split.[1][2] For years, hypotheses proposing Spain's 850,000-year-old Homo antecessor as the Last Common Ancestor struggled to explain why the core trunk leading to Homo sapiens subsequently evolved in Africa. By identifying the Thomas Quarry population as an African sister group to Homo antecessor, the Casablanca discovery resolves this biogeographic tension.[1][2]

While the European branch diverged north of the Mediterranean toward the specialized Neanderthal lineage, the North African population persisted across periodic Green Sahara corridors, evolving over the subsequent 450,000 years into the early Homo sapiens documented at Jebel Irhoud.[1][3] Ultimately, the Thomas Quarry I hominins

dismantle linear single-origin models: modern humanity emerged from a braided, continent-wide Pan-African evolutionary network in which the coastal savannas of the Maghreb were as generative as the rift valleys of East Africa.[1][2]

Updates... (Nov 24, 2026): Replaced two HTTP 403/anti-bot news URLs (Morocco World News and The Guardian) by consolidating redundant commentary onto the primary Nature and Max Planck Institute sources and substituting the CrossRef-verified primary chronostratigraphy study for 1.3 Ma Unit L1 Acheulean tools (Gallotti et al., Jul 28, 2021; Scientific Reports, 11(1), 15340; DOI 10.1038/s41598-021-94695-3); renumbered references [1]-[9] cleanly; and consolidated prior Nov 21, 2026 publication-date updates.

References & Primary Sources

* [1] Hublin, J. J., et al. (2026, Jan 7). Early hominins from Morocco basal to the Homo sapiens lineage. Nature. -- https://doi.org/10.1038/s41586-025-09914-y

* [2] Max Planck Institute for Evolutionary Anthropology (2026, Jan 7). Early hominins from Thomas Quarry I (Morocco) reveal an African lineage near the root of Homo sapiens. -- https://www.mpg.de/25951868/0105-evan-early-hominins-from-thomas-quarry-i-morocco-150495-x

* [3] Hublin, J. J., et al. (2017, Jun 8). New fossils from Jebel Irhoud, Morocco and the pan-African origin of Homo sapiens. Nature. -- https://doi.org/10.1038/nature22336

* [4] National Geographic (2026, Jan 7). This ancient human may be the root of the Homo sapiens family tree. -- https://www.nationalgeographic.com/science/article/ancient-human-origins-fossils-morocco

* [5] Channell, J. E. T., Hodell, D. A., Singer, B. S., & Xuan, C. (2010, Dec). Reconciling astrochronological and 40Ar/39Ar ages for the Matuyama-Brunhes boundary and late Matuyama Chron. Geochemistry, Geophysics, Geosystems. -- https://doi.org/10.1029/2010gc003203

* [6] Gallotti, R., Muttoni, G., Lef vre, D., Degeai, J.-P., et al. (2021, Jul 28). First high resolution chronostratigraphy for the early North African Acheulean at Casablanca (Morocco). Scientific Reports, 11(1), 15340. -- https://doi.org/10.1038/s41598-021-94695-3

* [7] Gallotti, R., Raynal, J.-P., & Mohib, A. (2023, Oct 20). Early North African Acheulean techno-economic systems at Thomas Quarry I - L1 (Casablanca, Morocco). Journal of Anthropological Sciences (PMID 36587282). -- https://pubmed.ncbi.nlm.nih.gov/36587282/

* [8] Geraads, D., et al. (2010, Mar). Mammalian faunas from the Pliocene and Pleistocene of Casablanca (Morocco). Historical Biology. -- https://doi.org/10.1080/08912960903458011

* [9] Daujeard, C., Geraads, D., Gallotti, R., et al. (2016, Apr 27). Pleistocene Hominins as a Resource for Carnivores: A c. 500,000-Year-Old Human Femur Bearing Tooth-Marks in North Africa (Thomas Quarry I, Morocco). PLOS One. -- https://doi.org/10.1371/journal.pone.0152284