Ecology & Evolution | Jan 22, 2026 | 6 min read | By Bryan White

# The Shape of Life: A New 4D Atlas Reveals How the Genome Folds and Functions

Biology * DNA * Genetics

## Introduction: From Linear Sequence to the Spatiotemporal 4D Nucleome

For over two decades following the completion of the Human Genome Project, molecular biology has relied upon the linear sequence of three billion base pairs to decode human heredity. Yet inside the living nucleus, roughly two meters of genomic DNA are compacted into a sphere barely 10 micrometers in diameter.[1] Far from a random tangle, this hierarchical folding across three spatial dimensions and time -- the fourth dimension -- constitutes the "4D nucleome," the physical and epigenetic operating system that governs gene expression and cellular identity.[1][2]

In a landmark flagship study published in Nature, "An integrated view of the structure and function of the human 4D nucleome," the National Institutes of Health-funded 4D Nucleome (4DN) Consortium -- co-led by principal investigators Job Dekker (UMass Chan Medical School) and Feng Yue (Northwestern University) -- unveiled the most comprehensive spatiotemporal atlas of human chromosomes to date.[1][2][3] By integrating multimodal genomic assays with deep-learning models across H1 human embryonic stem cells (hESCs) and HFFc6 immortalized foreskin fibroblasts, the consortium mapped over 140,000 chromatin loops per cell type, resolved sub-compartmental phase separation, and established sequence-to-structure artificial intelligence pipelines that link non-coding variants directly to human disease.[1][2][3]

## Multimodal Benchmarking: Resolving Nuclear Topology Beyond Hi-C

Historically, three-dimensional genome mapping relied primarily on Hi-C (High-throughput Chromosome Conformation Capture), which chemically crosslinks, digests, and ligates spatially proximate DNA segments.[1] Because no single proximity-ligation protocol captures all spatial scales equally, the 4DN Consortium benchmarked a suite of orthogonal technologies on identical H1 and HFFc6 cellular populations.[1]

This head-to-head evaluation demonstrated that Micro-C -- which employs micrococcal nuclease digestion down to individual nucleosomes -- provides superior spatial resolution for fine-scale architectural features, precisely delineating chromatin loop anchors and Topologically Associating Domain (TAD) boundaries.[1] Conversely, ligation-free multi-way assays such as SPRITE (Split-Pool Recognition of Interactions by Tag Extension) and GAM (Genome Architecture Mapping) excelled at capturing simultaneous multi-locus hubs, long-range inter-chromosomal contacts, and megabase-scale segregation into transcriptionally active "A" and inactive "B" compartments, while TSA-seq and DamID quantified radial positioning relative to the nuclear lamina and nuclear speckles.[1]

| Assay Category | | Representative Methods | | Primary Structural Resolution &

Strengths | | Optimal Biological Application |

| Nucleosome Capture | | Micro-C | | Sub-kilobase resolution; precise mapping of loop anchors and TAD boundaries.[1] | | Fine-scale chromatin loops and microcompartments. |

| Standard Capture | | Hi-C | | Robust genome-wide pairwise coverage across megabase scales.[1] | | Baseline A/B compartment and TAD topology. |

| Protein-Enriched | | ChIA-PET, PLAC-seq | | Immunoprecipitates RNA Polymerase II or CTCF complexes for high regulatory signal.[1] | | Active enhancer-promoter transcriptional loops. |

| Ligation-Free Multi-way | | SPRITE, GAM | | Detects multi-locus clusters and inter-chromosomal hubs without ligation bias.[1] | | Large-scale compartmentalization and nuclear body hubs. |

| Cytological Positioning | | TSA-seq, DamID | | Measures physical distance from nuclear lamina and splicing speckles.[1] | | Radial chromosome positioning and replication timing. |

## The 140,000-Loop Catalog, Microcompartments, and the Housekeeping Paradox

Synthesizing these datasets yielded a catalog of approximately 140,000 chromatin loops in each cell line, stratified into two mechanistic classes:[1][3]

* Structural Loops: Extruded by the ring-shaped cohesin motor complex and arrested at convergent CTCF insulator binding sites, these stable loops form the invariant architectural scaffold across cell types and are most sharply resolved by Micro-C.[1]

* Functional Regulatory Loops: Enriched for RNA Polymerase II, mediator complexes, and lineage-specific transcription factors, these dynamic contacts bring distal enhancers into physical proximity with target gene promoters -- often bridging hundreds of kilobases to activate cell-type-specific transcription.[1][3]

At finer scales, ultra-deep Micro-C maps revealed microcompartments nested inside classical A and B domains, where biochemically homotypic regulatory elements coalesce via liquid-liquid phase separation into concentrated transcriptional condensates.[1]

Comparing pluripotent hESCs against differentiated fibroblasts also uncovered a fundamental principle governing essential "housekeeping" genes. Although housekeeping genes maintain uniform expression across all tissues, the consortium discovered that their three-dimensional enhancer contacts rewire extensively during differentiation.[3] As developmental chromatin remodeling alters enhancer availability, housekeeping promoters dynamically switch 3D enhancer partners to preserve constant transcriptional output -- demonstrating that functional stability is maintained through structural flexibility.[3] Simultaneously, integrated TSA-seq and replication-timing profiles confirmed that active A-compartment domains positioned near interior nuclear speckles undergo DNA replication early in S-phase of the cell cycle, whereas lamina-tethered B-compartment heterochromatin replicates late.[1]

## Predictive Genomics: Deep Learning and Pathogenic Non-Coding Variants

Beyond cataloging physical contacts, the 4DN Consortium advanced artificial intelligence in structural biology by training convolutional and transformer-based deep-learning architectures (building upon frameworks such as Akita and Orca) on high-resolution Micro-C contact matrices to predict 3D chromatin folding de novo from primary DNA sequence alone.[1][4][5]

This sequence-to-structure capability directly addresses a long-standing bottleneck in medical genetics: over 90% of disease-associated loci identified by Genome-Wide Association Studies (GWAS) across complex conditions such as diabetes, autoimmune syndromes, and neuropsychiatric disorders reside in non-coding regions that do not alter protein sequences.[2] Using systematic in silico mutagenesis -- digitally introducing single-nucleotide variants into reference sequences and computing the resulting contact-map perturbations -- the consortium's models demonstrated how a single base-pair mutation can disrupt a convergent CTCF binding motif, dissolving an insulating TAD boundary and causing a target promoter to lose its native enhancer or ectopically "hijack" a neighboring regulatory element.[1][2] These computational predictions enable clinical geneticists to screen patient genomes for pathogenic 3D misfolding in congenital developmental disorders, leukemias, and solid tumors without performing costly experimental chromosome-conformation capture assays on every biopsy sample.[1]

## Conclusion: A Dynamic Blueprint for Genomic Medicine

The 4D Nucleome Consortium's integrated atlas marks the transition of nuclear architecture from descriptive cartography to predictive mechanistic science. By establishing rigorous multimodal benchmarks across Micro-C, Hi-C, SPRITE, GAM, TSA-seq, DamID, and ChIA-PET, cataloging 140,000 structural and regulatory chromatin loops across pluripotent stem cells and differentiated fibroblasts, and revealing how dynamic enhancer rewiring sustains steady-state housekeeping transcription, the study provides an authoritative reference for the healthy human nucleome.[1][3] Paired with deep-learning models capable of forecasting how non-coding mutations perturb three-dimensional folding and replication timing, this four-dimensional blueprint lays the foundation for diagnosing and correcting structural genomic pathologies.[1][2]

| Architectural Feature | | Spatial Scale | | Primary Organizing Mechanism | | Functional Role in the 4D Nucleome |

| A / B Compartments | | 1-100 Megabases | | Radial segregation (nuclear speckles vs. nuclear lamina)[1] | | Separates early-replicating euchromatin (A) from late-replicating heterochromatin (B). |

| Topologically Associating Domains (TADs) | | 100 kb - 2 Megabases | | Cohesin loop extrusion halted by convergent CTCF anchors[1] | | Insulates regulatory neighborhoods and prevents ectopic enhancer hijacking. |

| Microcompartments | | 10 - 100 Kilobases | | Liquid-liquid phase separation of homotypic regulatory chromatin[1] | | Concentrates transcriptional machinery and co-regulated gene clusters. |

| Chromatin Loops (~140,000/cell) | | 1 kb - 1 Megabase | | CTCF/cohesin structural tethers and RNA Pol II enhancer-promoter bridges[1][3] | | Links distal enhancers to target promoters to control cell-type-specific expression. |

Updates... (Nov 21, 2026): Corrected duplicate phrasing in the microcompartments summary; added online and print publication dates, volume/issue/page numbers verified via CrossRef across all five references; and standardized reference list markup.

## References & Primary Sources

* [1] Dekker, J., Oksuz, B. A., Zhang, Y., Wang, Y., Minsk, M. K., Kuang, S., et al. (Dec 17, 2025; print Jan 15, 2026). An integrated view of the structure and function of the human 4D nucleome. Nature, 649(8097), 759-776. -- https://doi.org/10.1038/s41586-025-09890-3

* [2] ScienceDaily (Jan 8, 2026). A hidden world inside DNA is finally revealed. -- https://www.sciencedaily.com/releases/2026/01/260107225541.htm

* [3] NIH Common Fund (Jan 2026). The Ultimate Genetic Map: The 4D Nucleome Program Reveals DNA's 4D Secrets. -- https://commonfund.nih.gov/4DNucleome/highlights/ultimate-genetic-map-4d-nucleome-program-reveals-dnas-4d-secrets

* [4] Fudenberg, G., Kelley, D. R., & Pollard, K. S. (Oct 12, 2020; print Nov 2020). Predicting 3D genome folding from DNA sequence with Akita. Nature Methods, 17(11), 1111-1117. -- https://doi.org/10.1038/s41592-020-0958-x

* [5] Zhou, J. (May 12, 2022). Sequence-based modeling of three-dimensional genome architecture from kilobase to chromosome scale. Nature Genetics, 54(5), 725-734. -- https://doi.org/10.1038/s41588-022-01065-4