Climate & Earth Systems | Nov 30, 2025 | 9 min read | By Bryan White

The Silent Shift: How the World's Most Stable Rainforest is Changing

Environment * Climate Change * Conservation

Introduction: The Planetary Lungs in the Anthropocene Rainforest

For nearly half a century, mature pantropical forests have operated as a vital net carbon sink, absorbing roughly 30% of anthropogenic carbon dioxide (CO2) emissions annually.[1] Within the global carbon cycle, African tropical forests -- centered on the Congo Basin -- were long regarded as the most stable of Earth's three major rainforest massifs, buffered against the drought-induced mortality afflicting the Amazon and the industrial plantation conversion across Southeast Asia.[3] While CO2 fertilization historically stimulated higher rates of photosynthesis and woody biomass accumulation across intact central African stands, a landmark late-2025 study in Scientific Reports led by Dr. Pedro Rodr guez-Veiga, Dr. Nezha Acil, and Professor Heiko Balzter at the National Centre for Earth Observation (NCEO) demonstrates that Africa's forests crossed a critical threshold around 2010.[4][8]

Titled "Loss of tropical moist broadleaf forest has turned Africa's forests from a carbon sink into a source," the research reveals that anthropogenic deforestation and canopy degradation have overwhelmed physiological carbon uptake, driving a net continental biomass loss of approximately 106 billion kilograms (106 Tg) annually between 2010 and 2017.[4] This transition signifies that feedback loops linking land-use pressure, atmospheric drying, and ecosystem respiration are accelerating beyond Earth System Model (ESM) projections, aligning Africa with the Amazon and Southeast Asia as a net contributor of atmospheric greenhouse gases.[6][7][4]

Methodological Paradigm Shift: L-Band SAR, LiDAR, and Machine Learning

Quantifying the continental carbon budget of Africa was historically limited by sparse ground plots across the Congo Basin and persistent cloud cover in the Intertropical Convergence Zone (ITCZ).[3][10][11] To achieve a methodological evolution in biomass tracking, Rodr guez-Veiga, Acil, Balzter, and colleagues integrated multi-sensor radar, LiDAR, and machine learning at 100-meter resolution across 2007-2017:[4][9]

* L-Band Synthetic Aperture Radar (SAR): Active microwave pulses (~23 cm wavelength) from ALOS PALSAR penetrated cloud cover and upper foliage to measure backscatter from woody trunks and branches.[11]

* LiDAR Calibration & Random Forests: Radar signals were calibrated against airborne LiDAR canopy profiles and validated with field plots via random forest regressors, detecting both clear-cutting and sub-canopy degradation from selective logging and charcoal harvesting.[4][9][13]

Quantifying the 2010 Reversal and Regional Hemorrhage

The aboveground biomass (AGB) time series revealed a sharp decadal inflection. From 2007 to 2010, African forests functioned as a net sink, gaining +439 +/- 66 Tg/yr.[12] Post-2010, extraction outpaced regeneration, driving net losses of 132 +/- 20 Tg/yr (2010-2015) and 41 +/- 6 Tg/yr (2015-2017) -- averaging ~106 Tg/yr.[6][12] Losses centered in three moist broadleaf hotspots: the Democratic Republic of Congo (DRC) (slash-and-burn farming, artisanal logging, and urban charcoal demand),[4] Madagascar (agricultural clearance and fuelwood extraction),[6] and West Africa's Upper Guinean Forests (cocoa and rubber expansion).[4] Although CO2 fertilization promoted shrub encroachment in savannas, dryland gains offset only a fraction of dense rainforest losses.[4][14]

| Assessment Period | | Net Biomass Flux (Tg/yr) | | Carbon Balance Status | | Dominant Biophysical / Anthropogenic Driver |

| 2007 - 2010 | | +439 +/- 66 | | Net Sink | | Intact forest growth and CO2 fertilization exceed extraction.[12] |

| 2010 - 2015 | | 132 +/- 20 | | Net Source | | Accelerated clearing in DRC, Madagascar, and West Africa.[12] |

| 2015 - 2017 | | 41 +/- 6 | | Net Source | | Continued canopy fragmentation and El Ni o fire encroachment.[12][22] |

| Post-2010 Mean (2010-2017) | | ~ 106 | | Net Source | | Moist broadleaf loss overwhelms savanna shrub thickening.[6][4] |

Pan-Tropical Mechanics, Vapor Pressure Deficit, and the Cuvette Centrale Peatlands

Africa's transition completes a pan-tropical decline: intact tropical forests, which removed ~17% of anthropogenic CO2 in the 1990s, absorbed just 6% by the 2010s.[5][7] Whereas Southeast Asia flipped via industrial oil-palm conversion and peat drainage[2][18] and the southeastern Amazon flipped through agribusiness clearing compounded by climate change and thermal drought mortality,[15][16][17] Africa's source flux is currently dominated by small-scale land-use change.[4]

| Tropical Basin | | Current Flux Status | | Primary Driver of Source Flux | | Structural Vulnerability |

| Southeast Asia | | Net Source | | Industrial plantations (palm oil/pulp) & peat oxidation | | High legacy emissions from drained coastal peatlands.[18] |

| Amazon Basin | | Net Source / Neutral | | Cattle/soy conversion & drought-induced tree mortality | | Hydrological recycling breakdown & savanna dieback risk.[16][17] |

| Congo / African Tropics | | Net Source (Post-2010) | | Smallholder agriculture, charcoal, mining & logging | | Edge desiccation, rising VPD, and Cuvette Centrale peat exposure.[6][27] |

Physiologically, African canopies face rising Vapor Pressure Deficit (VPD) -- the exponential increase in atmospheric drying power under warming.[19][20] High VPD

forces trees to close leaf stomata to prevent xylem cavitation (hydraulic failure), which simultaneously halts CO2 uptake and induces carbon starvation.[20][21] Fragmentation also dries forest edges, enabling savanna wildfires to penetrate moist stands during El Ni o anomalies.[19][22][23]

Crucially, canopy loss threatens the regional hydrology sustaining the Cuvette Centrale peatlands -- a 145,500-167,600 km subterranean swamp complex in the central Congo Basin storing 29 to 30 gigatons (Gt) of carbon (equivalent to three years of global fossil fuel emissions).[24][25][26] Radiocarbon re-dating in 2024-2025 established that these peatlands began accumulating over 40,000 years ago.[27] If regional rainfall declines due to surrounding forest loss or if overlapping hydrocarbon concessions construct drainage canals, oxidation of this irrecoverable carbon reservoir would trigger catastrophic emissions.[14][27][28]

Policy Architecture: The Tropical Forests Forever Facility (TFFF)

The failure of project-based REDD+ offsets catalyzed the launch of the Tropical Forests Forever Facility (TFFF) at COP30 in Bel m.[29][30] Structured as a $125 billion sovereign endowment ($25 billion public sponsor capital leveraging $100 billion in bonds), the TFFF uses investment returns to pay tropical nations ~$4 per hectare annually for standing forest, with automatic deforestation penalties and a mandatory 20% direct allocation to Indigenous Peoples and Local Communities (IPLCs).[30][31][32] With initial COP30 pledges reaching $6.7 billion, full capitalization remains urgent.[31]

Conclusion: Stanching the Anthropocene Hemorrhage

The NCEO findings confirm that intact tropical ecosystems can no longer passively buffer industrial emissions.[4][6] Yet because Africa's 106 Tg/yr biomass deficit is driven primarily by land-use pressures rather than irreversible climatic dieback, policy intervention retains agency.[4] Fully capitalizing the TFFF, expanding clean alternatives to urban charcoal, and legally safeguarding the Cuvette Centrale peatlands are essential to restore the Congo Basin's carbon shield.[27][31]

Updates... (Nov 24, 2026): Corrected the lead author's first name from Pablo to Dr. Pedro Rodr guez-Veiga (Scientific Reports, 15, 41744), removed an erroneous gamete-fertilization Wiki link on CO2 fertilization, added standardized id="references-list" and id="ref-1"-id="ref-33" reference parity with outbound link attributes, and linked Reference [16] directly to its CrossRef DOI. (Prior Nov 22, 2026 update documented late-2026 TFFF sovereign pledges reaching $7.8 billion across 66 countries and the EU, plus $88 million in COP31 Antalya philanthropic pledges on Nov 11, 2026[33].)

References & Primary Sources

* [1] Harris, N., & Rose, M. (Jul 24, 2025). World's Forest Carbon Sink Shrank to its Lowest Point in at Least 2 Decades, Due to Fires and Persistent Deforestation. World Resources Institute. -- https://www.wri.org/insights/forest-carbon-sink-shrinking-fires-deforestation

* [2] Harris, N., & Gibbs, D. (Jan 21, 2021). Forests Absorb Twice As Much Carbon As They Emit Each Year. World Resources Institute. --

https://www.wri.org/insights/forests-absorb-twice-much-carbon-they-emit-each-year

* [3] Grantham Research Institute on Climate Change and the Environment, LSE (Nov 17, 2025). Congo Basin rainforest at critical point and requires sustained investment from the international community. -- https://www.lse.ac.uk/granthaminstitute/news/congo-basin-rainforest-at-critical-point-and-requires-sustained-investment-from-the-in

* [4] Rodr guez-Veiga, P., et al. (Nov 28, 2025). Loss of tropical moist broadleaf forest has turned Africa's forests from a carbon sink into a source. Scientific Reports, 15, 41744. -- https://doi.org/10.1038/s41598-025-27462-3

* [5] Hubau, W., et al. (Mar 4, 2020; print Mar 5, 2020). Asynchronous carbon sink saturation in African and Amazonian tropical forests. Nature, 579(7797), 80-87. -- https://doi.org/10.1038/s41586-020-2035-0

* [6] Watts, J. (Nov 28, 2025). Africa's forests transformed from carbon sink to carbon source, study finds. The Guardian. -- https://www.theguardian.com/environment/2025/nov/28/africa-forests-transformed-carbon-sink-carbon-source-study

* [7] Energy Innovation (Aug 7, 2021). Tipping Points In Tropical Forests: Carbon Fluxes In The Amazon And Africa. -- https://energyinnovation.org/expert-voice/tipping-points-in-tropical-forests-carbon-fluxes-in-the-amazon-and-africa/

* [8] Ruehr, S., Keenan, T. F., Williams, C., et al. (Jul 25, 2023). Evidence and attribution of the enhanced land carbon sink. Nature Reviews Earth & Environment, 4(8), 518-534. -- https://doi.org/10.1038/s43017-023-00456-3

* [9] Thompson, T. (Nov 28, 2025). Turning Data into Action: How Global Collaboration Revealed Africa's Carbon Shift. National Centre for Earth Observation (NCEO). -- https://www.nceo.ac.uk/news-media/turning-data-into-action-how-global-collaboration-revealed-africas-carbon-shift/

* [10] Fisher, J. B., et al. (Sep 5, 2013). African tropical rainforest net carbon dioxide fluxes in the twentieth century. Philosophical Transactions of the Royal Society B, 368(1625), 20120376. -- https://doi.org/10.1098/rstb.2012.0376

* [11] Bouvet, A., et al. (Mar 2018). An above-ground biomass map of African savannahs and woodlands at 25 m resolution derived from ALOS PALSAR. Remote Sensing of Environment, 206, 156-173. -- https://doi.org/10.1016/j.rse.2017.12.030

* [12] The New Indian Express (Nov 29, 2025). Are Africa's forests turning from a carbon sink into a carbon source? -- https://www.newindianexpress.com/world/2025/Nov/29/are-africas-forests-turning-from-a-carbon-sink-into-a-carbon-source

* [13] Rodr guez-Veiga, P., et al. (Jul 24, 2020). Carbon Stocks and Fluxes in Kenyan Forests and Wooded Grasslands Derived from Earth Observation and Model-Data Fusion. Remote Sensing, 12(15), 2380. -- https://doi.org/10.3390/rs12152380

* [14] Nitnaware, H. (Nov 28, 2025). Africa's forests transitioned from carbon sink to net carbon source. Down To Earth. -- https://www.downtoearth.org.in/amp/story/forests/africas-forests-transitioned-from-carbon-sink-to-net-carbon-source

* [15] Veit, P., Gibbs, D., & Reytar, K. (Jan 6, 2023). Indigenous Forests Are Some of the Amazon's Last Carbon Sinks. World Resources Institute. -- https://www.wri.org/insights/amazon-carbon-sink-indigenous-forests

* [16] Basso, L. S., et al. (Sep 1, 2023). Atmospheric CO2 inversion reveals the Amazon as a minor carbon source caused by fire emissions, with forest uptake offsetting about half of these emissions. Atmospheric Chemistry and Physics, 23, 9685-9711. -- https://doi.org/10.5194/acp-23-9685-2023

* [17] Gatti, L. V., et al. (Jul 14, 2021; print Jul 15, 2021). Amazonia as a carbon source linked to deforestation and climate change. Nature, 595(7867), 388-393. -- https://doi.org/10.1038/s41586-021-03629-6

* [18] Project Drawdown (Nov 7, 2025; updated Aug 7, 2026). Why forests and farming are critical climate solutions in Southeast Asia. -- https://drawdown.org/insights/why-forests-and-farming-are-critical-climate-solutions-in-southeast-asia

* [19] Brando, P. M., et al. (May 30, 2019). Droughts, Wildfires, and Forest Carbon Cycling: A Pantropical Synthesis. Annual Review of Earth and Planetary Sciences, 47(1), 555-581. -- https://doi.org/10.1146/annurev-earth-082517-010235

* [20] Grossiord, C., et al. (Mar 20, 2020; print Jun 2020). Plant responses to rising vapor pressure deficit. New Phytologist, 226(6), 1550-1566. -- https://doi.org/10.1111/nph.16485

* [21] McDowell, N., et al. (Feb 16, 2018; print Aug 2018). Drivers and mechanisms of tree mortality in moist tropical forests. New Phytologist, 219(3), 851-869. -- https://doi.org/10.1111/nph.15027

* [22] Lehsten, V., et al. (Mar 6, 2009). Estimating carbon emissions from African wildfires. Biogeosciences, 6(3), 349-360. -- https://doi.org/10.5194/bg-6-349-2009

* [23] Gatti, L. V., et al. (Nov 1, 2010). Vertical profiles of CO2 above eastern Amazonia suggest a net carbon flux to the atmosphere and balanced biosphere between 2000 and 2009. Tellus B, 62(5). -- https://doi.org/10.3402/tellusb.v62i5.16606

* [24] Hawthorne, D., et al. (Dec 2025). Two contrasting swamp forest succession pathways in central Congo Basin peatlands. Quaternary Science Reviews, 369, 109637. -- https://doi.org/10.1016/j.quascirev.2025.109637

* [25] Cavallito, M. (Aug 11, 2025). Congo peatlands are twice as old as previously believed. Re Soil Foundation. -- https://resoilfoundation.org/en/environment/congo-tropical-peatlands-40k-years/

* [26] Lambrou, J. P. (Jul 21, 2022). True size of world's largest tropical peatland revealed for the first time. NMBU. -- https://www.nmbu.no/en/research/true-size-worlds-largest-tropical-peatland-revealed-first-time

* [27] University of Leeds (Jul 29, 2025). World's largest tropical peatlands revealed to be more than 40,000 years old. -- https://environment.leeds.ac.uk/faculty/news/article/5985/world-s-largest-tropical-peatlands-revealed-to-be-more-than-40-000-years-

* [28] Ruiz, S. (Dec 2, 2024). Peatlands in the DRC: A "Carbon Treasure Trove". Woodwell Climate Research Center. -- https://www.woodwellclimate.org/protecting-drc-peatlands-sustainable-economic-development/

* [29] COP30 Brasil (Oct 28, 2026). Tropical Forest Forever Facility, launched at COP30, receives USD 500 million pledge from the Netherlands. -- https://cop30.br/en/news-about-cop30/tropical-forest-forever-facility-launched-at-cop30-receives-usd-500-million-pledge-from-the-n

* [30] Barber, C., Davey, E., Zalles, V., & Oliveira, M. (Nov 26, 2025). The Tropical Forests Forever Facility Could Finally Finance Nature Conservation. Will Funders Back It? World Resources Institute. -- https://www.wri.org/insights/financing-nature-conservation-tropical-forest-forever-facility

* [31] de Santi, A. (Nov 24, 2025). Brazil's forest fund faces a slow takeoff at COP30 despite initial support. Mongabay. -- https://news.mongabay.com/2025/11/brazils-forest-fund-faces-a-slow-takeoff-at-cop30-despite-initial-support/

* [32] Ruiz, S. (Oct 28, 2025). TFFF Unlocks Finance for Forests. Woodwell Climate Research Center. -- https://www.woodwellclimate.org/tfff-tropical-forest-finance/

* [33] WWF International (Nov 11, 2026). New philanthropic pledges signal strong confidence in the TFFF. -- https://wwf.panda.org/wwf_news/?16541391/New-philanthropic-pledges-signal-strong-confidence-in-the-TFFF