Climate & Earth Systems | Feb 3, 2026 | 10 min read | By Bryan White

The West's Vanishing Snowpack: What the 2026 "Warm Drought" Means for Fire Season

Environment * Climate Change * Public Health

Introduction: The Decoupling of Precipitation and Snowpack Storage

The winter of 2025-2026 has emerged as a defining case study in the climatological transformation of the western United States. Characterized by a pervasive "snow drought," the water year has defied conventional expectations. Historically, the Sierra Nevada, the Cascades, and the Rocky Mountains function as passive natural infrastructure, capturing Pacific storm moisture within the snowpack until spring.[1] Despite near- or above-average precipitation in several northern watersheds, the Western cordillera entered February 2026 with record-low snow water equivalent (SWE) -- a decoupling of moisture input and solid-phase storage that threatens regional water security.[2]

On January 4, 2026, NASA Moderate Resolution Imaging Spectroradiometer (MODIS) satellite telemetry recorded snow cover across just 141,416 square miles of the western United States -- the lowest extent for that date in the MODIS record dating back to 2001.[2] Ground-based USDA SNOTEL sensors confirmed that over 80% of stations across Washington, Oregon, Colorado, Utah, Arizona, and New Mexico had fallen below the 20th percentile for SWE, with numerous basins registering historic lows.[2][5]

Hydrologists distinguish between two primary snow drought regimes:[1][6]

* Dry Snow Drought: Driven by blocking high-pressure systems that deflect storm tracks, starving mountains of total precipitation (predominant across the Desert Southwest and Southern Rockies in early 2026).[6]

* Warm Snow Drought: Near- or above-average precipitation accompanied by elevated freezing levels that partition moisture into liquid rain and trigger mid-winter melt.[3][6]

Water Year 2026 combines both regimes. In the Pacific Northwest and Northern Sierra, vigorous atmospheric river sequences delivered abundant subtropical moisture, yet warm wet-bulb temperatures forced rain-on-snow events at mid-altitude elevations, generating winter floodflows while depleting summer storage.[2][3][4]

Synoptic Architecture: Non-Canonical La Ni a and Persistent Ridging

The meteorological anomalies of 2025-2026 stemmed from planetary wave blocking, a modified El Ni o-Southern Oscillation (ENSO) teleconnection, and anthropogenic radiative forcing. Entering winter under a NOAA "La Ni a Advisory" (Ni o-3.4 sea surface temperatures near 0.5 deg C, with a 75% probability of transitioning to ENSO-neutral by spring), canonical models favored cold, snow-bearing storms in the Pacific Northwest.[7][8][9]

Instead, persistent upper-level ridging -- specifically a recurring "South-ridge"

centered over the Southwest -- dominated the synoptic pattern.[10][11] Zonal subtropical atmospheric rivers undercut or overtopped the ridge axis, delivering warm marine air to the West Coast without polar air support.[2][11] Subsiding air beneath the ridge also established a thermal "warm lid" with fewer cold troughs, suppressing nocturnal valley inversions.[12] With overnight minimums frequently remaining above 0 deg C, the snowpack failed to refreeze at night and stayed isothermal at 0 deg C, leaving the cryosphere vulnerable to rapid ablation from solar radiation and sensible heat.[2]

Thermodynamics of Ablation: Sublimation and Albedo Feedbacks

Snowpack depletion was further accelerated by two microphysical thermodynamic processes: diffusive sublimation and the snow-albedo feedback (SAF).[13][17] Governed by the Clausius-Clapeyron relation, warm anticyclonic air masses exhibited an elevated vapor pressure deficit (VPD), driving water molecules directly from the ice lattice into the vapor phase.[14][15] Enhanced by turbulent wind scouring, field campaigns such as SAIL show that sublimation can remove 10% to 90% of seasonal snowfall in arid Intermountain basins.[15][16]

Concurrently, warm spells degraded surface albedo through three physical triggers: (1) thin snow cover exposing dark ground early; (2) rain-on-snow events transferring sensible and latent heat that coarsened ice grains and lowered reflectivity from ~0.85 to ~0.60; and (3) extended dry spells preventing fresh snow from resetting surface reflectivity as dust and black carbon concentrated on the pack.[18][19][20]

| Energy Flux Component | | Historical Cold Winter | | 2026 Warm Snow Drought | | Thermodynamic Impact |

| Net Shortwave Radiation | | Low (Albedo reflects ~85%) | | High (Albedo reflects ~60%) | | Amplified solar melt input |

| Net Longwave Radiation | | Net nocturnal cooling loss | | Neutral / Net gain | | Loss of nocturnal refreezing |

| Sensible & Latent Heat | | Negative / Low exchange | | Positive / High VPD | | Conductive melt & sublimation |

| Precipitation Heat Flux | | Negligible (Snow 0 deg C) | | High (Rain > 2 deg C) | | Isothermal ripening & scouring |

Regional Hydro-Climatological Assessment and Reservoir Operations

Hydrological impacts across the West reflected regional interactions between storm latitude and antecedent storage:[2][27]

* Pacific Northwest: December 2025 atmospheric rivers delivered over 200% of normal rainfall (>20 inches liquid in mountain basins), scouring early snowpack and driving Oregon Cascade SWE below the 20th percentile.[2][4][21]

* California and the Sierra Nevada: California's hydrology stood at 59% of average SWE in early February 2026 (Phillips Station at 46%; Northern Sierra at 44%; Southern Sierra at 77%).[22][23] Carryover from prior wet years buffered Lake Oroville (127% of average) and Shasta Lake (122%), while adaptive "first flush" operations captured 15,000 acre-feet of winter flow.[24][25][26]

* Colorado River Basin: Upper Basin SWE languished at 55-70% of median (<25% in Arizona).[2][27] Lake Powell (26% full, projected near 3,535 feet) and Lake Mead (33% full) remain locked in the 7.48-million-acre-foot Mid-Elevation Release Tier amid urgent post-2026 negotiations.[28][29][30][31][32]

| Region / Reservoir | | Early 2026 Status | | Historical Context | | Dominant Driver |

| Pacific Northwest (OR/WA) | | 30-40% Median SWE | | < 20th Percentile | | Subtropical AR rain-on-snow scouring [2][21] |

| Sierra Nevada (CA) | | 59% Average SWE | | Phillips Station at 46% | | High freezing levels & dry Jan ridging [22][23] |

| Upper Colorado & Great Basin | | 50-65% Median SWE | | Record low Jan cover | | South-ridge blocking & sublimation [5][27] |

| Lake Oroville & Shasta (CA) | | ~70-80% Capacity | | 122-127% Hist. Avg | | Carryover & adaptive winter capture [24][26] |

| Lake Powell & Mead (AZ/UT/NV) | | 26% & 33% Capacity | | 48% & 60% Hist. Avg | | Structural deficit & soil moisture tax [28][32] |

Runoff Efficiency Penalties and the 2026 Wildfire Outlook

The translation of mountain snow deficits into spring streamflow is compounded by a severe penalty in basin runoff efficiency -- the ratio of snowmelt that reaches river channels versus water lost to soil infiltration and atmospheric evapotranspiration.[13][28] Following an anomalously hot, dry autumn in 2025, root-zone soil moisture profiles across the Rocky Mountains entered winter deeply depleted.[28] When the diminished 2026 snowpack melted in spring, parched soils exacted a heavy "soil moisture tax," absorbing meltwater before channelized runoff could materialize -- a nonlinear dynamic in which a 70% snowpack frequently yields only 50% of normal streamflow. Simultaneously, warmer late-winter temperatures broke vegetation dormancy early, accelerating transpirational water uptake and prompting the Colorado Basin River Forecast Center (CBRFC) to project April-July 2026 runoff at just 55-80% of average across Utah and Colorado (with only the Bear River basin approaching 90%).[8][13][33] Patchy high-altitude snow cover also challenges operational forecasting models such as California's Bulletin 120 (Snow-17), increasing reliance on Airborne Snow Observatory lidar surveys and forcing agricultural districts with junior water rights to expand groundwater pumping.[28][34][27][35]

Across the Western ecosystem, early snow disappearance removed a vital natural suppressant for the 2026 wildfire season while severely contracting winter ski and recreation economies across Arizona, California, Oregon, and Colorado.[2][21] While December rains temporarily moistened surface soils in the Pacific Northwest, they stimulated rapid growth of fine herbaceous fuels (grasses) at lower elevations that cured quickly under spring insolation, while leaving high-country timber and coarse woody fuels exposed to desiccation months ahead of schedule -- widening the hazardous "spring gap" prior to summer monsoons.[21]

Seasonal Outcome (April-November 2026): As the 2026 water year unfolded, record-warm

March heatwaves accelerated early melt, leaving Upper Colorado, Utah, and Colorado snowpack near historic lows (~20-25% of average in key basins) by April 1, 2026.[27][29] Severely reduced spring inflows pushed Lake Powell down to roughly 3,516 feet elevation (~21.8% capacity) in mid-September 2026 -- prompting emergency Bureau of Reclamation release restrictions and upstream drought-response transfers from Flaming Gorge to safeguard the 3,490-foot minimum power pool -- while Lake Mead fell into the 1,037-1,039-foot range (~26% capacity) by autumn 2026 amid mandatory Lower Basin delivery cuts.[29][32]

Conclusion: Structural Adaptation to a Rain-Dominated Cordillera

Water Year 2026 confirms the structural shift of the American West from a snow-dominated to a volatile rain-dominated regime. The convergence of a non-canonical La Ni a, persistent ridging, high VPD sublimation, and snow-albedo feedbacks stripped mountain storage even during active atmospheric river transport.[2][11][18] Adapting to recurring warm snow droughts requires pairing demand conservation with Forecast-Informed Reservoir Operations (FIRO), airborne snow telemetry, and conjunctive aquifer management.[26][28]

Updates (Nov 20, 2026): Added seasonal outcome through November 2026 documenting April 1 Upper Colorado Basin SWE lows, Lake Powell's mid-September drop to ~3,516 ft and emergency Flaming Gorge transfers, and Lake Mead's fall 2026 decline to ~1,037-1,039 ft; replaced blocked Smithsonian, AMS XML, ResearchGate, California DWR, and USDA legacy URLs with verified primary DOIs (including Pye & Pu 2025, Gibson et al. 2020, Lundquist et al. 2024, Thackeray & Fletcher 2016, Kaufmann & Pretis 2023, and Painter et al. 2016), NOAA Climate.gov, CDEC, and USDA NWCC sources; unnumbered headings, added publication dates across all 35 references, and enforced 1:1 inline parity.

References & Primary Sources

* [1] NIDIS / Drought.gov (2026). Snow Drought. -- https://www.drought.gov/what-is-drought/snow-drought

* [2] NIDIS / Drought.gov (Jan 8, 2026). Snow Drought Current Conditions and Impacts in the West. -- https://www.drought.gov/drought-status-updates/snow-drought-current-conditions-and-impacts-west-2026-01-08

* [3] Harpold, A. A., Dettinger, M., & Rajagopal, S. (Feb 28, 2017). Defining Snow Drought and Why It Matters. Eos. -- https://eos.org/opinions/defining-snow-drought-and-why-it-matters

* [4] NOAA Climate.gov (2026). Snow drought across the western United States. -- https://www.climate.gov/news-features/event-tracker/snow-drought-across-western-united-states

* [5] NASA Earth Observatory (Jan 29, 2026). The West Faces Snow Drought. -- https://science.nasa.gov/earth/earth-observatory/the-west-faces-snow-drought/

* [6] Harpold, A. A., Dettinger, M., & Rajagopal, S. (2017). Defining Snow Drought and Why It Matters. University of Nevada, Reno Extension (SP-17-01). -- https://naes.agnt.unr.edu/PMS/Pubs/687_2017_33.pdf

* [7] NOAA Climate Prediction Center (2026). ENSO Diagnostic Discussion. --

https://www.cpc.ncep.noaa.gov/products/analysis_monitoring/enso_advisory/ensodisc.shtml

* [8] NOAA Colorado Basin River Forecast Center (Jan 2026). January 2026 Water Supply Briefing. -- https://www.cbrfc.noaa.gov/present/2026/cbrfcwsupjan2026.pdf

* [9] NOAA Climate Prediction Center (2026). ENSO: Recent Evolution, Current Status and Predictions. -- https://www.cpc.ncep.noaa.gov/products/analysis_monitoring/lanina/enso_evolution-status-fcsts-web.pdf

* [10] Pye, M. J., & Pu, Z. (Aug 2025). The Synoptic-Scale Circulation during the Western U.S. Drought of 2021 and 2022. Journal of Applied Meteorology and Climatology, 64(8). -- https://doi.org/10.1175/jamc-d-24-0059.1

* [11] Gibson, P. B., Waliser, D. E., Guan, B., DeFlorio, M. J., Ralph, F. M., & Swain, D. L. (Apr 1, 2020). Ridging Associated with Drought across the Western and Southwestern United States: Characteristics, Trends, and Predictability Sources. Journal of Climate, 33(7), 2485-2508. -- https://doi.org/10.1175/jcli-d-19-0439.1

* [12] Teng, H., Leung, R., et al. (2022). Fewer Troughs, Not More Ridges, Have Led to a Drying Trend in the Western United States. OSTI / Pacific Northwest National Laboratory. -- https://www.osti.gov/pages/servlets/purl/1978525

* [13] NIDIS / Drought.gov (May 20, 2025). Special Snow Drought Update: Rapid Snowmelt. -- https://www.drought.gov/drought-status-updates/special-snow-drought-update-rapid-snowmelt-2025-05-20

* [14] Jambon-Puillet, E., Shahidzadeh, N., & Bonn, D. (Oct 10, 2018). Singular sublimation of ice and snow crystals. Nature Communications, 9, 4191. -- https://doi.org/10.1038/s41467-018-06689-x

* [15] Lundquist, J. D., Vano, J., Gutmann, E., et al. (Jun 2024). Sublimation of Snow. Bulletin of the American Meteorological Society, 105(6). -- https://doi.org/10.1175/bams-d-23-0191.1

* [16] Aspen Global Change Institute (2023). Sublimation of Snow Resources (SOS Project). -- https://www.agci.org/projects/sublimation-of-snow-sos-project/sublimation-of-snow-resources

* [17] Thackeray, C. W., & Fletcher, C. G. (Jan 6, 2016). Snow albedo feedback. Progress in Physical Geography: Earth and Environment, 40(3), 392-408. -- https://doi.org/10.1177/0309133315620999

* [18] Rupper, S., et al. (Apr 30, 2020). An Examination of Physical Processes That Trigger the Albedo-Feedback on Glacier Surfaces and Implications for Regional Glacier Mass Balance Across High Mountain Asia. Frontiers in Earth Science, 8, 129. -- https://www.frontiersin.org/journals/earth-science/articles/10.3389/feart.2020.00129/full

* [19] Letcher, T. W., & Minder, J. R. (Sep 29, 2015). Characterization of the Simulated Regional Snow Albedo Feedback Using a Regional Climate Model over Complex Terrain. Journal of Climate, 28(19), 7576-7595. -- https://doi.org/10.1175/jcli-d-15-0166.1

* [20] Kaufmann, R. K., & Pretis, F. (Jul 27, 2023). An empirical estimate for the snow albedo feedback effect. Climatic Change, 176, 108. --

https://doi.org/10.1007/s10584-023-03572-7

* [21] National Interagency Fire Center (Feb 2, 2026). National Significant Wildland Fire Potential Outlook (January through April 2026). -- https://www.nifc.gov/nicc-files/predictive/outlooks/monthly_seasonal_outlook.pdf

* [22] California Department of Water Resources / CDEC (2026). Statewide Snow Water Content (SWE) Current Conditions. -- https://cdec.water.ca.gov/snowapp/sweq.action

* [23] USDA NRCS National Water and Climate Center (2026). Snow and Climate Monitoring Portal. -- https://nwcc-apps.sc.egov.usda.gov/

* [24] Medina, M. / SFGATE (Dec 30, 2025). Where California's reservoir levels stand going into 2026. -- https://www.sfgate.com/bayarea/article/where-california-reservoir-levels-stand-going-2026-21268544.php

* [25] Engaging Data (2026). Interactive California Reservoir Levels Dashboard. -- https://engaging-data.com/ca-reservoir-dashboard/

* [26] California Natural Resources Agency (Jan 2026). December Storms, Improved Flexibility Allow DWR to Increase State Water Project Allocation. -- https://content.govdelivery.com/accounts/CNRA/bulletins/406bd33

* [27] USDA NRCS National Water and Climate Center (2026). Western U.S. SNOTEL Snow/Precipitation Monitoring. -- https://nwcc-apps.sc.egov.usda.gov/

* [28] NIDIS / Drought.gov (Jan 15, 2026). Drought Status Update for the Intermountain West. -- https://www.drought.gov/drought-status-updates/drought-status-update-intermountain-west-2026-01-15

* [29] U.S. Bureau of Reclamation (2026). Glen Canyon Dam Water Operations. -- https://www.usbr.gov/uc/water/crsp/cs/gcd.html

* [30] Newsweek (Aug 20, 2025). Lake Mead, Lake Powell Water Level Update as Warning Issued. -- https://www.newsweek.com/lake-mead-lake-powell-water-level-update-warning-issued-2116365

* [31] Pagosa Daily Post (Feb 2, 2026). Colorado River Governors Cautiously Optimistic Following 'Historic' DC Meeting. -- https://pagosadailypost.com/2026/02/02/colorado-river-governors-cautiously-optimistic-following-historic-dc-meeting/

* [32] U.S. Bureau of Reclamation (Jan 2026). Most Probable 24-Month Study: January 2026. -- https://www.usbr.gov/lc/region/g4000/24mo.pdf

* [33] Western Water Assessment (2026). Intermountain West Dashboard. -- https://wwa.colorado.edu/resources/intermountain-west-dashboard

* [34] Painter, T. H., Berisford, D. F., Boardman, J. W., et al. (Oct 2016). The Airborne Snow Observatory: Fusion of scanning lidar, imaging spectrometer, and physically-based modeling for mapping snow water equivalent and snow albedo. Remote Sensing of Environment, 184, 139-152. -- https://doi.org/10.1016/j.rse.2016.06.018

* [35] U.S. Bureau of Reclamation (2026). Northern California Area Office Daily Operational Data. -- https://www.usbr.gov/mp/cvo/vungvari/daily.pdf