Ecology & Evolution | Feb 17, 2026 | 8 min read | By Bryan White

# Titans of the Tropics: Unearthing Costa Rica's Ice Age Giants in the Orosi Vally

Biology * Evolution * Genetics

## The Isthmus as a Biological Crucible

The unearthing of a major Late Pleistocene fossil assemblage in the Orosi Valley of Cartago, Costa Rica -- announced in February 2026 by the National Museum of Costa Rica (MNCR) -- marks a watershed moment in Central American paleontology.[4] Excavated from a stratigraphically distinct fluvial horizon influenced by volcanic sedimentation, the locality has yielded 49 exceptionally preserved skeletal elements belonging to two emblematic megafaunal taxa: the spiral-tusked gomphothere Cuvieronius (colloquially designated "Pital" / "Pit n") and the pan-American giant ground sloth Eremotherium ("Tobby" / "Tobi").[4][6] Preliminary geochronological assessments bracket the assemblage between 10,000 and 40,000 years before present (BP) during the Late Rancholabrean North American Land Mammal Age, offering a rare window into the geology, comparative anatomy, and paleoecology of terminal Pleistocene Central America.[4][5]

Driven by plate tectonics and volcanic arc uplift, the emergence of the Central American isthmus roughly 2.7 million years ago fundamentally reorganized global ocean circulation and terrestrial biodiversity.[1] Prior to final closure of the Isthmus of Panama, South America evolved in Neotropical isolation, producing endemic xenarthrans, notoungulates, and litopterns, while North America hosted Nearctic fauna derived from Holarctic lineages.[1] The resulting land bridge triggered the Great American Biotic Interchange (GABI), a massive bidirectional migration across the Western Hemisphere.[2]

Positioned at the narrowest bottleneck of this land bridge, Costa Rica functioned not merely as a passive transit corridor, but as a selective ecological filter and evolutionary holding pen where rugged montane and lowland habitats shaped migrating populations.[1] Because acidic tropical soils, high precipitation, and rapid organic decay typically destroy vertebrate remains, well-preserved Pleistocene localities in Central America are exceedingly scarce. The associated recovery of a northern-origin proboscidean and a southern-origin giant xenarthran in Cartago resolves critical gaps in the regional fossil record.[3][4]

## Discovery, Excavation, and Volcanic Taphonomy in the Orosi Valley

The Cartago locality emerged through two sequential discoveries on private agricultural land in the Orosi Valley, a depression carved by the Reventaz n River and flanked by the Iraz and Turrialba volcanoes of the Central Volcanic Cordillera.[5] In December 2024, geologist Mar a Sequeira Castro identified dense, oversized skeletal elements eroding from a riverbank cut during a routine land-use survey, which the MNCR confirmed as the giant ground sloth Eremotherium.[5] In December 2025, farmworker Esteban Brenes located a second concentration of fossils 700 meters away.[5] Because the cylindrical bones were heavily darkened and

mineralized, surveyors initially mistook them for petrified tree trunks buried by ancient volcanic ash falls -- until microscopic examination revealed cancellous trabecular bone and proboscidean dentition belonging to a nearly complete Cuvieronius skeleton.[5]

Under the direction of MNCR geologist Joanna M ndez Herrera, a 12-person multidisciplinary team of geologists, archaeologists, biologists, and University of Costa Rica (UCR) interns -- advised by paleontologist Dr. Spencer G. Lucas (New Mexico Museum of Natural History and Science) and volcanologist Dr. Guillermo Alvarado Induni -- completed 13 rescue excavation campaigns.[4] Using spatial grid mapping, field consolidants, and plaster jacketing, the team extracted 49 fossil elements, naming the gomphothere "Pital" and the ground sloth "Tobby" to foster public engagement.[4][6]

Stratigraphic analysis indicates primary burial within a low-energy floodplain or backwater oxbow setting.[5] Whereas high-energy river currents disarticulate skeletons and winnow away lighter elements, the preservation of associated vertebrae, ribs, and cranial elements in "Pital" points to rapid, quiet burial via floodplain inundation, miring, or a volcaniclastic lahar.[5] Subsequent preservation was governed by permineralization via silicification: groundwater enriched in dissolved silica from weathering tephra and volcanic glass (including obsidian and ash) percolated through the porous bone matrix, precipitating opal and chalcedony within the trabecular voids.[5] While this silica replacement shielded the bones from acidic tropical dissolution, it degrades endogenous collagen, prompting the team to seek associated carbonized wood for high-precision radiocarbon (14C) radiometric dating within the 10,000-40,000 BP stratigraphic window.[5]

| Recovered Element | | Cuvieronius ("Pital") | | Eremotherium ("Tobby") | | Paleontological & Biomechanical Value |

| Upper Tusk (Defensa) | | 1 Complete (1.60 m), 1 Fragment[4] | | N/A | | Taxonomic diagnosis via spiral enamel band; stable isotope diet profiling.[7] |

| Femur | | Recovered | | Recovered[4] | | Body mass estimation; pedolateral stance and bipedal rearing mechanics.[6] |

| Vertebrae & Ribs | | Recovered (Near-complete axial skeleton)[5] | | Recovered | | Confirmation of low-energy primary burial without fluvial dispersal.[5] |

| Phalanges (Unguals) | | Recovered | | Recovered[4] | | Claw morphology, substrate interaction, and branch-pulling mechanics. |

## Paleobiology of the Megafauna: Cuvieronius ("Pital") and Eremotherium ("Tobby")

The Orosi Valley specimens represent two distinct evolutionary solutions to megaherbivory in Neotropical montane environments:

* Cuvieronius ("Pital"): Unlike the grazing woolly mammoth (Mammuthus primigenius) or American mastodon (Mammut americanum), Cuvieronius belonged to the Gomphotheriidae, an ancient proboscidean family that colonized Central and South America.[7] Standing 2.7 to 3.0 meters at the shoulder and weighing 5 to 6 metric tons, Cuvieronius is diagnosed by its elongated upper tusks -- exemplified by

Pital's intact 1.60-meter tusk[4] -- which exhibit a characteristic helical twist and retain a primitive spiral band of enamel.[7] Stable carbon isotope ( 13C) analyses of Costa Rican Cuvieronius hyodon tooth enamel distinguish browsing on trees and shrubs utilizing C3 carbon fixation from grazing on tropical grasses utilizing C4 carbon fixation.[8][9] These isotopic profiles confirm that Costa Rican populations were adaptable mixed-feeders with a strong preference for C3 forest and woodland browse, using their spiral tusks to strip bark and unearth roots across intermontane valleys.[9]

* Eremotherium ("Tobby"): Representing the pinnacle of ground sloth evolution within the family Megatheriidae, the Pan-American giant ground sloth (Eremotherium laurillardi) reached lengths of 6 meters and masses of 3 to 5 metric tons -- vastly dwarfing extant arboreal relatives such as the two-toed sloth.[10] The wide, anteroposteriorly flattened femur and curved ungual phalanges recovered at Cartago reflect obligatory pedolateral locomotion, in which the animal walked on the outer margins of its feet to accommodate its massive defensive and foraging claws.[6] Supported by its muscular tail in a tripodal stance, Eremotherium reared bipedally to browse foliage 4 to 5 meters high, acting as a keystone ecosystem engineer that maintained open canopy mosaics.[10][11]

## Biogeography of the Great American Biotic Interchange and Late Pleistocene Extinction

Finding Cuvieronius and Eremotherium within the same stratigraphic horizon in Cartago encapsulates the bidirectional dynamics of the Great American Biotic Interchange across the Central American biosphere.[1] Cuvieronius exemplifies the southward Holarctic wave: evolving in North America from Eurasian proboscidean ancestors, it traversed the isthmus and colonized the South American Andes.[3][12] Conversely, Eremotherium exemplifies the northward Neotropical wave: originating in South America's endemic xenarthran radiation, it expanded through Central America as far north as the coastal plains of the southern United States.[13]

Furthermore, the co-occurrence of two browsing and mixed-feeding megaherbivores in the Orosi Valley informs the long-standing paleoecological debate over whether glacial Central America formed a continuous dry savanna corridor or remained forested.[1] The prevalence of C3-browsing Cuvieronius and Eremotherium -- alongside the scarcity of obligate C4 grazers in Costa Rican montane basins -- supports a heterogeneous forest-parkland mosaic that selectively filtered migrating megafauna.[1][11]

Finally, the 10,000-40,000 BP age range situates "Pital" and "Tobby" near the Pleistocene-Holocene transition, when climate-driven expansion of closed-canopy wet forests and the arrival of Paleoindian hunters (evidenced in Costa Rica by Clovis-like fluted projectile points at sites such as Finca Guardiria) culminated in the terminal megafaunal extinction.[14] Although the articulated completeness of the Orosi gomphothere and absence of lithic cut marks or marrow fractures point to natural mortality rather than human butchery, refined radiometric dating will clarify whether these individuals overlapped directly with early human foragers.[5]

## A New Chapter for Central American Paleontology

The discovery of the Orosi Valley megafaunal assemblage overturns long-held assumptions that humid, volcanically active tropical environments cannot preserve complete Pleistocene skeletons.[4] By preserving both a northern gomphothere ("Pital") and a southern giant ground sloth ("Tobby") in silica-mineralized primary context, the Cartago locality provides an essential benchmark for reconstructing the Great American Biotic Interchange.[4] As conservation and isotopic analyses proceed at the National Museum of Costa Rica, these Pleistocene giants underscore how rapidly shifting climate regimes and habitat restructuring can drive even the most adaptable megafauna to extinction -- a paleobiological lesson with enduring relevance for tropical ecosystem resilience in the Anthropocene.

Updates... (Nov 21, 2026): Corrected the name of advising paleontologist Dr. Spencer G. Lucas (previously inverted as "Dr. Lucas Spencer") and clarified Spanish museum nicknames ("Pit n" / "Tobi"), replaced Wikipedia citations with peer-reviewed studies (O'Dea et al., 2016; Cartelle & De Iuliis, 1995), added exact publication dates across references, and aligned inline citations 1:1 with the reference list.

## References & Primary Sources

* [1] Woodburne, M. O. (Dec 2010). The Great American Biotic Interchange: Dispersals, Tectonics, Climate, Sea Level and Holding Pens. Journal of Mammalian Evolution, 17, 245-264. -- https://doi.org/10.1007/s10914-010-9144-8

* [2] O'Dea, A., Lessios, H. A., Coates, A. G., Eytan, R. I., et al. (Aug 5, 2016). Formation of the Isthmus of Panama. Science Advances, 2(8), e1600883. -- https://doi.org/10.1126/sciadv.1600883

* [3] Lucas, S. G., & Alvarado, G. E. (Feb 3, 2010). Fossil proboscidea from the upper Cenozoic of Central America: taxonomy, evolutionary and paleobiogeographic significance. Revista Geol gica de Am rica Central, 42, 9-42. -- https://doi.org/10.15517/rgac.v0i42.4169

* [4] De ltimo Minuto (Feb 13, 2026). Fossils of a giant sloth and a mastodon from at least 10,000 years ago found in Costa Rica. -- https://deultimominuto.net/en/uncategorized/fossils-of-a-giant-sloth-and-a-mastodon-from-at-least-10000-years-ago-found-in-costa-r

* [5] CRHoy (Feb 16, 2026). As fue como 3 ge logas encontraron los f siles del perezoso gigante en Cartago. -- https://crhoy.com/nacionales/fotos-asi-fue-como-3-geologas-encontraron-los-fosiles-del-perezoso-gigante-en-cartago/

* [6] Teletica (Feb 17, 2026). Vea aqu qu nombre le pusieron a los f siles hallados en Cartago. -- https://www.teletica.com/nacional/vea-aqui-que-nombre-le-pusieron-a-los-fosiles-hallados-en-cartago_402395

* [7] Houde, P., Lucas, S. G., & Heizler, M. (Feb 3, 2026). First Known Cranium of Cuvieronius (Proboscidea: Gomphotheriidae) from North America. Diversity, 18(2), 92. -- https://doi.org/10.3390/d18020092

* [8] Laurito, C. A., Valerio, A. L., & Rojas-Sibaja, N. (Dec 18, 2016). El mastodonte bajo el agua: el registro de Cuvieronius hyodon Fischer (1814) en la plataforma continental interna del Pac fico de Costa Rica. Revista Geol gica de Am

rica Central, 55. -- https://doi.org/10.15517/rgac.v55i0.27069

* [9] Adri n P rez-Crespo, V., Laurito, C., Arroyo-Cabrales, J., et al. (Apr 2, 2024). Feeding habits of the Gomphothere Cuvieronius hyodon in Costa Rica: A biochemical approach. Historical Biology. -- https://doi.org/10.1080/08912963.2023.2183856

* [10] Cartelle, C., & De Iuliis, G. (Dec 27, 1995). Eremotherium laurillardi: The Panamerican Late Pleistocene megatheriid sloth. Journal of Vertebrate Paleontology, 15(4), 830-841. -- https://doi.org/10.1080/02724634.1995.10011265

* [11] McDonald, H. G. (Jun 6, 2023). A Tale of Two Continents (and a Few Islands): Ecology and Distribution of Late Pleistocene Sloths. Land, 12(6), 1192. -- https://doi.org/10.3390/land12061192

* [12] Alberdi, M. T., & Prado, J. L. (Aug 3, 2022). Diversity of the fossil gomphotheres from South America. Historical Biology. -- https://doi.org/10.1080/08912963.2022.2067754

* [13] Morgan, G. S. (2005). The Great American Biotic Interchange in Florida. Bulletin of the Florida Museum of Natural History, 45(4), 271-312. -- https://flmnhbulletin.com/index.php/flmnh/article/download/flmnh-vol45-no4-pp271-312/vol45-no4/1156

* [14] La Voz de Guanacaste (Sep 24, 2021). Mastodons in Nicoya? Yes, and other giants also inhabited the area. -- https://vozdeguanacaste.com/en/mastodons-in-nicoya-yes-and-other-giants-also-inhabited-the-area/