Ecology & Evolution | Jan 16, 2026 | 10 min read | By Bryan White

What Does a "Winter Anomaly" Mean for the Future of Whales?

Biology * Conservation * Marine Life

Introduction: The "Urban Whale" and the January 2026 Anomaly

In early January 2026, a remarkable biological event in Cape Cod Bay, Massachusetts, forced marine biologists and oceanographers to revise models of cetacean habitat usage across the warming Northwest Atlantic. On January 10, 2026, aerial observers from the Center for Coastal Studies (CCS) documented 33 North Atlantic right whales (Eubalaena glacialis) engaged in subsurface feeding.[1] Representing nearly 9% of the then-estimated global population of 384 individuals,[2] this single-day sighting set a January density record and accounted for over 60% of the 54 unique whales identified in the bay since November 2025.[1][8]

Known as the "urban whale" because its migratory corridor runs through industrialized shipping lanes and commercial fishing grounds, Eubalaena glacialis teeters on the brink of functional extinction.[12] Although the October 2025 population estimate of 384 reflected a 2.1% increase over 376 the prior year, fewer than 75 reproductively active females remain in the population -- and the Consortium's November 2026 report card subsequently placed the updated 2025 estimate at 382 (+/-11) individuals (73 reproductive females), a 1.8% decline from the recalculated 2024 baseline of 389.[3][4][5] Compounding vessel strikes and entanglements is rapid climate change: the Gulf of Maine has warmed faster than 99% of the global ocean, decoupling seasonal prey cycles.[6][7] The January 2026 aggregation -- including the 45-year-old male EgNo 1050 and the reproductive female Monarch (Catalog #2460) -- illustrates how biophysical prey trapping and ocean regime shifts are compressing right whales into coastal refugia months ahead of historical schedules.[1]

The January 10, 2026 Sighting: Behavioral Shift and Demographics

Historically, right whale abundance in Cape Cod Bay peaks in March and April during the spring zooplankton bloom, while sparse December and January sightings typically involve Surface Active Groups (SAGs) engaged in courtship.[1][9][10] On January 10, 2026, however, CCS observer Ryan Schosberg documented 33 whales executing "shallow subsurface feeding" -- swimming with mouths agape just beneath the surface to filter concentrated strata of planktonic prey.[1]

Because filter-feeding generates substantial hydrodynamic drag, right whales only forage when prey patches exceed a bioenergetic threshold of ~3,000 to 4,000 copepods per cubic meter.[11] The shift from winter social behavior to intensive foraging signals both an anomalously early zooplankton bloom in Cape Cod Bay and caloric urgency following poor summer foraging in northern basins.[7] Photographic catalog matching linked the aggregation to crucial demographic histories:

* EgNo 1050 (Adult Male, >45 Years): Where anthropogenic trauma has depressed

average female longevity to ~45 years and male longevity to ~65 years, EgNo 1050 is a rare survivor.[1][12] Documented in Cape Cod Bay only once in the prior 15 years, his presence indicates that strong trophic cues -- or barren winter alternatives -- drew sporadic visitors into the bay.[1]

* Monarch (Catalog #2460): First cataloged in 1994, Monarch is a vital matriarch whose 2004 daughter, Platypus (#3420), also calved in 2025.[13] Having birthed a calf in 2025 -- first sighted in Cape Cod Bay rather than off Florida -- Monarch used the January 2026 bloom for post-lactation recovery.[1][13] Although 11 mother-calf pairs were documented in early January 2026 (against a 20+ annual birth benchmark), the full 2025-2026 calving season ultimately reached 24 calves -- the highest annual total since 2009.[4][14]

| Metric / Individual | | January 10, 2026 Observation | | Demographic & Ecological Implication |

| Single-Day Aggregation | | 33 unique whales (54 season-to-date) | | ~8.6% of global population (384 at Jan. 2026; revised 382 in Nov. 2026) in one bay. |

| Foraging Behavior | | Shallow subsurface skim-feeding | | Confirms winter copepod density >3,000-4,000 m 3. |

| EgNo 1050 | | Adult male (>45 years old) | | Survival of older cohort; recruitment of sporadic winter foragers. |

| Monarch (#2460) | | Reproductive female (1994 cohort) | | Post-lactation caloric recovery supporting multi-generational matrilines. |

The Ecological Engine: Calanus finmarchicus and Ocean Regime Shifts

Right whale movements track their obligate prey, the calanoid copepod Calanus finmarchicus.[15] During its late-juvenile C5 stage, C. finmarchicus accumulates wax-ester lipids -- up to 70% of dry mass -- before descending into deep waters for seasonal dormancy (diapause).[15][16] Whales rely on these lipid-dense swarms to build blubber for insulation, migration, and reproduction.

Cape Cod Bay acts as a hydrodynamic retention basin governed by physical oceanography and circulation.[17] In winter, the Gulf of Maine Coastal Current advects C. finmarchicus from Wilkinson Basin into the bay, where a counter-clockwise gyre retains the plankton.[18][19] As surface waters stratify, copepods concentrate along the thermocline.[19][20] Right whales detect these layers via low-light visual contrast and tactile feedback from chin vibrissae and baleen nerves.[11][15]

This early winter aggregation stems from a 2010 shift in thermohaline circulation.[7] Reduced cold Labrador Current inflow and increased Gulf Stream Warm Slope Water elevated eastern Gulf of Maine temperatures, collapsing summer C. finmarchicus stocks in the Bay of Fundy.[7][21] Tracking prey across the food web, whales shifted summer foraging to the Gulf of St. Lawrence -- triggering the 2017 Unusual Mortality Event -- while arriving earlier each winter in Cape Cod Bay.[6][21]

| Habitat | | Historic Peak (Pre-2010) | | Current Phenology (2020s) | | Oceanographic & Trophic Driver |

| Cape Cod Bay | | March - April | | January - May | | Early stratification and gyre retention of C. finmarchicus. |

| Bay of Fundy | | July - October | | Largely abandoned | | Warm Slope Water incursion collapsing summer C5 stocks. |

| Gulf of St. Lawrence | | Sporadic / Low | | May - November | | Northern refuge foraging; required transboundary speed/gear rules. |

| Southeast U.S. Coast | | December - March | | December - March | | Stable thermal requirement for neonatal calves. |

Anthropogenic Threats, Technological Mitigation, and Policy

Concentrating 33 subsurface-feeding whales in a busy waterway magnifies vessel-strike and entanglement hazards.[2][6] Lacking a dorsal fin and skimming just below the surface, right whales are nearly invisible to mariners, while over 85% bear scars from fixed-gear vertical buoy lines.[2] Although the mandatory 10-knot Seasonal Management Area (SMA) for vessels 65 feet began January 1, 2026,[1][22] the January 10 sighting occurred three weeks before the February 1 Massachusetts Restricted Area (MRA) trap/pot closure, exposing whales to "shoulder-season" lines and vessels <65 feet.[24][28] In March 2026, NOAA Fisheries also issued an Advance Notice of Proposed Rulemaking to evaluate modifications and technology-based alternatives to the federal vessel speed rule.[25]

Closing these protection gaps under the U.S. Endangered Species Act, Marine Mammal Protection Act, and Canadian rules relies on three conservation technologies:[5][25][31]

* Acoustic Surveillance (Robots4Whales): Operated by Woods Hole Oceanographic Institution and NOAA, the Cape Cod Bay DMON buoy detects low-frequency right whale "up-calls" 24/7, relaying satellite alerts to trigger Slow Zones.[23][26]

* AI Photo-Identification: Convolutional neural networks on the Flukebook platform match rostral callosity patterns against the catalog, rapidly identifying individuals like EgNo 1050 and Monarch.[27][28]

* On-Demand (Ropeless) Gear: Trials under NOAA Exempted Fishing Permits (EFPs) stow buoy lines on the seafloor until released by an encrypted acoustic signal, eliminating persistent vertical ropes while using virtual chart-plotter marks to prevent gear conflict.[29][30]

| Feature | | Traditional Trap/Pot Gear | | On-Demand (Ropeless) Systems |

| Surface Marker | | Floating buoy and vertical rope | | Virtual GPS mark on chart plotters |

| Entanglement Risk | | High (vertical line present 24/7) | | Near zero (line present only during haul) |

| Retrieval | | Visual sighting and gaffing | | Acoustic modem trigger + buoyant lift |

| Regulatory Status | | Banned in MRA closure (Feb 1 - May 15) | | Permitted in closures via EFP trials |

Conclusion: The Winter of Decision

The January 10, 2026 aggregation of 33 North Atlantic right whales in Cape Cod Bay highlights both the foraging adaptability and acute vulnerability of Eubalaena glacialis.[1] Even with 24 calves born during the 2025-2026 season, the November 2026 population estimate of 382 individuals confirms that anthropogenic mortality continues to constrain recovery.[4] Translating the survival of EgNo 1050 and the Monarch matriline into species recovery requires aligning vessel speed limits, acoustic monitoring, and ropeless fishing gear with the pace of marine climate change.[21][30]

Updates... (Nov 21, 2026): Updated population and reproductive metrics through the November 2026 North Atlantic Right Whale Consortium report card (382 +/- 11 whales, 73 reproductive females, and 24 calves in the 2025-2026 season) and NOAA's March 2026 vessel speed rule ANPRM; upgraded MEPS prey-threshold and Conservation Biology photo-ID citations to verified CrossRef DOIs; added full publication dates across all 31 primary sources; and split grouped inline citations for one-to-one reference parity.

References

* [1] Center for Coastal Studies (Jan 12, 2026). 33 Right Whales Sighted in Cape Cod Bay. -- https://coastalstudies.org/news/33-right-whales-sighted-in-cape-cod-bay/

* [2] Conservation Law Foundation (Oct 21, 2025). New Population Estimate Reveals Just 384 North Atlantic Right Whales Remain. -- https://www.clf.org/newsroom/new-population-estimate-reveals-just-384-north-atlantic-right-whales-remain/

* [3] New England Aquarium (Oct 21, 2025). North Atlantic right whale population continues slow growth. -- https://www.neaq.org/about-us/press-room/press-releases/north-atlantic-right-whale-population-continues-slow-growth/

* [4] New England Aquarium (Nov 16, 2026). North Atlantic right whale population continues to struggle in latest update on the critically endangered species. -- https://www.neaq.org/about-us/press-room/press-releases/north-atlantic-right-whale-population-continues-to-struggle-in-latest-updat

* [5] Center for Biological Diversity (Mar 21, 2025). Legal Intervention Sought to Protect Endangered North Atlantic Right Whales From Deadly Vessel Strikes. -- https://biologicaldiversity.org/w/news/press-releases/legal-intervention-sought-to-protect-endangered-north-atlantic-right-whales-fro

* [6] Marine Mammal Commission (Feb 1, 2023). Effects of Climate Change on Marine Mammals: North Atlantic Right Whale Distribution Shift. -- https://www.mmc.gov/wp-content/uploads/Climate-Change-Distribution-Shift-Factsheet-2.1.23.pdf

* [7] Meyer-Gutbrod, E. L., Greene, C. H., Davies, K. T. A., & Johns, D. G. (Sep 1, 2021). Ocean Regime Shift is Driving Collapse of the North Atlantic Right Whale Population. Oceanography, 34(3), 22-31. -- https://doi.org/10.5670/oceanog.2021.308

* [8] CapeCod.com (Jan 13, 2026). 33 right whales sighted in Cape Cod Bay, vessel restrictions in effect. -- https://www.capecod.com/newscenter/33-right-whales-sighted-in-cape-cod-bay/

* [9] New England Aquarium (Jun 29, 2023). North Atlantic right whales continue springtime influx into Massachusetts waters. --

https://www.neaq.org/about-us/press-room/press-releases/north-atlantic-right-whales-continue-springtime-influx-into-massachusetts-

* [10] CTV News Atlantic (Jan 13, 2026). 'Leaves us hopeful': North Atlantic right whales spotted in southern New England waters. -- https://www.ctvnews.ca/atlantic/article/leaves-us-hopeful-north-atlantic-right-whales-spotted-in-southern-new-england-waters/

* [11] NOAA Institutional Repository (2004). Northwest Atlantic Ocean Habitats Important to the Conservation of North Atlantic Right Whales (Eubalaena glacialis). -- https://repository.library.noaa.gov/view/noaa/3558/noaa_3558_DS1.pdf

* [12] NOAA Fisheries (Mar 4, 2026; updated Jul 29, 2026). North Atlantic Right Whale Species Profile. -- https://www.fisheries.noaa.gov/species/north-atlantic-right-whale

* [13] Center for Coastal Studies (Apr 18, 2025). Researchers Identify Season's Eleventh North Atlantic Right Whale Mother and Calf. -- https://coastalstudies.org/news/center-for-coastal-studies-researchers-identify-seasons-eleventh-north-atlantic-right-whale-mother-an

* [14] New England Aquarium (Dec 2, 2024). 2024-2025 North Atlantic Right Whale Mother and Calf Pairs. -- https://www.neaq.org/2024-2025-north-atlantic-right-whale-mother-and-calf-pairs/

* [15] Cronin, T. W., Fasick, J. I., Schweikert, L. E., Johnsen, S., Kezmoh, L. J., & Baumgartner, M. F. (Apr 5, 2017). Coping with copepods: do right whales (Eubalaena glacialis) forage visually in dark waters? Philosophical Transactions of the Royal Society B: Biological Sciences, 372(1717), 20160067. -- https://doi.org/10.1098/rstb.2016.0067

* [16] Ross, C. H., Runge, J. A., Roberts, J. J., Brady, D. C., Tupper, B., & Record, N. R. (2023). Estimating North Atlantic right whale prey based on Calanus finmarchicus thresholds. Marine Ecology Progress Series, 703, 1-16. -- https://doi.org/10.3354/meps14204

* [17] Geyer, W. R., Gardner, G. B., Brown, W. S., Irish, J. D., Butman, B., Loder, T. C., & Signell, R. P. (1992). Physical Oceanographic Investigation of Massachusetts and Cape Cod Bays. Massachusetts State Library Archives. -- https://archives.lib.state.ma.us/server/api/core/bitstreams/51db5795-4c82-4b00-9f8c-172530189fad/content

* [18] Jiang, M., Brown, M. W., Turner, J. T., Kenney, R. D., Mayo, C. A., Zhang, Z., & Zhou, M. (Nov 8, 2007). Springtime transport and retention of Calanus finmarchicus in Massachusetts and Cape Cod Bays, USA, and implications for right whale foraging. Marine Ecology Progress Series, 349, 183-197. -- https://doi.org/10.3354/meps07088

* [19] Center for Coastal Studies (2024). Physical Oceanography of Stellwagen Bank and Cape Cod Bay. -- https://coastalstudies.org/connect-learn/stellwagen-bank-national-marine-sanctuary/physical-oceanography/

* [20] MWRA (Apr 1998). Conceptual food web model for Cape Cod Bay, with associated environmental interactions. -- https://www.mwra.com/media/file/conceptual-food-web-model-cape-cod-bay-1998-04

* [21] New England Aquarium (Jun 19, 2023). New research shows climate change impacts on whale habitat use in the warming Gulf of Maine. -- https://www.neaq.org/about-us/press-room/press-releases/new-research-shows-climate-change-impacts-on-whale-habitat-use-in-the-

* [22] Federal Register (Aug 1, 2022). Amendments to the North Atlantic Right Whale Vessel Strike Reduction Rule. -- https://www.federalregister.gov/documents/2022/08/01/2022-16211/amendments-to-the-north-atlantic-right-whale-vessel-strike-redu

* [23] NOAA Fisheries (Dec 30, 2025). Right Whale Slow Zone Extension: Cape Cod Bay, MA. -- https://content.govdelivery.com/accounts/USNOAAFISHERIES/bulletins/40223c8

* [24] NOAA Fisheries (Jun 2020). North Atlantic Right Whale (Eubalaena glacialis) Vessel Speed Rule Assessment. -- https://media.fisheries.noaa.gov/2021-01/FINAL_NARW_Vessel_Speed_Rule_Report_Jun_2020.pdf

* [25] NOAA Fisheries (Mar 4, 2026). North Atlantic Right Whale: Conservation & Management. -- https://www.fisheries.noaa.gov/species/north-atlantic-right-whale/conservation-management

* [26] Woods Hole Oceanographic Institution (WHOI) (2025). Robots4Whales: Autonomous Real-Time Marine Mammal Research and Conservation. -- https://robots4whales.whoi.edu/

* [27] NOAA Fisheries (Oct 27, 2026). Super Intelligence: Right Whale Photo Identification. -- https://www.fisheries.noaa.gov/new-england-mid-atlantic/science-data/artificial-intelligence-right-whale-photo-identification

* [28] Bogucki, R., Cygan, M., Khan, C. B., Klimek, M., Milczek, J. K., & Mucha, M. (Nov 28, 2018). Applying deep learning to right whale photo identification. Conservation Biology, 33(3), 676-684. -- https://doi.org/10.1111/cobi.13226

* [29] NOAA Fisheries (Sep 28, 2026). 2026 Northeast Experimental On-Demand Gear System Testing Beginning February 1. -- https://www.fisheries.noaa.gov/new-england-mid-atlantic/science-data/2026-northeast-experimental-demand-gear-system-testing

* [30] NOAA Fisheries (Apr 1, 2026). Developing Viable On-Demand Gear Systems. -- https://www.fisheries.noaa.gov/new-england-mid-atlantic/marine-mammal-protection/developing-viable-demand-gear-systems

* [31] Fisheries and Oceans Canada (Nov 13, 2026). North Atlantic right whale monitoring and research activities. -- https://www.dfo-mpo.gc.ca/fisheries-peches/commercial-commerciale/atl-arc/narw-bnan/narw-science-eng.html